

# UNIQUENESS OF LIMIT MODELS IN CLASSES WITH AMALGAMATION

RAMI GROSSBERG, MONICA VANDIEREN, AND ANDRÉS VILLAVECES

ABSTRACT. We prove:

**Main Theorem:** *Let  $\mathcal{K}$  be an abstract elementary class and  $\mu > \text{LS}(\mathcal{K})$ . Suppose  $\mathcal{K}$  satisfies the disjoint amalgamation property for models of cardinality  $\mu$ . If  $\mathcal{K}$  is  $\mu$ -Galois-stable, does not have long splitting chains, and satisfies locality of splitting, then any two  $(\mu, \sigma_\ell)$ -limits over  $M$  for  $(\ell \in \{1, 2\})$  are isomorphic over  $M$ .*

This theorem extends results of Shelah from [Sh 394], [Sh 576], [Sh 600], Kolman and Shelah in [KoSh] and Shelah and Villaveces from [ShVi]. Our uniqueness theorem has been used by Grossberg and VanDieren to prove a case of Shelah’s categoricity conjecture for tame abstract elementary classes in [GrVa2]. Our Main Theorem stands in for a beginning of superstability for categorical abstract elementary classes (a subject still quite open).

## 1. INTRODUCTION

We work in the general context of abstract elementary classes (AECs) with the amalgamation property (AP), the disjoint amalgamation property, the joint embedding property (JEP), and Galois-stability at one fixed cardinality  $\mu$  above the Löwenheim-Skolem number. We prove the uniqueness of limit models under superstability-like assumptions of the  $\mu$ -splitting dependence relation.

The basic model theory of abstract elementary classes (fundamental definitions, the role of the AP and the JEP, the existence of “monster models”  $\mathfrak{C}$ , Galois types and the foundational development of stability theory in that context) can be checked in the monograph [Gr2] and the books [Ba], [Sh i]).

---

*Date:* January 2, 2012.

*AMS Subject Classification:* Primary: 03C45, 03C52, 03C75. Secondary: 03C05, 03C55 and 03C95.

The second author was partially sponsored for this work by grant DMS 0801313 of the *National Science Foundation*.

The third author was partially sponsored for this work by grant No. 1101-05-13605 of the *Instituto Colombiano para el Desarrollo de la Ciencia y la Tecnología*, Colciencias.

For the sake of completeness, we include some of the fundamentals of this context here.

In 1977, Shelah, building on the work of Fraïssé and Jónsson, identified a non-elementary context in which a model theoretic analysis could be carried out. Shelah began to study classes of models, together with a partial ordering of the class, which exhibit many of the properties that the models of a first order theory have with respect to the elementary submodel relation. Such classes were named abstract elementary classes. They include classes of models axiomatizable in  $L_{\omega_1, \omega}(\mathbf{Q})$ . Both classification theory and stability theory may be carried out to some extent within these classes. One strong advantage is that there are no a priori compactness assumptions. We reproduce the definition here.

**Definition 1.1.** Let  $\mathcal{K}$  be a class of structures all in the same similarity type  $L(\mathcal{K})$ , and let  $\prec_{\mathcal{K}}$  be a partial order on  $\mathcal{K}$ . The ordered pair  $\langle \mathcal{K}, \prec_{\mathcal{K}} \rangle$  is an *abstract elementary class, AEC for short* iff

A0 (Closure under isomorphism)

- (a) For every  $M \in \mathcal{K}$  and every  $L(\mathcal{K})$ -structure  $N$  if  $M \cong N$  then  $N \in \mathcal{K}$ .
- (b) Let  $N_1, N_2 \in \mathcal{K}$  and  $M_1, M_2 \in \mathcal{K}$  such that there exist  $f_l : N_l \cong M_l$  (for  $l = 1, 2$ ) satisfying  $f_1 \subseteq f_2$  then  $N_1 \prec_{\mathcal{K}} N_2$  implies that  $M_1 \prec_{\mathcal{K}} M_2$ .

A1 For all  $M, N \in \mathcal{K}$  if  $M \prec_{\mathcal{K}} N$  then  $M \subseteq N$ .

A2 Let  $M, N, M^*$  be  $L(\mathcal{K})$ -structures in  $\mathcal{K}$ . If  $M \subseteq N$ ,  $M \prec_{\mathcal{K}} M^*$  and  $N \prec_{\mathcal{K}} M^*$ , then  $M \prec_{\mathcal{K}} N$ .

A3 (Downward Löwenheim-Skolem)  $LS(\mathcal{K})$  is the minimal infinite cardinal  $\geq |L(\mathcal{K})|$  such that for every  $M \in \mathcal{K}$  and for every  $A \subseteq |M|$  there exists  $N \in \mathcal{K}$  such that  $N \prec_{\mathcal{K}} M$ ,  $|N| \geq A$  and  $\|N\| \leq |A| + LS(\mathcal{K})$ .

A4 (Tarski-Vaught Chain)

- (a) For every regular cardinal  $\mu$  and every  $N \in \mathcal{K}$  if  $\{M_i \prec_{\mathcal{K}} N : i < \mu\} \subseteq \mathcal{K}$  is  $\prec_{\mathcal{K}}$ -increasing (i.e.  $i < j \implies M_i \prec_{\mathcal{K}} M_j$ ) then  $\bigcup_{i < \mu} M_i \in \mathcal{K}$  and  $\bigcup_{i < \mu} M_i \prec_{\mathcal{K}} N$ .
- (b) For every regular  $\mu$ , if  $\{M_i : i < \mu\} \subseteq \mathcal{K}$  is  $\prec_{\mathcal{K}}$ -increasing then  $\bigcup_{i < \mu} M_i \in \mathcal{K}$  and  $M_0 \prec_{\mathcal{K}} \bigcup_{i < \mu} M_i$ .

For  $M$  and  $N \in \mathcal{K}$  a monomorphism  $f : M \rightarrow N$  is called an  $\mathcal{K}$ -embedding iff  $f[M] \prec_{\mathcal{K}} N$ . Thus,  $M \prec_{\mathcal{K}} N$  is equivalent to “ $\text{id}_M$  is a  $\mathcal{K}$ -embedding from  $M$  into  $N$ .”

For  $M_0 \prec_{\mathcal{K}} M_1$  and  $N \in \mathcal{K}$ , the formula  $f : M_1 \xrightarrow{M_0} N$  stands for  $f$  is a  $\mathcal{K}$ -embedding such that  $f \upharpoonright M_0 = \text{id}_{M_0}$ .

For a class  $\mathcal{K}$  and a cardinal  $\mu \geq \text{LS}(\mathcal{K})$  let

$$\mathcal{K}_{\mu} := \{M \in \mathcal{K} : \|M\| = \mu\}.$$

In reality, abstract elementary classes were not as approachable as one would hope and much work in non-elementary model theory takes place in contexts which additionally satisfy the amalgamation property:

**Definition 1.2.** Let  $\mu \geq \text{LS}(\mathcal{K})$ . We say that  $\mathcal{K}$  has the  $\mu$ -amalgamation property ( $\mu$ -AP) iff for any  $M_{\ell} \in \mathcal{K}_{\mu}$  (for  $\ell \in \{0, 1, 2\}$ ) such that  $M_0 \prec_{\mathcal{K}} M_1$  and  $M_0 \prec_{\mathcal{K}} M_2$  there are  $N \in \mathcal{K}_{\mu}$  and  $\mathcal{K}$ -embeddings  $f_{\ell} : M_{\ell} \rightarrow N$  such that  $f_{\ell} \upharpoonright M_0 = \text{id}_{M_0}$  for  $\ell = 1, 2$ .

A model  $M_0 \in \mathcal{K}_{\mu}$  satisfying the above requirement is called an *amalgamation base*.

We say that  $\mathcal{K}$  has the *amalgamation property* (AP) iff any triple of models from  $\mathcal{K}_{\geq \text{LS}(\mathcal{K})}$  can be amalgamated.

**Remark 1.3.** (1) Using the isomorphism axioms we can see that  $\mathcal{K}$  has the  $\lambda$ -AP iff for any  $M_{\ell} \in \mathcal{K}_{\lambda}$  (for  $\ell \in \{0, 1, 2\}$ ) such that  $M_0 \prec_{\mathcal{K}} M_{\ell}$  (for  $\ell \in \{1, 2\}$ ) there are  $N \in \mathcal{K}_{\lambda}$  and  $f : M_1 \xrightarrow{M_0} N$  such that  $N \succ_{\mathcal{K}} M_2$ .

(2) Using the axioms of AECs it is not difficult to prove that if  $\mathcal{K}$  has the  $\lambda$ -AP for every  $\lambda \geq \text{LS}(\mathcal{K})$  then  $\mathcal{K}$  has the AP.

The roots of the following fact can be traced back to Jónsson’s 1960 paper [Jo], the present formulation is from [Gr1]:

**Fact 1.4.** Let  $\langle \mathcal{K}, \prec_{\mathcal{K}} \rangle$  be an AEC and  $\lambda \geq \kappa > \text{LS}(\mathcal{K})$  such that  $\mathcal{K}_{< \lambda}$  has the AP and the JEP. Suppose  $M \in \mathcal{K}$ . If  $\lambda^{< \kappa} = \lambda \geq \|M\|$  then there exists  $N \succ M$  of cardinality  $\lambda$  which is  $\kappa$ -model-homogeneous.

Thus if an AEC  $\mathcal{K}$  has AP and JEP, then like in first-order stability theory we may assume that there is a large model-homogeneous  $\mathfrak{C} \in \mathcal{K}$ , that acts like a monster model.

We will refer to the model  $\mathfrak{C}$  as the *monster model*. All models considered will be of size less than  $\|\mathfrak{C}\|$ , and we will find realizations of types we construct inside this monster model. From now on, we assume that the monster model  $\mathfrak{C}$  has been fixed.

Notions of types as sets of formulas, even when the class is described in some infinitary logic, do not behave as nicely as in first-order logic. A replacement was introduced by Shelah in [Sh 394]. In order to avoid confusion between this and the classical, syntactic notion, we will use the terminology in [Gr2] and call this newer, different notion the *Galois type*.

Since in this paper we deal only with AECs with the AP property, the notion of Galois type has a simpler definition than in the general case.

**Definition 1.5** (Galois types). Suppose that  $\mathcal{K}$  has the AP.

- (1) Given  $M \in \mathcal{K}$  consider the action of  $\text{Aut}_M(\mathfrak{C})$  on  $\mathfrak{C}$ , for an element  $a \in |\mathfrak{C}|$  let  $\text{ga-tp}(a/M)$  denote the *Galois type of  $a$  over  $M$*  which is defined as the orbit of  $a$  under  $\text{Aut}_M(\mathfrak{C})$ .
- (2) For  $M \in \mathcal{K}$ , we let

$$\text{ga-S}(M) = \{ \text{ga-tp}(a/M) : a \in |\mathfrak{C}| \}.$$

- (3)  $\mathcal{K}$  is  $\lambda$ -Galois-stable iff

$$N \in \mathcal{K}_\lambda \implies |\text{ga-S}(N)| \leq \lambda.$$

- (4) Given  $p \in \text{ga-S}(M)$  and  $N \in \mathcal{K}$  such that  $N \succ_{\mathcal{K}} M$ , we say that  $p$  is *realized* by  $a \in N$ , iff  $\text{ga-tp}(a/M) = p$ . Just as in the first-order case we will write  $a \models p$  when  $a$  is a realization of  $p$ .

For a more detailed discussion of Galois types, their extensions, restrictions, equivalent forms and generalizations, the reader may consult [Gr2].

While the amalgamation property is useful for dealing with Galois types, in this paper we require a stronger version of AP for one of the steps in the proof of the uniqueness of limit models. Specifically, we use  $\mu$ -disjoint amalgamation over limit models to prove that relatively full towers are limit models (Theorem 4).

**Definition 1.6.** Let  $\mathcal{K}$  be an abstract elementary class.  $\mathcal{K}$  has the  $\mu$ -disjoint amalgamation property ( $\mu$ -DAP) iff for every  $M_\ell \in \mathcal{K}_\mu$  (for  $\ell = 0, 1, 2$ ) such that  $M_0 \prec_{\mathcal{K}} M_\ell$  (for  $\ell = 1, 2$ ) there are  $N \in \mathcal{K}_\mu$  which is a  $\mathcal{K}$ -extension of  $M_2$  and a  $\mathcal{K}$ -embedding  $f : M_1 \xrightarrow[M_0]{} N$  such that  $f[M_1] \cap M_2 = M_0$ .

We say that a class has the *disjoint amalgamation property* iff it has the  $\mu$ -disjoint amalgamation property for every  $\mu \geq \text{LS}(\mathcal{K})$ . We write DAP for short. In this paper we only require that disjoint amalgamation hold for the subclass of all limit models of  $\mathcal{K}_\mu$ .

**1.1. Limit Models.** The next notion to consider is that of a saturated model. In homogeneous abstract elementary classes (see, for example, [GrLe]) where one may study classes of models omitting given sets of types, the existence of a saturated model presents some problems. One solution is to consider models which realize as many types as possible. Such models are called Galois-saturated. More formally, a model  $M$  of size  $\kappa > \text{LS}(\mathcal{K})$  is *Galois-saturated* if it realizes all Galois types over submodels  $N \prec_\mathcal{K} M$  of cardinality  $< \kappa$ . When stability theory has been ported to contexts more general than first order logic, many situations have appeared when Galois-saturated models do not fulfill the main roles that saturated models play in elementary classes.

The main concept of this paper is Shelah's *limit model* which (among other things) serves as a substitute for the role of saturation in stability theory (see [Gr2],[ShVi],[Sh i], etc.) or at least serves as a stepping stone to prove the properties of Galois-saturated models. For example, under the assumption of categoricity with reasonable stability conditions, the uniqueness of Galois-saturated models is not straightforward and is proved by first considering limit models [Sh 394]. In some contexts limit models have been successfully used as “tools” towards finding Galois-saturated models ([KoSh] and [Sh 472]). Furthermore, the notion of limit model refines the notion of saturation; more detailed information is given on the particular way one model is embedded inside another.

Limit models appear in [KoSh] and in [Sh 576] under the name  $(\mu, \alpha)$ -saturated models. In [Sh 600], Shelah calls this notion *brimmed*. Later papers, beginning with Villaveces [ShVi], adopt the name *limit models*. We use the more recent terminology. Before defining limit models, we must introduce their building block, universal extensions.

**Definition 1.7.** (1) Let  $\kappa$  be a cardinal  $\geq \text{LS}(\mathcal{K})$ . We say  $M^* \succ_\mathcal{K} N$  is  $\kappa$ -universal over  $N$  iff for every  $N' \in \mathcal{K}_\kappa$  with  $N \prec_\mathcal{K} N'$  there exists a  $\mathcal{K}$ -embedding  $g : N' \xrightarrow[N]{} M^*$  such that:

$$\begin{array}{ccc}
& N' & \\
\text{id} \uparrow & \searrow g & \\
N & \xrightarrow{\text{id}} & M^*
\end{array}$$

- (2) We say  $M^*$  is *universal over  $N$*  or  $M^*$  is a *universal extension of  $N$*  iff  $M^*$  is  $\|N\|$ -universal over  $N$ .

**Definition 1.8.** [Limit models] Let  $\mu \geq \text{LS}(\mathcal{K})$  and  $\alpha < \mu^+$  a limit ordinal and  $N \in \mathcal{K}_\mu$ . We say that  $M$  is  $(\mu, \alpha)$ -*limit over  $N$*  iff there exists an increasing and continuous chain  $\{M_i \mid i < \alpha\} \subseteq \mathcal{K}_\mu$  such that  $M_0 = N$ ,  $M = \bigcup_{i < \alpha} M_i$ ,  $M_i$  is a proper  $\mathcal{K}$ -submodel of  $M_{i+1}$  and  $M_{i+1}$  is universal over  $M_i$  for all  $i < \alpha$ .

From Theorem 1 we get that for  $\alpha \leq \mu^+$  there always exists a  $(\mu, \alpha)$ -limit model provided  $\mathcal{K}$  has the AP, has no maximal models and is  $\mu$ -Galois-stable. This theorem was stated without proof as Claim 1.16 in [Sh 600], for a proof see [GrVa1] or [Gr1].

**Theorem 1** (Existence). *Let  $\mathcal{K}$  be an AEC without maximal models and suppose it is Galois-stable in  $\mu$ . If  $\mathcal{K}$  has the amalgamation property then for every  $N \in \mathcal{K}_\mu$  there exists  $M^* \succ_{\mathcal{K}} N$ , universal over  $N$  of cardinality  $\mu$ .*

The following theorem partially clarifies the analogy with saturated models:

**Theorem 2.** *Let  $T$  be a stable, complete, first-order theory and let  $\mathcal{K}$  be the elementary class of models of  $T$  with the usual notion of elementary submodel. If  $M$  is a  $(\mu, \delta)$ -limit model for  $\delta$  a limit ordinal with  $\text{cf } \delta \geq \kappa(T)$ , then  $M$  is saturated.*

*Proof.* Use an argument similar to the proof of [Sh e, Theorem III 3.11].  $\dashv$

Thus in elementary classes superstability implies that limit models are saturated, in particular are unique. This raises the following natural question for AECs:

**Question 1.9** (Uniqueness problem). *Let  $\mathcal{K}$  be an AEC,  $\mu \geq \text{LS}(\mathcal{K})$ ,  $\sigma_1, \sigma_2 < \mu^+$ ,  $M \in \mathcal{K}_\mu$  and suppose that  $N_\ell$   $(\mu, \sigma_\ell)$ -limit models over  $M$ . What “reasonable” assumptions on  $\mathcal{K}$  will imply that  $\exists f : N_1 \cong_M N_2$ ?*

Question 1.9 is non-trivial only for the case where  $\text{cf}(\sigma_1) \neq \text{cf}(\sigma_2)$ . Using a back and forth argument one can show that when  $\text{cf}(\sigma_1) = \text{cf}(\sigma_2)$  then we get uniqueness without any assumptions on  $\mathcal{K}$ . More precisely:

**Fact 1.10.** *Let  $\mu \geq \text{LS}(\mathcal{K})$  and  $\sigma < \mu^+$ . If  $M_1$  and  $M_2$  are  $(\mu, \sigma)$ -limits over  $M$ , then there exists an isomorphism  $g : M_1 \rightarrow M_2$  such that  $g \upharpoonright M = \text{id}_M$ . Moreover if  $M_1$  is a  $(\mu, \sigma)$ -limit over  $M_0$ ;  $N_1$  is a  $(\mu, \sigma)$ -limit over  $N_0$  and  $g : M_0 \cong N_0$ , then there exists a  $\mathcal{K}$ -embedding,  $\hat{g}$ , extending  $g$  such that  $\hat{g} : M_1 \cong N_1$ .*

**Fact 1.11.** *Let  $\mu$  be a cardinal and  $\sigma$  a limit ordinal with  $\sigma < \mu^+$ . If  $M$  is a  $(\mu, \sigma)$ -limit model, then  $M$  is a  $(\mu, \text{cf}(\sigma))$ -limit model.*

The main result of this paper provides an answer to Question 1.9.

**Theorem 3** (Main Theorem). *Let  $\mathcal{K}$  be an AEC without maximal models, and  $\mu > \text{LS}(\mathcal{K})$ . Suppose  $\mathcal{K}$  satisfies the  $\mu$ -DAP. If  $\mathcal{K}$  is  $\mu$ -Galois-stable, does not have long splitting chains, and satisfies locality of splitting<sup>1</sup>, then any two  $(\mu, \sigma_\ell)$ -limits over  $M$  for  $(\ell \in \{1, 2\})$  are isomorphic over  $M$ .*

The last section of this paper (see pages 25 and ff.) describes different approaches to Question 1.9.

We thank John Baldwin, Tapani Hyttinen and Pedro Zambrano for helping to clarify the presentation. We also thank the referee for valuable suggestions, remarks and an example of an  $\aleph_1$ -categorical AEC failing DAP over countable models but having DAP over limit models.

## 2. THE SETTING

In what follows,  $\mathcal{K}$  is assumed to be an AEC, and  $\mu$  is a cardinal  $\geq \text{LS}(\mathcal{K})$ . In this section we summarize all of the assumptions that will be made on the class  $\mathcal{K}$ , and in the subsequent sections we introduce three of the main components of the proof of the uniqueness of limit models: strong types and towers.

We will prove the uniqueness of limit models in  $\mu$ -Galois stable AECs equipped with a moderately well-behaved dependence relation. We will use  $\mu$ -splitting as the dependence relation, but any dependence relation which is local and has existence, uniqueness and extension properties suffices.

---

<sup>1</sup>See Assumption 2.5 for the precise description of long splitting chains and locality.

**Definition 2.1.** A type  $p \in \text{ga-S}(M)$   $\mu$ -splits over  $N \in \mathcal{K}_{\leq \mu}$  if and only if  $N$  is a  $\prec_{\mathcal{K}}$ -submodel of  $M$  and there exist  $N_1, N_2 \in \mathcal{K}_{\mu}$  and a  $\mathcal{K}$ -mapping  $h$  such that  $N \prec_{\mathcal{K}} N_l \prec_{\mathcal{K}} M$  for  $l = 1, 2$  and  $h : N_1 \rightarrow N_2$  with  $h \upharpoonright N = \text{id}_N$  and  $p \upharpoonright N_2 \neq h(p \upharpoonright N_1)$ .

The existence property for non- $\mu$ -splitting types follows from Galois stability in  $\mu$ :

**Fact 2.2** (Existence - Claim 3.3 of [Sh 394]). *Assume  $\mathcal{K}$  has AP and is Galois-stable in  $\mu$ . For every  $M \in \mathcal{K}_{\geq \mu}$  and  $p \in \text{ga-S}(M)$ , there exists  $N \in \mathcal{K}_{\mu}$  such that  $p$  does not  $\mu$ -split over  $N$ .*

The uniqueness and extension property of non- $\mu$ -splitting types holds for types over limit models:

**Fact 2.3** (Uniqueness - Theorem I.4.15 of [Va]). *Let  $N \prec_{\mathcal{K}} M \prec_{\mathcal{K}} M'$  be models in  $\mathcal{K}_{\mu}$  such that  $M'$  is universal over  $M$  and  $M$  is universal over  $N$ . If  $p \in \text{ga-S}(M)$  does not  $\mu$ -split over  $N$ , then there is a unique  $p' \in \text{ga-S}(M')$  such that  $p'$  extends  $p$  and  $p'$  does not  $\mu$ -split over  $N$ .*

A variation of this fact is later used in an induction construction in the proof of Theorem 6. We state it explicitly here:

**Fact 2.4** (Theorem I.4.10 of [Va]). *Let  $M, N, M^*$  be models in  $\mathcal{K}_{\mu}$ . Suppose that  $M$  is universal over  $N$  and that  $M^*$  is universal over  $M$ . If a type  $p = \text{ga-tp}(a/M)$  does not  $\mu$ -split over  $N$  then there exists an automorphism  $g$  of  $\mathfrak{C}$  fixing  $N$  such that  $\text{ga-tp}(g(a)/M^*)$  does not  $\mu$ -split over  $N$  and  $\text{ga-tp}(g(a)/M) = p$ .*

Here are the assumptions of the paper:

**Assumption 2.5.**  $\mathcal{K}$  is an AEC with the  $\mu$ -DAP over limit models and JEP, and  $\mathcal{K}$  satisfies the following properties:

- (1)  $\mathcal{K}$  is stable in  $\mu$ .
- (2) All models are submodels of a fixed monster model  $\mathfrak{C}$ .<sup>2</sup>
- (3)  $\mu$ -splitting in  $\mathcal{K}$  satisfies the following locality (sometimes called continuity) and “no long splitting chains” properties.

For all infinite  $\alpha$ , for every sequence  $\langle M_i \mid i < \alpha \rangle$  of limit models of cardinality  $\mu$  and for every  $p \in \text{ga-S}(M_{\alpha})$ , where  $M_{\alpha} = \bigcup_{i < \alpha} M_i$ , we have that

<sup>2</sup>Notice that this already implies the full AP.



- (a) If for every  $i < \alpha$ , the type  $p \upharpoonright M_i$  does not  $\mu$ -split over  $M_0$ , then  $p$  does not  $\mu$ -split over  $M_0$ .
- (b) There exists  $i < \alpha$  such that  $p$  does not  $\mu$ -split over  $M_i$ .

In the context of an AEC with the full amalgamation property and JEP, categoricity in a cardinal  $\lambda > \mu$  implies all parts of Assumption 2.5. For a proof of Assumption 2.5.1 from categoricity, see Claim 1.7 of [Sh 394] or [Ba]. The observation that assumption 2.5(3a) follows from categoricity is a consequence of Observation 6.2 and Main Lemma 9.4 of [Sh 394]. Lemma 6.3 of [Sh 394] is the statement that assumption 2.5(3b) follows from categoricity when the cofinality of the categoricity cardinal is larger than  $\mu$ .

Assumption 2.5 also holds in contexts without the assumption of categoricity. The  $\mu$ -DAP over limit models holds for free in first order classes of the form  $(\text{Mod}(T), \prec)$  for complete  $T^3$ . DAP also holds in homogeneous classes (see [Sh 3] or [Po]), in excellent classes (see [Sh 87b]) and is an axiom in the definition of finitary classes (see [HyKe]). It also holds for cats consisting of existentially closed models of positive Robinson theories ([Za]). In each of these contexts dependence relations satisfying Assumption 2.5 have been developed. Finally, the locality and existence of non- $\mu$ -splitting extensions are akin to consequences of superstability in first order logic.

**Fact 2.6** (“No long splitting chains” follows from stability in FO). *Suppose that  $T$  is first order complete. If  $T$  is stable then Assumption 2.5(3b) holds for  $\alpha$  such that  $\text{cf}(\alpha) \geq |T|^+$ .*

*Proof.* Let  $\langle M_i | i \leq \alpha \rangle$  be an increasing sequence of saturated models and  $p \in S(M_\alpha)$  be such that  $\forall i < \alpha$ ,  $p$   $\mu$ -splits over  $M_i$ . Let  $\varphi_i(\bar{x}, \bar{y})$  be a formula witnessing the splitting of  $p \upharpoonright M_{i+1}$  over  $M_i$ . As  $\text{cf}(\alpha) \geq |T|^+$ , there exists  $S \subset \alpha$  infinite such that  $i, j \in S \Rightarrow \varphi_i = \varphi_j$ .

Without loss of generality, suppose that  $\langle M_n | n \leq \omega \rangle$  is an increasing sequence of saturated models, and  $p \in S_\varphi(M_\omega)$  is such that  $\bar{a}_i, \bar{b}_i \in M_{i+1}$  witness that  $p \upharpoonright M_{i+1}$  splits over  $M_i$ . Then  $p(x_1, \bar{y}_1, \bar{z}_1, x_2, \bar{y}_2, \bar{z}_2)$  and  $\{\bar{d}_i | i < \omega\}$  witness that  $p$  has the order property, where  $\bar{d}_i = \bar{a}_i \hat{\ } \bar{b}_i \hat{\ } c_i$ ,  $c_i \in M_{i+2}$  and

$$c_i \models p \upharpoonright \{\bar{a}_k, \bar{b}_k | k \leq i\} \cup \{d_k | k < i\}.$$

<sup>3</sup>As the referee has pointed out, this is not necessarily true over arbitrary models: the class of structures for a single binary relation  $E$ , where  $E$  is an equivalence relation, and such that each class has at most two elements and at most one class has exactly one element, and if there is a one-element class then the structure is countable, with  $\prec_K$  being ordinary substructure (thus, axiomatizable in  $L_{\omega_1, \omega}$ ). This is an  $\aleph_1$ -categorical AEC with AP that does not have DAP over countable models - but has DAP over limit models.

Now use [Gr1, Lemma VII, 2.12].  $\dashv$

### 3. STRONG TYPES

Under the assumption of  $\mu$ -stability, we can define *strong types* as in [ShVi]. These strong types will allow us to achieve a better control of extensions of towers of models than what we obtain using just Galois types.

**Definition 3.1** (Definition 3.2.1 of [ShVi]). For  $M$  a  $(\mu, \theta)$ -limit model, let

$$\mathfrak{St}(M) := \left\{ (p, N) \left| \begin{array}{l} N \prec_{\mathcal{K}} M; \\ N \text{ is a } (\mu, \theta)\text{-limit model}; \\ M \text{ is universal over } N; \\ p \in \text{ga-S}(M) \text{ is non-algebraic} \\ \text{and } p \text{ does not } \mu\text{-split over } N. \end{array} \right. \right\}$$

Elements of  $\mathfrak{St}(M)$  are called *strong types*. Two strong types  $(p_1, N_1) \in \mathfrak{St}(M_1)$  and  $(p_2, N_2) \in \mathfrak{St}(M_2)$  are *parallel* iff for every  $M'$  of cardinality  $\mu$  extending  $M_1$  and  $M_2$  there exists  $q \in \text{ga-S}(M')$  such that  $q$  extends both  $p_1$  and  $p_2$  and  $q$  does not  $\mu$ -split over  $N_1$  nor over  $N_2$ . We use the notation  $(p_1, N_1) \parallel (p_2, N_2)$  for  $(p_1, N_1)$  is parallel to  $(p_2, N_2)$ .

**Remark 3.2.** Under the assumption of the existence of universal extensions, it is equivalent to say two strong types  $(p_1, N_1) \in \mathfrak{St}(M_1)$  and  $(p_2, N_2) \in \mathfrak{St}(M_2)$  are parallel iff for some  $M'$  of cardinality  $\mu$  universal over some common extension of  $M_1$  and  $M_2$  there exists  $q \in \text{ga-S}(M')$  such that  $q$  extends both  $p_1$  and  $p_2$  and  $q$  does not  $\mu$ -split over  $N_1$  and  $N_2$ .

**Lemma 3.3** (Monotonicity of parallel types). *Suppose  $M_0, M_1 \in \mathcal{K}_\mu$  and  $M_0 \prec_{\mathcal{K}} M_1$  and  $(p, N) \in \mathfrak{St}(M_1)$ . If  $M_0$  is universal over  $N$ , then we have  $(p \upharpoonright M_0, N) \parallel (p, N)$ .*

*Proof.* Straightforward using the uniqueness of non- $\mu$ -splitting extensions.  $\dashv$

**Notation 3.4.** Let  $M, M' \in \mathcal{K}_\mu$ , Suppose that  $M \prec_{\mathcal{K}} M'$ . For  $(p, N) \in \mathfrak{St}(M')$ , if  $M$  is universal over  $N$ , we define the restriction  $(p, N) \upharpoonright M \in \mathfrak{St}(M)$  to be  $(p \upharpoonright M, N)$ .

If we write  $(p, N) \upharpoonright M$ , we mean that  $p$  does not  $\mu$ -split over  $N$  and  $M$  is universal over  $N$ .

We often denote by  $\sim$  the parallelism relation between strong types in  $\mathfrak{St}(M)$ , for fixed  $M$ .

Notice that  $\sim$  is an equivalence relation on  $\mathfrak{St}(M)$  (see [Va]). Stability in  $\mu$  implies that there are few strong types over any model of cardinality  $\mu$ :

**Fact 3.5** (Claim 3.2.2 (3) of [ShVi]). *If  $\mathcal{K}$  is Galois-stable in  $\mu$ , then for any  $M \in \mathcal{K}_\mu$ ,  $|\mathfrak{St}(M)/\sim| \leq \mu$ .*

#### 4. TOWERS

To each  $(\mu, \theta)$ -limit model  $M$  we can naturally associate a  $\prec_{\mathcal{K}}$ -increasing chain  $\bar{M} = \langle M_i \in \mathcal{K}_\mu \mid i < \theta \rangle$  witnessing that  $M$  is a  $(\mu, \theta)$ -limit model (that is,  $\bigcup_{i < \theta} M_i = M$  and  $M_{i+1}$  is universal over  $M_i$ ). Furthermore, by Facts 1.10 and 1.11 we can require that this chain satisfies additional requirements such as  $M_{i+1}$  is a limit model over  $M_i$ . In this section we will be considering a related chain of models which we will refer to as a tower (see Definition 4.1). But first, we will describe how towers will be used to prove the main theorem of this paper.

To prove the uniqueness of limit models we will construct a model which is simultaneously a  $(\mu, \theta_1)$ -limit model over some fixed model  $M$  and a  $(\mu, \theta_2)$ -limit model over  $M$ . Notice that, by Fact 1.10, it is enough to construct a model  $M^*$  that is simultaneously a  $(\mu, \omega)$ -limit model and a  $(\mu, \theta)$ -limit model for arbitrary ordinal  $\theta < \mu^+$ . By Fact 1.11 we may assume that  $\theta$  is a limit ordinal  $< \mu^+$  such that  $\theta = \mu \cdot \theta$ .

So, we actually construct an array of models with  $\omega + 1$  rows and the number of columns of this array will have the same cofinality as  $\theta$ . See the big picture of the construction on page 24. We intend to carry out the construction **down** and **to the right** in that picture. In the array, the bottom right hand corner ( $M^*$ ) will be a  $(\mu, \omega)$ -limit model witnessed by a chain of models as described in the first paragraph of this section. This chain will appear in the last column of the array. We will see that  $M^*$  is a  $(\mu, \theta)$ -limit model by examining the last (the  $\omega^{\text{th}}$ ) row of the array. This last row will be an  $\prec_{\mathcal{K}}$ -increasing sequence of models,  $\bar{M}^*$  whose length will have the same cofinality as  $\theta$ . However we will not be able to guarantee that  $M_{i+1}^*$  is universal over  $M_i^*$  in this last row. Thus we need another method to conclude that  $M^*$  is a  $(\mu, \theta)$ -limit model. This involves attaching more information to our sequence  $\bar{M}^*$ . We call this accessorized sequence of models a tower (see Definition 4.1 below). Each row in our construction of the array of models will be such a tower.

Under the assumption of Galois-stability, given any sequence  $\langle a_i \mid i < \theta \rangle$  of elements with  $a_i \in M_{i+1} \setminus M_i$ , we can identify some  $N_i \prec_{\mathcal{K}} M_i$  such that

$\text{ga-tp}(a_i/M_i)$  does not  $\mu$ -split over  $N_i$ . Furthermore, by Assumption 2.5, we may choose this  $N_i$  such that  $M_i$  is a limit model over  $N_i$ . We abbreviate this situation by the triple  $(\bar{M}, \bar{a}, \bar{N})$ .

**Definition 4.1** (Towers). Let  $(I, <)$  be a well ordering of cardinality  $< \mu^+$ . For cleaner notation, we will identify  $I$  with  $\theta$ , its order-type, and we will denote the successor of  $i$  in the ordering  $I$  by  $i + 1$  when it is clear. Then, we define a *tower* to be a triple  $(\bar{M}, \bar{a}, \bar{N})$  where  $\bar{M} = \langle M_i \mid i < \theta \rangle$  is a  $\prec_{\mathcal{K}}$ -increasing sequence of limit models of cardinality  $\mu$ ;  $\bar{a} = \langle a_i \mid i < \theta \rangle$  and  $\bar{N} = \langle N_i \mid i < \theta \rangle$  satisfy  $a_i \in M_{i+1} \setminus M_i$ ;  $\text{ga-tp}(a_i/M_i)$  does not  $\mu$ -split over  $N_i$ ;  $M_i$  is a  $(\mu, \sigma)$ -limit model over  $N_i$ .

**Notation 4.2.** We denote by  $\mathcal{K}_{\mu, I}^*$  the set of towers of the form  $(\bar{M}, \bar{a}, \bar{N})$  where the sequences  $\bar{M}$ ,  $\bar{a}$  and  $\bar{N}$  are indexed by  $I$ . Occasionally,  $I$  will be an ordinal  $\theta$  with the usual ordering, and we write  $\mathcal{K}_{\mu, \theta}^*$  for this set of towers. At times, we will be considering towers based on different well orderings  $I$  and  $I'$  simultaneously. In these contexts if  $i \in I \cap I'$ , the notation  $i + 1$  is not necessarily well-defined so we will use the notation  $\text{succ}_I(i)$  for the successor of  $i$  in the ordering  $I$ . Finally when  $I$  is a sub-order of  $I'$  for any  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I'}^*$  we write  $(\bar{M}, \bar{a}, \bar{N}) \upharpoonright I$  for the tower in  $\mathcal{K}_{\mu, I}^*$  given by the subsequences  $\langle M_i \mid i \in I \rangle$ ,  $\langle N_i \mid i \in I \rangle$  and  $\langle a_i \mid i \in I \rangle$ .

In addition to having control over the last row of the array, we also need to be able to guarantee that the last column of the tower witnesses that  $M^*$  is a  $(\mu, \omega)$ -limit model. This will be done by prescribing the following ordering on rows of the array:

**Definition 4.3.** For towers  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$  and  $(\bar{M}', \bar{a}', \bar{N}') \in \mathcal{K}_{\mu, I'}^*$  with  $I \subseteq I'$ , we write  $(\bar{M}, \bar{a}, \bar{N}) < (\bar{M}', \bar{a}', \bar{N}')$  if and only if for every  $i \in I$ ,  $a_i = a'_i$ ,  $N_i = N'_i$  and  $M'_i$  is a proper universal extension of  $M_i$ .

**Remark 4.4.** The ordering  $<$  on towers is identical to the ordering  $<_{\mu}^c$  defined in [ShVi]. The superscript was used by Shelah and Villaveces to distinguish this ordering from others. We only use one ordering on towers, so we omit the superscripts and subscripts here.

Once we have established an ordering on towers, we can define a specific tower which will be called a *union of an increasing sequence of towers*. Suppose that  $\langle (\bar{M}, \bar{a}, \bar{N})^\gamma \in \mathcal{K}_{\mu, I_\gamma}^* \mid \gamma < \beta \rangle$  is an increasing sequence of towers such that the index set  $I_\gamma$  of  $(\bar{M}, \bar{a}, \bar{N})^\gamma$  is a sub-ordering of the

index set  $I_{\gamma'}$  for  $(\bar{M}, \bar{a}, \bar{N})^{\gamma'}$  whenever  $\gamma < \gamma'$ . Let  $I_\beta := \bigcup_{\gamma < \beta} I_\gamma$ . Then denote by  $(\bar{M}, \bar{a}, \bar{N})^\beta \in \mathcal{K}_{\mu, I_\beta}^*$  the “union” of the sequence of towers where

$$\begin{aligned}\bar{a}_i^\beta &= a_i^{\min\{\gamma | i \in I_\gamma\}}, \\ \bar{N}_i^\beta &= N_i^{\min\{\gamma | i \in I_\gamma\}}\end{aligned}$$

and

$$\bar{M}^\beta = \langle M_i^\beta \mid i \in \bigcup_{\gamma < \beta} I_\gamma \rangle$$

with

$$M_i^\beta = \bigcup_{\gamma < \beta} \bigcup_{i \in I_\gamma} M_i^\gamma.$$

By assumption 2.5.3a,  $(\bar{M}, \bar{a}, \bar{N})^\beta$  is indeed a tower.

Notice that we do not assume an individual tower to be continuous. Nor do we assume that inside of a tower  $M_{i+1}$  is universal over  $M_i$ . If one considers the approach of defining an array of models row by row, then generally (even in the first order case) even if all rows are continuous and satisfy the universality property mentioned in this paragraph, it is not necessarily true that the union of these rows will be a tower in which every model is universal over its predecessors.

For a tower  $(\bar{M}, \bar{a}, \bar{N})$ , it was shown in [ShVi], that even if  $M_{i+1}$  is not universal over  $M_i$ , one can conclude that  $\bigcup_{i < \theta} M_i$  is a  $(\mu, \theta)$ -limit model provided that all types over each of the  $M_i$  are realized by a sufficient number of  $a_j$ s in the tower. Unfortunately constructing such a tower meeting these along with all of our other requirements is beyond reach. However, in [Va], VanDieren showed that slightly less was needed (see Definition 4.5). In [Va], the amalgamation property is not assumed resulting in noise that can be avoided in our context. Thus because we have at our disposal the AP, we provide a complete, undistracted proof here.

**Definition 4.5** (Relatively Full Towers). Suppose that  $I$  is a well-ordered set such that there exists a cofinal sequence  $\langle i_\alpha \mid \alpha < \theta \rangle$  of  $I$  of order type  $\theta$  such that there are  $\mu \cdot \omega$  many elements between  $i_\alpha$  and  $i_{\alpha+1}$ .

Let  $(\bar{M}, \bar{a}, \bar{N})$  be a tower indexed by  $I$  such that each  $M_i$  is a  $(\mu, \sigma)$ -limit model. For each  $i$ , let  $\langle M_i^\gamma \mid \gamma < \sigma \rangle$  witness that  $M_i$  is a  $(\mu, \sigma)$ -limit model. The tower  $(\bar{M}, \bar{a}, \bar{N})$  is *full relative to*  $(M_i^\gamma)_{\gamma < \sigma, i \in I}$  iff for every  $\gamma < \sigma$  and every  $(p, M_i^\gamma) \in \mathfrak{St}(M_i)$  with  $i_\alpha \leq i < i_{\alpha+1}$ , there exists  $j \in I$  with  $i \leq j < i_{\alpha+1}$  such that  $(\text{ga-tp}(a_j/M_j), N_j)$  and  $(p, M_i^\gamma)$  are parallel.

**Theorem 4** (Relatively full towers provide limit models). *Let  $\theta$  be a limit ordinal  $< \mu^+$  satisfying  $\theta = \mu \cdot \theta$ . Suppose that  $I$  is a well-ordered set as in Definition 4.5.*

*Let  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$  be a tower made up of  $(\mu, \sigma)$ -limit models, for some fixed  $\sigma < \mu^+$ . If  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$  is full relative to  $(M_i^\gamma)_{i \in I, \gamma < \sigma}$ , then  $M := \bigcup_{i \in I} M_i$  is a  $(\mu, \theta)$ -limit model.*

*Proof.* Without loss of generality we may assume that  $\bar{M}$  is continuous. Let  $M'$  be a  $(\mu, \theta)$ -limit model over  $M_{i_0}$  witnessed by  $\langle M'_\alpha \mid \alpha < \theta \rangle$ . By  $\mu$ -DAP, we may assume that  $M' \cap M = M_{i_0}$ . Since  $\theta = \mu \cdot \theta$ , we may also arrange things so that the universe of  $M'_\alpha$  is  $\mu \cdot \alpha$  and  $\alpha \in M'_{\alpha+1}$ .

We will construct an isomorphism between  $M$  and  $M'$  by induction on  $\alpha < \theta$ . Define an increasing and continuous sequence of  $\prec_{\mathcal{K}}$ -mappings  $\langle h_\alpha \mid \alpha < \theta \rangle$  such that

- (1)  $h_\alpha : M_{i_\alpha+j} \rightarrow M'_{\alpha+1}$  for some  $j < \mu \cdot \omega$
- (2)  $h_0 = \text{id}_{M_{i_0}}$  and
- (3)  $\alpha \in \text{rg}(h_{\alpha+1})$ .

For  $\alpha = 0$  take  $h_0 = \text{id}_{M_{i_0}}$ . For  $\alpha$  a limit ordinal let  $h_\alpha = \bigcup_{\beta < \alpha} h_\beta$ . Since  $\bar{M}$  is continuous, the induction hypothesis gives us that  $h_\alpha$  is a  $\prec_{\mathcal{K}}$ -mapping from  $M_{i_\alpha}$  into  $M'_\alpha$  allowing us to satisfy condition (1) of the construction.

Suppose that  $h_\alpha$  has been defined. Let  $j < \mu \cdot \omega$  be such that  $h_\alpha : M_{i_\alpha+j} \rightarrow M'_{\alpha+1}$ . There are two cases: either  $\alpha \in \text{rg}(h_\alpha)$  or  $\alpha \notin \text{rg}(h_\alpha)$ . First suppose that  $\alpha \in \text{rg}(h_\alpha)$ . Since  $M'_{\alpha+2}$  is universal over  $M'_{\alpha+1}$ , it is also universal over  $h_\alpha(M_{i_\alpha+j})$ . This allows us to extend  $h_\alpha$  to  $h_{\alpha+1} : M_{i_{\alpha+1}} \rightarrow M'_{\alpha+2}$ .

Now consider the case when  $\alpha \notin \text{rg}(h_\alpha)$ . Since  $\langle M_{i_\alpha+j}^\gamma \mid \gamma < \sigma \rangle$  witnesses that  $M_{i_\alpha+j}$  is a  $(\mu, \sigma)$ -limit model, by Assumption 2.5, there exists  $\gamma < \sigma$  such that  $\text{ga-tp}(\alpha/M_{i_\alpha+j})$  does not  $\mu$ -split over  $M_{i_\alpha+j}^\gamma$ . By our choice of  $\bar{M}'$  disjoint from  $\bar{M}$  outside of  $M_{i_0}$ , we know that  $\alpha \notin M_{i_\alpha+j}$ . Thus  $\text{ga-tp}(\alpha/M_{i_\alpha+j})$  is non-algebraic. By relative fullness of  $(\bar{M}, \bar{a}, \bar{N})$ , there exists  $j'$  with  $j \leq j' < i_{\alpha+1}$  such that  $(\text{ga-tp}(\alpha/M_{i_\alpha+j'}), M_{i_\alpha+j'}^\gamma)$  is parallel to  $(\text{ga-tp}(a_{i_{\alpha+1}+j'}/M_{i_{\alpha+1}+j'}), N_{i_{\alpha+1}+j'})$ . In particular we have that

$$(*) \quad \text{ga-tp}(a_{i_{\alpha+1}+j'}/M_{i_{\alpha+1}+j'}) = \text{ga-tp}(\alpha/M_{i_\alpha+j}).$$

We can extend  $h_\alpha$  to an automorphism  $h'$  of  $\mathfrak{C}$ . An application of  $h'$  to  $(*)$  gives us

$$(**) \quad \text{ga-tp}(h'(a_{i_{\alpha+1}+j'})/h_\alpha(M_{i_\alpha+j})) = \text{ga-tp}(\alpha/h_\alpha(M_{i_\alpha+j})).$$

Since  $M'_{\alpha+2}$  is universal over  $h_\alpha(M_{i_\alpha})$ , we may extend  $h_\alpha$  to a  $\mathcal{K}$ -mapping  $h_{\alpha+1} : M_{i_{\alpha+1}+j'} \rightarrow M'_{\alpha+2}$  such that  $h_{\alpha+1}(a_{i_{\alpha+1}+j'}) = \alpha$ .

Let  $h := \bigcup_{\alpha < \theta} h_\alpha$ . Clearly  $h : M \rightarrow M'$ . To see that  $h$  is an isomorphism, notice that condition (3) of the construction forces  $h$  to be surjective.  $\dashv$

## 5. UNIQUENESS OF LIMIT MODELS

We now begin the construction of our array of models and  $M^*$ . Let  $\theta$  be an ordinal as in the previous section. The goal will be to build an array of models with  $\omega + 1$  rows so that the bottom row of the array is a relatively full tower indexed by a set of cofinality  $\theta$ . To do this, we will be adding elements to the index set of towers row by row so that at stage  $n$  of our construction the tower that we build is indexed by  $I_n$  described here:

**Notation 5.1.** The index sets  $I_n$  will be defined inductively so that  $\langle I_n \mid n < \omega + 1 \rangle$  is an increasing and continuous chain of well-ordered sets. We fix  $I_0$  to be an index set of order type  $\theta + 1$  and will denote it by  $\langle i_\alpha \mid \alpha \leq \theta \rangle$ . We will refer to the members of  $I_0$  by name in many stages of the construction. These indices serve as anchors for the members of the remaining index sets in the array. Next we demand that for each  $n < \omega$ ,  $\{j \in I_n \mid i_\alpha < j < i_{\alpha+1}\}$  has order type  $\mu \cdot n$  such that each  $I_n$  has supremum  $i_\theta$ . An example of such  $\langle I_n \mid n \leq \omega \rangle$  is  $I_n = \theta \times (\mu \cdot n) \cup \{i_\theta\}$  ordered lexicographically, where  $i_\theta$  is an element  $\geq$  each  $i \in \bigcup_{n < \omega} I_n$ . Also, let  $I = \bigcup_{n < \omega} I_n$ .

To prove the main theorem of the paper, we need to prove that for a fixed  $M \in \mathcal{K}$  of cardinality  $\mu$  any  $(\mu, \theta)$ -limit and  $(\mu, \omega)$ -limit model over  $M$  are isomorphic over  $M$ . Let us begin by fixing  $M \in \mathcal{K}_\mu$  and  $\theta$  such that  $\mu \cdot \theta = \theta$ . Without loss of generality,  $M$  is a limit model. We define by induction on  $n \leq \omega$  a  $<$ -increasing and continuous sequence of towers  $(\bar{M}, \bar{a}, \bar{N})$  such that

- (1)  $(\bar{M}, \bar{a}, \bar{N})^0$  is a tower with  $M_0^0 = M$ .
- (2)  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I_n}^*$
- (3) For every  $(p, N) \in \mathfrak{St}(M_i^n)$  with  $i_\alpha \leq i < i_{\alpha+1}$  there is  $j \in I_{n+1}$  with  $i < j < i_{\alpha+1}$  so that  $(\text{ga-tp}(a_j/M_j^{n+1}), N_j^{n+1})$  and  $(p, N)$  are parallel.
- (4)  $M_{i_{\alpha+1}}^{n+1}$  is a  $(\mu, \mu)$ -limit model over  $\bigcup_{j < i_{\alpha+1}} M_j^{n+1}$ .

Given  $M$ , we can find a tower  $(\bar{M}, \bar{a}, \bar{N})^0 \in \mathcal{K}_{\mu, I_0}^*$  with  $M_0^0 = M$  because of the existence of universal extensions and because of Assumption 2.5.3b. The last pages (Page 24 onward) of this section provide a picture of this construction of an array of models, explanations for carrying out the final

stage of the construction and a proof that this is sufficient to prove the main theorem. We spend most of the remainder of this section verifying that it is possible to carry out the induction step of the construction. This is a particular case of Theorem II.7.1 of [Va]. But since our context is somewhat easier, we do not encounter so many obstacles as in [Va] and we provide a different, more direct proof here:

**Theorem 5** (Dense  $<$ -extension property). *Given  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I_n}^*$  there exists  $(\bar{M}', \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I_{n+1}}^*$  such that  $(\bar{M}', \bar{a}, \bar{N}) < (\bar{M}, \bar{a}, \bar{N})$  and for each  $(p, N) \in \mathfrak{St}(M_i)$  with  $i_\alpha \leq i < i_{\alpha+1}$ , there exists  $j \in I_{n+1}$  with  $i < j < i_{\alpha+1}$  such that  $(\text{ga-tp}(a_j/M'_j), N_j)$  and  $(p, N)$  are parallel. Here, the  $M_i$ 's are defined for  $i \in I_n$  and the  $M'_j$  are defined for  $j \in I_{n+1}$ .*

Before we prove Theorem 5, we prove a slightly weaker extension property, one in which we can find an extension of the tower  $(\bar{M}, \bar{a}, \bar{N})$  of the same index set:

**Lemma 5.2** ( $<$ -extension property). *Given  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$ , there exists a  $<$ -extension  $(\bar{M}', \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$  of  $(\bar{M}, \bar{a}, \bar{N})$  such that for each limit  $i$ ,  $M'_i$  is a  $(\mu, \mu)$ -limit model over  $\bigcup_{j < i} M'_j$ .*

*Proof.* Given  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$  we will define a  $<$ -extension  $(\bar{M}', \bar{a}, \bar{N})$  by induction on  $i \in I$ . Notice that a straightforward induction proof is not sufficient here for if we have defined  $\langle M_j \mid j \leq i \rangle$  as a tower extending  $(\bar{M}, \bar{a}, \bar{N})$  restricted to  $\langle j \mid j \leq i \rangle$  and are at the stage of defining  $M'_{i+1}$ , we may be faced with an impossible task: during our construction we may have inadvertently placed inside  $M'_i$  witnesses for the splitting of the type of  $a_{i+1}$  over  $N_{i+1}$ ; this would prevent us from extending  $M'_i$  to  $M'_{i+1}$  so that  $\text{ga-tp}(a_{i+1}/M'_{i+1})$  does not  $\mu$ -split over  $N_{i+1}$ . Therefore, we will instead define approximations  $(M_i^+)$  for  $M'_i$  by induction on  $i \in I$  and at each stage  $i$  of the induction we will make adjustments of the previously defined approximation  $M_j^+$  for  $j < i$ . This leads us into defining  $M_i^+$  and a directed system of  $\prec_{\mathcal{K}}$ -embeddings  $\langle f_{j,i} \mid j < i \in I \rangle$  such that for  $i \in I$ ,  $M_i \prec_{\mathcal{K}} M_i^+$  for  $j \leq i$ ,  $f_{j,i} : M_j^+ \rightarrow M_i^+$  and  $f_{j,i} \upharpoonright M_j = \text{id}_{M_j}$ . We further require that  $M_{i+1}^+$  is a limit model over  $f_{i,i+1}(M_i^+)$  and  $\text{ga-tp}(a_i/f_{i,i+1}(M_i^+))$  does not  $\mu$ -split over  $N_i$ . When  $i$  is a limit, we choose  $M_i^+$  to be a  $(\mu, \mu)$ -limit model over  $\bigcup_{j < i} f_{j,i}(M_j^+)$ .

This construction is done by induction on  $i \in I$  using the existence of non- $\mu$ -splitting extensions. Suppose that  $\langle M_k^+ \mid k \leq i \rangle$  and  $\langle f_{k,l} \mid k \leq l \leq i \rangle$



have been defined. We explain how to define  $M_{i+1}^+$  and  $f_{i,i+1}$ . The rest of the definitions required for the  $i + 1^{\text{st}}$  stage are dictated by the requirement that we are forming a directed system. Let  $M_{i+1}^*$  be a limit model over both  $M_i^+$  and  $M_{i+1}$ . Since  $\text{ga-tp}(a_{i+1}/M_{i+1})$  does not  $\mu$ -split over  $N_{i+1}$ , by Fact 2.4 there exists  $f \in \text{Aut}(\mathfrak{C}/M_{i+1})$  so that  $\text{ga-tp}(a_{i+1}/f(M_{i+1}^*))$  does not  $\mu$ -split over  $N_{i+1}$ . Take  $M_{i+1}^+ := f(M_{i+1}^*)$  and  $f_{i,i+1} := f \upharpoonright M_i^+$ .

At limit stages we take direct limits so that  $f_{j,i} \upharpoonright M_j = \text{id}_{M_j}$ . This is possible by Subclaims II.7.10 and II.7.11 of [Va] or see Claim 2.17 of [GrVa2]. Take an extension of the direct limit that is both universal over  $M_i$  and is a  $(\mu, \mu)$ -limit over  $\bigcup_{j < i} f_{j,i}(M_j)$  and call this  $M_i^+$ . Notice that we do not obtain a continuous tower; continuity will be recovered later using reduced towers.

Let  $f_{j,\sup\{I\}}$  and  $M'_{\sup\{I\}}$  be the direct limit of this system such that  $f_{j,\sup\{I\}} \upharpoonright M_j = \text{id}_{M_j}$ . We can now define  $M'_j := f_{j,\sup\{I\}}(M_j^+)$  for each  $j \in I$ . By construction, we have that  $\text{ga-tp}(a_i/f_{i,i+1}(M_i^+))$  does not  $\mu$ -split over  $N_i$ . Mapping into  $M_{\sup(I)}$  by  $f_{i+1,\sup(I)}$ , and noting that both  $a_i$  and  $N_i$  are fixed by  $f_{i+1,\sup(I)}$ , we conclude that  $\text{ga-tp}(a_i/M'_i)$ , as required.  $\dashv$

We can now use the extension property for towers of the same index set from Lemma 5.2 to prove the dense extension property which allows us to grow the index set as we add elements to the models in the extension.

*Proof of Theorem 5.* Given  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I_n}^*$ , let  $(\bar{M}', \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I_n}^*$  be an extension of  $(\bar{M}, \bar{a}, \bar{N})$  as in Lemma 5.2 so that each  $M'_{i_{\alpha+1}}$  is a  $(\mu, \mu)$ -limit model over  $\bigcup_{j < i_{\alpha+1}} M'_j$ .

For each  $i_\alpha$ , let  $\langle M'_l \mid l \in I_{n+1}, i_\alpha + \mu \cdot n < l < i_{\alpha+1} \rangle$  witness that  $M'_{i_{\alpha+1}}$  is a  $(\mu, \mu)$ -limit model over  $\bigcup_{j < i_{\alpha+1}} M'_j$ . Without loss of generality we may assume that each of these  $M'_l$  is a limit model over its predecessor.

Fix  $\{(p, N)_{i_\alpha}^l \mid i_\alpha + \mu \cdot n < l < i_{\alpha+1}\}$  an enumeration of  $\bigcup \{\text{St}(M_i) : i \in I_n, i_\alpha \leq i < i_{\alpha+1}\}$ . By our choice of  $I_{n+1}$  and stability in  $\mu$ , such an enumeration is possible. Since  $M'_{\text{succ}_{I_{n+1}}(l)}$  is universal over  $M'_l$ , there exists a realization in  $M'_{\text{succ}_{I_{n+1}}(l)}$  of the non- $\mu$ -splitting extension of  $p_{i_\alpha}^l$  to  $M'_l$ . Let  $a_l$  be this realization and take  $N_l := N_{i_\alpha}^l$ .

Notice that  $(\langle M'_j \mid j \in I_{n+1} \rangle, \langle a_j \mid j \in I_{n+1} \rangle, \langle N_j \mid j \in I_{n+1} \rangle)$  provide the desired extension of  $(\bar{M}, \bar{a}, \bar{N})$  in  $\mathcal{K}_{\mu, I_{n+1}}^*$ .  $\dashv$

We are almost ready to carry out the complete construction. However, notice that Theorem 5 does not provide us with a continuous extension. Therefore the bottom (i.e. the  $\omega + 1^{st}$ ) row of our array may not be continuous which would prevent us from applying Theorem 4 to conclude that  $M^*$  is a  $(\mu, \theta)$ -limit model. So we will further require that the towers that occur in the rows of our array are all continuous. This can be guaranteed by restricting ourselves to reduced towers as in [ShVi] and [Va].

**Definition 5.3.** A tower  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$  is said to be *reduced* provided that for every  $(\bar{M}', \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$  with  $(\bar{M}, \bar{a}, \bar{N}) \leq (\bar{M}', \bar{a}, \bar{N})$  we have that for every  $i \in I$ ,

$$(*)_i \quad M'_i \cap \bigcup_{j \in I} M_j = M_i.$$

If we take a  $<$ -increasing chain of reduced towers, the union will be reduced. The following fact appears as Theorem 3.1.14 of [ShVi]. We provide the proof for completeness.

**Fact 5.4.** Let  $\langle (\bar{M}, \bar{a}, \bar{N})^\gamma \in \mathcal{K}_{\mu, I_\gamma}^* \mid \gamma < \beta \rangle$  be a  $<$ -increasing and continuous sequence of reduced towers such that the sequence is continuous in the sense that for a limit  $\gamma < \beta$ , the tower  $(\bar{M}, \bar{a}, \bar{N})^\gamma$  is the union of the towers  $(\bar{M}, \bar{a}, \bar{N})^\zeta$  for  $\zeta < \gamma$ . Then the union of the sequence of towers  $\langle (\bar{M}, \bar{a}, \bar{N})^\gamma \in \mathcal{K}_{\mu, I_\gamma}^* \mid \gamma < \beta \rangle$  is itself a reduced tower.

*Proof.* Suppose that  $(\bar{M}, \bar{a}, \bar{N})^\beta$  is not reduced. Let  $(\bar{M}', \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I_\beta}^*$  witness this. Then there exists an  $i \in I_\beta$  and an element  $b$  such that  $b \in (M'_i \cap \bigcup_{j \in I_\beta} M_j^\beta) \setminus M_i^\beta$ . There exists  $\gamma < \beta$  such that  $b \in \bigcup_{j \in I_\gamma} M_j^\gamma \setminus M_i^\gamma$ . Notice that  $(\bar{M}', \bar{a}, \bar{N}) \upharpoonright I_\gamma$  witnesses that  $(\bar{M}, \bar{a}, \bar{N})^\gamma$  is not reduced.  $\dashv$

The following appears in [ShVi] (Theorem 3.1.13).

**Fact 5.5** (Density of reduced towers). *There exists a reduced  $<$ -extension of every tower in  $\mathcal{K}_{\mu, I}^*$ .*

*Proof.* Assume for the sake of contradiction that no  $<$ -extension of  $(\bar{M}, \bar{a}, \bar{N})$  is reduced. This allows us to construct a  $\leq$ -increasing and continuous sequence of towers  $\langle (\bar{M}, \bar{a}, \bar{N})^\zeta \in \mathcal{K}_{\mu, I}^* \mid \zeta < \mu^+ \rangle$  such that  $(\bar{M}, \bar{a}, \bar{N})^{\zeta+1}$  witnesses that  $(\bar{M}, \bar{a}, \bar{N})^\zeta$  is not reduced. The construction is done inductively in the obvious way.

For each  $b \in \bigcup_{\zeta < \mu^+, i \in I} M_i^\zeta$  define

$$i(b) := \min \{ i \in I \mid b \in \bigcup_{\zeta < \mu^+} \bigcup_{j \leq i} M_j^\zeta \} \text{ and}$$

$$\zeta(b) := \min \{ \zeta < \mu^+ \mid b \in M_{i(b)}^\zeta \}.$$

$\zeta(\cdot)$  can be viewed as a function from  $\mu^+$  to  $\mu^+$ . Since  $|I| = \mu$  and each  $M_i^\zeta$  has cardinality  $\mu$ , there exists a club  $E = \{ \delta < \mu^+ \mid \forall b \in \bigcup_{i \in I} M_i^\delta, \zeta(b) < \delta \}$ . Actually, all we need is for  $E$  to be non-empty.

Fix  $\delta \in E$ . By construction  $(\bar{M}, \bar{a}, \bar{N})^{\delta+1}$  witnesses the fact that  $(\bar{M}, \bar{a}, \bar{N})^\delta$  is not reduced. So we may fix  $i \in I$  and  $b \in M_i^{\delta+1} \cap \bigcup_{j \in I} M_j^\delta$  such that  $b \notin M_i^\delta$ . Since  $b \in M_i^{\delta+1}$ , we have that  $i(b) \leq i$ . Since  $\delta \in E$ , we know that there exists  $\zeta < \delta$  such that  $b \in M_{i(b)}^\zeta$ . Because  $\zeta < \delta$  and  $i(b) \leq i$ , this implies that  $b \in M_i^\delta$  as well. This contradicts our choice of  $i$  and  $b$  witnessing the failure of  $(\bar{M}, \bar{a}, \bar{N})^\delta$  to be reduced.  $\dashv$

By revising the proof of Lemma 5.2, we can conclude:

**Lemma 5.6.** *Suppose that  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$  is reduced. If  $I_0$  is an initial segment of  $I$ , then  $(\bar{M}, \bar{a}, \bar{N}) \restriction I_0$  is reduced.*

*Proof.* Suppose that  $(\bar{M}, \bar{a}, \bar{N}) \restriction I_0$  is not reduced. Let  $(\bar{M}', \bar{a} \restriction I_0, \bar{N} \restriction I_0)$  and  $\delta < j \in I_0$  with  $b \in (M'_\delta \cap M_j) \setminus M_\delta$  witness this. We can apply the inductive step of Lemma 5.2 (replacing an initial segment of the construction there with  $\bar{M}'$ ), to find  $(\bar{M}'', \bar{a}, \bar{N})$  an extension of  $(\bar{M}, \bar{a}, \bar{N})$  such that there is a  $\prec_{\mathcal{K}}$ -mapping  $f$  from the models of  $\bar{M}'$  into the models of  $\bar{M}''$  with  $f \restriction M_j = \text{id}_{M_j}$ . Notice that  $(\bar{M}'', \bar{a}, \bar{N})$  and  $b, \delta, j$  will witness that  $(\bar{M}, \bar{a}, \bar{N})$  is not reduced.  $\dashv$

**Theorem 6** (Reduced towers are continuous). *If  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$  is reduced, then it is continuous, namely for each limit  $i$  in  $I$ ,  $M_i = \bigcup_{j < i} M_j$ .*

*Proof of Theorem 6.* Suppose the claim fails for  $\mu$ . Let  $\delta$  be the minimal limit ordinal such that there exists an  $I$  and  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, I}^*$  a reduced tower discontinuous at the  $\delta^{\text{th}}$  element of  $I$ . We can apply Lemma 5.6 to assume without loss of generality that  $I = \delta + 1$ . Fix  $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, \delta+1}^*$  reduced and discontinuous at  $\delta$  with  $b \in M_\delta \setminus \bigcup_{i < \delta} M_i$ .

**Claim 5.7.** *There exists a  $<$ -extension of  $(\bar{M}, \bar{a}, \bar{N}) \restriction \delta$ , say  $(\bar{M}', \bar{a} \restriction \delta, \bar{N} \restriction \delta) \in \mathcal{K}_{\mu, \delta}^*$  containing  $b$ .*

Notice that Claim 5.7 yields Theorem 6: Let  $M'_\delta \prec_{\mathcal{K}} \mathfrak{C}$  be a limit model universal over  $M_\delta$  containing  $\bigcup_{i < \delta} M'_i$ . Notice that  $(\bar{M}', \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, \delta+1}^*$  is an extension of  $(\bar{M}, \bar{a}, \bar{N})$  witnessing that  $(\bar{M}, \bar{a}, \bar{N})$  is not reduced.

The proof of Claim 5.7 requires several steps:

*Proof of Claim 5.7.* We use the minimality of  $\delta$  and the density of reduced towers (Fact 5.5) to build a  $<$ -increasing and continuous sequence of reduced (and continuous) towers  $\langle (\bar{M}, \bar{a}, \bar{N})^\zeta \in \mathcal{K}_{\mu, \delta}^* \mid \zeta < \delta \rangle$  such that  $(\bar{M}, \bar{a}, \bar{N})^0 := (\bar{M}, \bar{a}, \bar{N}) \upharpoonright \delta$ . This gives us a  $\delta$  by  $\delta$  array of models. If  $b$  appears in this array, we are done. So let us suppose that  $\text{ga-tp}(b/\bigcup_{i < \delta} M_i^i)$  is non-algebraic. Since  $\bigcup_{i < \delta} M_i^i$  is a  $(\mu, \delta)$ -limit model (witnessed by the diagonal of this array), we conclude by Assumption 2.5.3b that there exists  $\xi < \delta$  such that  $\text{ga-tp}(b/\bigcup_{i < \delta} M_i^i)$  does not  $\mu$ -split over  $M_\xi^\xi$ .

We will find a  $<$ -extension of  $(\bar{M}, \bar{a}, \bar{N}) \upharpoonright \delta$  by defining an  $\prec_{\mathcal{K}}$ -increasing chain of models  $\langle N'_i \mid i < \alpha \rangle$  and an increasing chain of  $\prec_{\mathcal{K}}$ -mappings  $\langle h_i \mid i < \delta \rangle$ . The pre-image of  $N'_i$  under an extension of  $\bigcup_{i < \delta} h_i$  will form a sequence  $\bar{M}'$  such that  $(\bar{M}, \bar{a}, \bar{N}) \upharpoonright \delta < (\bar{M}', \bar{a} \upharpoonright \delta, \bar{N} \upharpoonright \delta)$ ,  $b \in M'_{\xi+1}$  and  $M'_i = M_i^i$  for all  $i < \xi$ .

We choose by induction on  $i < \delta$  a  $\prec_{\mathcal{K}}$ -increasing and continuous chain of limit models  $\langle N'_i \in \mathcal{K}_\mu \mid i < \delta \rangle$  and an increasing and continuous sequence of  $\prec_{\mathcal{K}}$ -mappings  $\langle h_i \mid i < \delta \rangle$  satisfying

- (1)  $N'_{i+1}$  is a limit model and is universal over  $N'_i$
- (2)  $h_i : M_i^i \rightarrow N'_i$
- (3)  $M_i^{i+1}$  is a limit model over  $h_i(M_i^i)$
- (4)  $\text{ga-tp}(h_{i+1}(a_i)/N'_i)$  does not  $\mu$ -split over  $h_i(N_i)$
- (5) for  $i \leq \xi$ ,  $N'_i = M_i^i$  with  $h_i = \text{id}_{M_i^i}$
- (6) for  $i = \xi + 1$ ,  $b \in N'_{\xi+1}$  and
- (7) for  $i = \xi + 1$ ,  $N'_{\xi+1}$  is a limit model over  $\bigcup_{j < \delta} M_j^j$ .

THE CONSTRUCTION: Condition (5) determines immediately the definition of  $N'_i$  and  $h_i$  for  $i \leq \xi$ . We proceed with the rest of the construction by induction on  $i$  for  $\xi < i < \delta$ . To guarantee continuity of the  $N'$  and  $h$ 's, if  $i$  is a limit ordinal  $\geq \xi$ , let  $N'_i = \bigcup_{j < i} N'_j$  and  $h_i = \bigcup_{j < i} h_j$ .

At readers request, we have provided the details of the construction for the critical steps  $i = \xi + 1$  and  $i = \xi + 2$ . To carry out the construction for successors beyond  $\xi + 2$ , follow the steps for stage  $\xi + 2$  described below.

DEFINING  $h_{\xi+1}$  AND  $N'_{\xi+1}$ : Suppose that  $N'_\xi$  and  $h_\xi$  have been defined according to the construction, specifically satisfying condition (5). Since  $(\bar{M}, \bar{a}, \bar{N})$  is a tower,  $\text{ga-tp}(a_\xi/M_\xi^\xi)$  does not  $\mu$ -split over  $N_\xi$ . Since  $N'_\xi = M_\xi^\xi$

and  $h_\xi = \text{id}_{M_\xi^\xi}$ , we get condition (4) by invariance so long as  $h_{\xi+1}$  extends  $h_\xi (= \text{id}_{M_\xi^\xi})$ . So it suffices to choose  $N'_{\xi+1}$  to be some limit model over  $\bigcup_{i < \delta} M_i^i$  containing  $b$ . Because  $M_{\xi+1}^{\xi+2}$  is universal over  $M_{\xi+1}^\xi$  we can fix a mapping  $h_{\xi+1} : M_{\xi+1}^{\xi+1} \rightarrow M_{\xi+1}^{\xi+2}$  which is the identity on  $M_{\xi+1}^{\xi+1}$ .

DEFINING  $h_{\xi+2}$  AND  $N'_{\xi+2}$ : For this next stage of the construction we need to take a little more care to obtain condition (4). We will begin with finding the mapping  $h_{\xi+2}$ . First, fix  $\bar{h}_{\xi+1}$  an automorphism of  $\mathfrak{C}$  extending  $h_{\xi+1}$ . Notice we may assume that  $\bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2}) \prec_{\mathcal{K}} N'_{\xi+1}$  because of the universality of  $N'_{\xi+1}$  over  $h_{\xi+1}(M_{\xi+1}^{\xi+1})$  guaranteed by conditions (7) and (3). Moreover, we can arrange the mapping  $\bar{h}_{\xi+1}$  so that

$$(\triangle) \quad N'_{\xi+1} \text{ is a limit model over } \bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2}).$$

Because we started out with a  $\delta$  by  $\delta$  array originating from a sequence of towers, we have that  $\text{ga-tp}(a_{\xi+1}/M_{\xi+1}^{\xi+1})$  does not  $\mu$  split over  $N_{\xi+1}$ . Thus invariance gives us

$$(*) \quad p := \text{ga-tp}(\bar{h}_{\xi+1}(a_{\xi+1})/h_{\xi+1}(M_{\xi+1}^{\xi+1})) \text{ does not } \mu\text{-split over } h_{\xi+1}(N_{\xi+1}).$$

**Subclaim 5.8.** *We can find  $g \in \text{Aut}(\mathfrak{C}/h_{\xi+1}(M_{\xi+1}^{\xi+1}))$  so that*

- $M_{\xi+2}^{\xi+3}$  is universal over  $g(\bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2}))$  and
- $\text{ga-tp}(g(\bar{h}_{\xi+1}(a_{\xi+1}))/N'_{\xi+1})$  does not  $\mu$ -split over  $h_{\xi+1}(N_{\xi+1})$ .

*Proof of Subclaim 5.8.* First fix  $f \in \text{Aut}(\mathfrak{C}/h_{\xi+1}(N_{\xi+1}))$  mapping  $N'_{\xi+1}$  into  $h_{\xi+1}(M_{\xi+1}^{\xi+1})$ . An application of  $f^{-1}$  to  $(*)$  along with monotonicity of non-splitting tell us that

$$(*)_{f^{-1}} \quad \text{ga-tp}(f^{-1}(\bar{h}_{\xi+1}(a_{\xi+1}))/N'_{\xi+1}) \text{ does not } \mu\text{-split over } h_{\xi+1}(N_{\xi+1}).$$

Furthermore, by the definition of non-splitting,  $(*)_{f^{-1}}$  implies that

$$(\sharp) \quad \text{ga-tp}(f^{-1}(\bar{h}_{\xi+1}(a_{\xi+1}))/h_{\xi+1}(M_{\xi+1}^{\xi+1})) = \text{ga-tp}(\bar{h}_{\xi+1}(a_{\xi+1})/h_{\xi+1}(M_{\xi+1}^{\xi+1})).$$

To see this, in the definition of splitting, take  $N_1 := N'_{\xi+1}$  and  $N_2 := h_{\xi+1}(M_{\xi+1}^{\xi+1})$  with the mapping  $f$ .

Using the equality of types  $(\sharp)$ , we can find  $g_1 \in \text{Aut}(\mathfrak{C}/h_{\xi+1}(M_{\xi+1}^{\xi+1}))$  such that  $g_1(f^{-1}(\bar{h}_{\xi+1}(a_{\xi+1}))) = \bar{h}_{\xi+1}(a_{\xi+1})$ . Notice that the inverse  $g_1^{-1}$  would satisfy the second requirement of our subclaim through  $(*)_{f^{-1}}$ , as  $g^{-1}(h_{\xi+1}(a_{\xi+1})) = f^{-1}(h_{\xi+1}(a_{\xi+1}))$ :

$$(*)_{g_1^{-1}} \quad \text{ga-tp}(g_1^{-1}(\bar{h}_{\xi+1}(a_{\xi+1}))/N'_{\xi+1}) \text{ does not } \mu\text{-split over } h_{\xi+1}(N_{\xi+1}).$$

However,  $g_1^{-1}$  may not necessarily satisfy the second requirement of the subclaim. To satisfy the first we requirement, we perform a sequence of mappings.

The next mapping we find is  $g_2$  which takes  $g_1^{-1}(\bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2}))$  into  $M_{\xi+2}^{\xi+3}$  over  $M_{\xi+1}^{\xi+2}$  in such a way that

$$(\dagger) \quad M_{\xi+2}^{\xi+3} \text{ is universal over } g_2 \circ g_1^{-1}(\bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2})).$$

This is possible since  $M_{\xi+2}^{\xi+3}$  is universal over  $M_{\xi+1}^{\xi+2}$  which contains  $h_{\xi+1}(M_{\xi+1}^{\xi+1})$ . We are not yet done, since  $g_2 \circ g_1^{-1}$  does not necessarily satisfy the first requirement of the subclaim. However, we have not yet exhausted all of our induction hypotheses.

We consider one final mapping

$$g_3 : g_2(g_1^{-1}(N'_{\xi+1})) \cong_{g_2 \circ g_1^{-1}(\bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2}))} N'_{\xi+1}.$$

This isomorphism exists because both  $g_2(g_1^{-1}(N'_{\xi+1}))$  and  $N'_{\xi+1}$  are limit models of the same cofinality over  $g_2 \circ g_1^{-1}(\bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2}))$ . To see why this is true, recall the statement  $(\Delta)$  that  $N'_{\xi+1}$  is a limit model over  $\bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2})$  by our original selection of  $\bar{h}_{\xi+1}$ . Then by invariance and an application of  $g_2 \circ g_1^{-1}$  to  $(\Delta)$ , we conclude that  $g_2(g_1^{-1}(N'_{\xi+1}))$  is a limit model of the same cofinality as  $N'_{\xi+1}$  over  $g_2 \circ g_1^{-1}(\bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2}))$ . Notice that because  $N'_{\xi+1}$  was selected to be a limit model over  $\bigcup_{i < \delta} M_i^i$ , by  $(\dagger)$  it is also a limit model (of the same cofinality) over  $g_2 \circ g_1^{-1}(\bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2}))$ .

Now notice that  $g := g_3 \circ g_2 \circ g_1^{-1}$  satisfies the first requirement of the subclaim because of  $(\dagger)$  and the fact that  $g_3$  fixes  $g_2 \circ g_1^{-1}(\bar{h}_{\xi+1}(M_{\xi+2}^{\xi+2}))$ . On the other hand, invariance of non-splitting and an application of  $g_3 \circ g_2$  to  $(*)_{g_1^{-1}}$  gives us

$$(*)_g \quad \text{ga-tp}(g(\bar{h}_{\xi+1}(a_{\xi+1}))/N'_{\xi+1}) \text{ does not } \mu\text{-split over } h_{\xi+1}(N_{\xi+1})$$

because  $g(N'_{\xi+1}) = N'_{\xi+1}$  and  $g \upharpoonright h_{\xi+1}(N_{\xi+1})$  is the identity mapping.

This completes the proof of Subclaim 5.8.  $\dashv$

Let  $h_{\xi+2} := g \circ \bar{h}_{\xi+1} \upharpoonright M_{\xi+2}^{\xi+2}$ . All that is required now is to choose  $N'_{\xi+2}$  so that it is a limit model over  $N'_{\xi+1}$ .

This concludes the construction; we now argue that the existence of the sequences  $\langle N'_i \mid i < \delta \rangle$  and  $\langle h_i \mid i < \delta \rangle$  is enough to find a  $<$ -extension,  $(\bar{M}', \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, \delta}^*$ , of  $(\bar{M}, \bar{a}, \bar{N}) \upharpoonright \delta$  such that  $b \in M'_\zeta$  for some  $\zeta < \delta$ .

Let  $h_\delta := \bigcup_{i < \delta} h_i$ . Note that  $\text{dom}(h_\delta) = \bigcup_{i < \delta} M_i^i$ . We will be defining for  $i < \delta$ ,  $M'_i$  to be pre-image of  $N'_i$  under some extension of  $h_\delta$ . The following claim allows us to choose the pre-image so that  $M'_\zeta$  contains  $b$  for some  $\zeta < \delta$ .

**Claim 5.9.** *There exists  $h \in \text{Aut}(\mathfrak{C})$  extending  $h_\delta$  such that  $h(b) = b$ .*

*Proof of Claim 5.9.* From our choice of  $\xi$ , we know that  $\text{ga-tp}(b/\bigcup_{i < \delta} M_i^i)$  does not  $\mu$ -split over  $M_\xi^\xi$ . Extend  $h_\delta$  to an automorphism  $h^*$  of  $\mathfrak{C}$ . We will show:

**Subclaim 5.10.**  $\text{ga-tp}(b/h^*(\bigcup_{i < \delta} M_i^i)) = \text{ga-tp}(h^*(b)/h^*(\bigcup_{i < \delta} M_i^i))$ .

*Proof of Subclaim 5.10.* We will use non-splitting to derive the subclaim. Following notation used in the definition of splitting, let  $N^1 = \bigcup_{i < \delta} M_i^i$ ,  $N^2 = h^*(\bigcup_{i < \delta} M_i^i)$  and  $p = \text{ga-tp}(b/N^1)$ . Since  $h_i(M_i^i) \prec_{\mathcal{K}} M_i^{i+1} \prec_{\mathcal{K}} \bigcup_{j < \delta} M_j^j$ , we have  $N^2 \prec_{\mathcal{K}} N^1$ . Since  $p$  does not  $\mu$ -split over  $M_\xi^\xi$ , we have that  $p \upharpoonright N^2 = h^*(p \upharpoonright N^1)$ . In other words,

$$\text{ga-tp}(b/h^*(\bigcup_{i < \delta} M_i^i), \mathfrak{C}) = \text{ga-tp}(h^*(b)/h^*(\bigcup_{i < \delta} M_i^i), \mathfrak{C}),$$

as desired. This ends the proof of Subclaim 5.10.  $\dashv$

From Subclaim 5.10, we can find an automorphism  $f$  of  $\mathfrak{C}$  such that  $f(h^*(b)) = b$  and  $f \upharpoonright h^*(\bigcup_{i < \delta} M_i^i) = \text{id}_{h^*(\bigcup_{i < \delta} M_i^i)}$ . Notice that  $h := f \circ h^*$  satisfies the conditions of Claim 5.9:  $h(b) = f(h^*(b)) = b$  and  $h \supset h_\delta$  as  $f \upharpoonright h^*(\bigcup_{i < \delta} M_i^i) = \text{id}_{h^*(\bigcup_{i < \delta} M_i^i)}$ . This ends the proof of Claim 5.9.  $\dashv$

Now that we have an automorphism  $h$  fixing  $b$ , we can define for each  $i < \delta$ ,  $M'_i := h^{-1}(N'_i)$ .

**Claim 5.11.**  $(\bar{M}', \bar{a}, \bar{N})$  is a  $<$ -extension of  $(\bar{M}, \bar{a}, \bar{N})$  such that  $b \in M'_{\xi+1}$ .

*Proof of Claim 5.11.* By construction  $b \in N'_{\xi+1}$ . Since  $h(b) = b$ , this implies  $b \in M'_{\xi+1}$ . To verify that we have a  $\leq$ -extension we need to show for  $i < \delta$ :

- i.  $M'_i$  is universal over  $M_i$
- ii.  $a_i \in M'_{i+1} \setminus M_i$  for  $i+1 < \delta$  and
- iii.  $\text{ga-tp}(a_i/M'_i)$  does not  $\mu$ -split over  $N_i$  whenever  $i, i+1 \leq \delta$ .

Item i. follows from the fact that  $M_i^i$  is universal over  $M_i$  and  $M_i^i \prec_{\mathcal{K}} M'_i$ .

Item iii. follows from invariance and our construction of the  $N'_i$ 's. Finally,

recalling that a non-splitting extension of a non-algebraic type is also non-algebraic, we see that Item iii. implies  $a_i \notin M'_i$ . By our choice of  $h_{i+1}(a_i) \in M_{i+1}^{i+2} \prec_{\mathcal{K}} N'_{i+1}$ , we have that  $a_i \in M'_{i+1}$ . Thus Item ii. is satisfied as well. This ends the proof of Claim 5.11.  $\dashv$

With this, we also end the proof of Claim 5.7.  $\dashv$

... and also, we end the proof of Theorem 6.  $\dashv$

**Corollary 5.12.** *In Theorem 5, we can choose  $(\bar{M}, \bar{a}, \bar{N})$  to be reduced, and hence continuous.*

Now we return to the construction in the proof of the Main Theorem.

$$(\bar{M}, \bar{a}, \bar{N})^0 \quad i_0 \quad i_1 \quad i_\alpha \quad i_{\alpha+1} \quad (\theta \times (\omega + 1)\text{-towers})$$

$$(\bar{M}, \bar{a}, \bar{N})^1$$

$$(\bar{M}, \bar{a}, \bar{N})^n$$

$$(\bar{M}, \bar{a}, \bar{N})^{n+1}$$

$$M^*$$

$$M_{i_\alpha}^n$$

$$M_{i_{\alpha+1}}^n$$

$$\mu \cdot (n + 1)$$

$$M_{s(i_\alpha)}^{n+1}$$

$$M_{i_\alpha}^{n+1}$$

$$\mu \cdot (n + 1) \quad \mu$$

$$M_{i_{\alpha+1}}^{n+1}$$

$$M_{i_\alpha}^{n+2}$$

$$M_{i_{\alpha+1}}^{n+2}$$

Corollary 5.12 tells us that the construction of our array of models as an increasing sequence of towers is possible in successor cases. In the limit case,



let  $I_\omega = \bigcup_{m < \omega} I_m$ , and simply define,  $(\bar{M}, \bar{a}, \bar{N})^\omega \in \mathcal{K}_{\mu, \mathbf{I}_\omega}^*$  to be the union of the towers  $(\bar{M}, \bar{a}, \bar{N})^n$ .

To see that the construction satisfies our requirements, first notice that the last column of the array,  $\langle M_{i_\theta}^n \mid n < \omega \rangle$ , witnesses that  $M^*$  is a  $(\mu, \omega)$ -limit model. In light of Theorem 4 we need only verify that the last row of the array is a relatively full tower of cofinality  $\theta$ .

**Claim 5.13.**  $(\bar{M}, \bar{a}, \bar{N})^\omega$  is full relative to  $(M_i^n)_{n < \omega, i \in I_\omega}$ .

*Proof.* Given  $i$  with  $i_\alpha \leq i < i_{\alpha+1}$ , let  $(p, M_i^n)$  be some strong type in  $\mathfrak{St}(M_i^\omega)$ . Notice that by monotonicity of non-splitting  $(p \upharpoonright M_i^{n+1}, M_i^n) \in \mathfrak{St}(M_i^{n+1})$ . By construction there is a  $j \in I_{n+1}$  with  $i < j < i_{\alpha+1}$  such that  $(\text{ga-tp}(a_j/M_j^{n+2}), N_j^{n+2})$  is parallel to  $p \upharpoonright M_i^{n+1}$ . We will show that  $(\text{ga-tp}(a_j/M_j^\omega), N_j^\omega)$  is parallel to  $(p, N)$ .

First notice that  $\text{ga-tp}(a_j/M_j^\omega)$  does not  $\mu$ -split over  $N_j^\omega = N_j^{n+2}$  because  $(\bar{M}, \bar{a}, \bar{N})^\omega$  is a tower. Since  $(\text{ga-tp}(a_j/M_j^{n+2}), N_j^{n+2})$  is parallel to  $(p \upharpoonright M_i^{n+1}, M_i^n)$  there is  $q \in \text{ga-S}(M_j^\omega)$  such that  $q$  extends both  $p \upharpoonright M_i^{n+1}$  and  $\text{ga-tp}(a_j/M_j^{n+2})$ . By two separate applications of the uniqueness of non- $\mu$ -splitting extensions we know that  $q \upharpoonright M_i^\omega = p$  and  $q = \text{ga-tp}(a_j/M_j^\omega)$ . To see that  $(q, N_j^\omega)$  is parallel to  $(p, M_i^n)$ , let  $M'$  be an extension of  $M_j^\omega$  of cardinality  $\mu$ . Since  $(p \upharpoonright M_i^{n+1}, M_i^n)$  and  $(q \upharpoonright M_j^{n+2}, N_j^{n+2})$  are parallel, there is  $q' \in \text{ga-S}(M')$  extending both  $p \upharpoonright M_i^{n+1}$  and  $q \upharpoonright M_j^{n+2}$  and not  $\mu$ -splitting over both  $M_i^n$  and  $N_j^{n+2}$ . By the uniqueness of non- $\mu$ -splitting extensions, we have that  $q'$  is also an extension of  $q$  and  $p$ . Thus  $q'$  witnesses that  $(q, N_j^\omega)$  and  $(p, M_i^n)$  are parallel.  $\dashv$

This completes the proof of Theorem 3.

## 6. CONCLUDING REMARKS

In this section we discuss other results related to Question 1.9. First to understand the boundaries of Question 1.9, consider the elementary case. Limit models are not necessarily unique even for first order complete stable theories.

**Theorem 7.** Suppose  $T$  is a complete, stable theory. Let  $\mu \geq 2^{|T|}$  such that  $\mu^{|T|} = \mu$ . If  $T$  is not superstable, then no  $(\mu, \omega)$ -limit model is isomorphic to any  $(\mu, \kappa)$ -limit model for any  $\kappa$  with  $\text{cf}(\kappa) \geq \kappa(T)$ .

*Proof.* Let  $T$  be a stable, but not superstable, complete theory, and fix  $\kappa$  and  $\mu$  as in the statement of the theorem. As  $T$  is not superstable,

by [Sh e, Lemma VII, 3.5 (2)], for  $\lambda := (2^\mu)^+$ , there are  $\langle \bar{a}_\eta | \eta \in {}^\omega \geq \lambda \rangle$  and  $\langle \varphi_n(\bar{x}, \bar{y}_n) | n < \omega \rangle$  such that for every  $n < \omega$  and all  $\eta \in {}^\omega \lambda$ ,

$$(\mathfrak{C} \models \varphi_n[\bar{a}_\eta, \bar{a}_\nu]) \iff \nu = \eta \upharpoonright n.$$

By induction on  $n < \omega$  define  $\langle M_n | n < \omega \rangle$  all of cardinality  $\mu$  and  $\langle \eta_n, \nu_n | n < \omega \rangle$  such that

- (1)  $M_{n+1}$  is universal over  $M_n$  and saturated of cardinality  $\mu$ ,
- (2)  $\eta_{n+1} > \eta_n$ ,  $\nu_{n+1} > \eta_n$ , and  $\eta_{n+1} \neq \nu_{n+1}$ ,
- (3)  $\bar{a}_{\eta_{n+1}}, \bar{a}_{\nu_{n+1}} \in M_{n+1}$  and
- (4)  $\text{tp}(\bar{a}_{\eta_{n+1}}/M_n) = \text{tp}(\bar{a}_{\nu_{n+1}}/M_n)$ .

THIS CONSTRUCTION IS ENOUGH: Let  $N' \models T$  be a  $(\mu, \kappa)$ -limit over  $M_0$ . By Theorem 2,  $N'$  must be saturated. Let  $N = \bigcup_{n < \omega} M_n$ . Clearly  $N$  is a  $(\mu, \omega)$ -limit over  $M_0$ . To conclude that  $N$  and  $N'$  are non-isomorphic, it is enough to show that  $N$  is not saturated. Consider  $p := \{\varphi_{n+1}(\bar{x}; \bar{a}_{\eta_{n+1}}) \wedge \neg \varphi_{n+1}(\bar{x}; \bar{a}_{\nu_{n+1}}) | n < \omega\}$ . The set of formulas  $p$  is a type since it is realized in  $\mathfrak{C}$  by  $\bar{a}_\eta$  where  $\eta := \bigcup_{n < \omega} \eta_n$ . Notice that  $N$  cannot satisfy  $p$ . If  $\bar{a} \in N$  would satisfy  $p$ , then  $M_n$  realizes  $p$  for some  $n < \omega$ . Thus by condition (4), we would have

$$\mathfrak{C} \models \varphi_{n+1}[\bar{a}, \bar{a}_{\eta_{n+1}}] \iff \mathfrak{C} \models \varphi_{n+1}[\bar{a}, \bar{a}_{\nu_{n+1}}]$$

which would contradict the assumption that  $\bar{a}$  satisfies  $p$ .

THIS IS POSSIBLE: By stability and  $\mu^{|T|} = \mu$ , using the proof of [Sh e, Th. III 3.12], every model of cardinality  $\mu$  has a saturated proper elementary extension. Let  $M_0$  be a saturated model of cardinality  $\mu$  and take  $\eta_0 = \nu_0 := \langle \rangle$ . Given  $\eta_n, \nu_n, M_n$ , using Theorem 1 let  $M^*$  be universal over  $M_n$  of cardinality  $\mu$ . Let  $M^{**} \succ M^*$  of cardinality  $\mu$  containing  $\bar{a}_{\eta_n}$  and  $\bar{a}_{\nu_n}$ . By [Sh e, Th. III 3.12], we can take  $M_{n+1} \succ M^{**}$  saturated of cardinality  $\mu$ . Clearly it is universal over  $M_n$ . For  $n < \omega$ , consider  $F_n(\alpha) := \text{tp}(\bar{a}_{\eta_n \hat{\ } \alpha}/M_n)$ . As  $\lambda$  is regular and  $\lambda > |S(M_n)|$ , there is  $S \subset \lambda$  of cardinality  $\lambda$  such that  $\alpha \neq \beta \in S \Rightarrow F_n(\alpha) \neq F_n(\beta)$ . Pick  $\alpha \neq \beta \in S$  and define  $\eta_{n+1} := \eta_n \hat{\ } \alpha$  and  $\nu_{n+1} := \eta_n \hat{\ } \beta$ .  $\dashv$

In the non-elementary setting, many authors have considered approximations to Theorem 3. Several authors have proved and used the uniqueness of limit models in AECs under the assumption of categoricity: [Sh 394] [Ba], [KoSh], [Sh 576], [ShVi] and [Va]. Also, Shelah's [Sh i] examines (as an aside) the uniqueness of limit models in good frames. These are abstract elementary classes under much stronger assumptions than we make here.

In Theorem 6.5 of [Sh 394], Shelah claims uniqueness of limit models of cardinality  $\mu$  for classes with the amalgamation property under little more than categoricity in some  $\lambda > \mu > \text{LS}(\mathcal{K})$  together with existence of arbitrarily large models. Shelah's claim in Theorem 6.5 of [Sh 394] (isomorphism over the base) seems too strong for the proof that he suggests. Instead, he proves that  $(\mu, \kappa)$ -limit models are Galois saturated. The argument in [Sh 394] depends in a crucial way on an analysis of Ehrenfeucht-Mostowski models. In our paper, we cannot employ Ehrenfeucht-Mostowski machinery because we do not assume here categoricity or the existence of models above the Hanf number.

Under similar assumptions as those in [Sh 394], more recently, Baldwin in [Ba] (Chapter 11) has used methods based on [Sh 394] to prove that if  $M_1$  and  $M_2$  are  $(\mu, \sigma_1)$ - and  $(\mu, \sigma_2)$ -limit models over  $N$ , respectively, then  $M_1 \cong M_2$ . Baldwin, however, does not prove that  $M_1$  and  $M_2$  are isomorphic *over*  $N$ . Our result is therefore much stronger than that in [Ba].

Kolman and Shelah in [KoSh] prove the uniqueness of limit models of cardinality  $\mu$  in  $\lambda$ -categorical AECs that are axiomatized by a  $L_{\kappa, \omega}$ -sentence where  $\lambda > \mu$  and  $\kappa$  is a measurable cardinal. Then Kolman and Shelah use this uniqueness result to prove that amalgamation occurs below the categoricity cardinal in  $L_{\kappa, \omega}$ -theories with  $\kappa$  measurable. Both the measurability of  $\kappa$  and the categoricity are used integrally in their proof of uniqueness.

Shelah in [Sh 576] (see Claim 7.8) proved a special case of the uniqueness of limit models under the assumption of  $\mu$ -AP, categoricity in  $\mu$  and in  $\mu^+$  as well as assuming  $K_{\mu^{++}} \neq \emptyset$ . In that paper Shelah needs to produce *reduced types* and use some of their special properties.

In [ShVi], Shelah and Villaveces attempted to prove a uniqueness theorem without assuming any form of amalgamation; however, they assumed that  $\mathcal{K}$  is categorical in some sufficiently large  $\lambda$ , that every model in  $\mathcal{K}$  has a proper extension and that  $2^\lambda < 2^{\lambda^+}$ . VanDieren in [Va] managed to prove the uniqueness statement under the assumptions of [ShVi] together with the additional assumption that  $\mathcal{K}^{am} := \{M \in \mathcal{K}_\mu \mid M \text{ is an amalgamation base}\}$  is closed under unions of increasing  $\prec_{\mathcal{K}}$  chains.

In [Sh i] the most important new concept is that of a  $\lambda$ -good frame, which is an axiomatization of the notion of superstability, *with hypothesis on just one cardinal*  $\lambda$ . Its full definition is more than a page long. Shelah's assumptions on the AEC include, among other things, the amalgamation property, the existence of a forking like dependence relation and of a family of types

playing a role akin to that of regular types in first order superstable theories – Shelah calls them *bs*-types. One of the axioms of a good frame is the existence of a non-maximal super-limit model. This axiom along with  $\mu$ -stability implies the uniqueness of limit models of cardinality  $\mu$ . In Lemma II.4.8 of [Sh i] he states that in a good frame, limit models are unique. (While we don't claim that we understand Shelah's proof or believe in its correctness, he explicitly uses the interplay between *bs*-types and the forking notion as well as no long forking chains and continuity of forking.)

The formal differences between our approach and Shelah's [Sh i] can be summarized as follows:

- (1) Suppose that  $\mathcal{K}$  is an AEC with no maximal models satisfying the disjoint amalgamation property over limit models and is categorical in  $\lambda^+$  for some  $\lambda > \text{LS}(\mathcal{K})$ ; we then get uniqueness of limit models. By way of comparison, in order to get a uniqueness of limit models, Shelah needs results of [Sh 576] (a 99 pages-long paper) and significant parts of his book [Sh i] along with the stronger assumptions of categoricity in several consecutive cardinals together with several additional set-theoretic axioms. All our results are in ZFC.
- (2) When specialized to the case when  $\mathcal{K}$  is the class of models of a complete first order theory  $T$ , Shelah's proof in [Sh i, Lemma II.4.8] really uses the full power of assuming that  $T$  is *superstable*. The proof of uniqueness in this paper just needs, in addition to the stability of  $T$ , no splitting chains of length  $\omega$ . As the main interest of our theorem is for the general case of AEC, rather than just for first order theories, the difference between this paper and [Sh i, Lemma II.4.8] is clearer when understood in light of the greater picture.

We are particularly interested in Theorem 3 not only for the sake of generalizing Shelah's result from [Sh 576] but due to the fact that the first and second author use this uniqueness theorem along with tools from [Sh 394] in a crucial step to prove:

**Theorem 8** (Upward categoricity theorem, [GrVa2]<sup>4</sup>). *Suppose that  $\mathcal{K}$  has arbitrarily large models, is  $\chi$ -tame and satisfies the amalgamation and joint embedding properties. Let  $\lambda$  be such that  $\lambda > \text{LS}(\mathcal{K})$  and  $\lambda \geq \chi$ . If  $\mathcal{K}$  is categorical in  $\lambda^+$  then  $\mathcal{K}$  is categorical in all  $\mu \geq \lambda^+$ .*

<sup>4</sup>Some time after Grossberg and VanDieren announced Theorem 8, Baldwin circulated an alternative proof of Theorem 8 that eventually appeared in [Ba]. Lessmann in [Les05] proved the result for  $\mathcal{K}$  with  $\text{LS}(\mathcal{K}) = \aleph_0$  beginning with categoricity in  $\aleph_1$ .

Grossberg and VanDieren’s use of the uniqueness of limit models in this theorem hints at a connection between classical definitions of superstability in first order logic and the uniqueness of limit models. This link is explored in further work of VanDieren.

## REFERENCES

- [Ba] Baldwin, John T. **Categoricity**. University Lecture Series. American Mathematical Society 50 (2009). Print.
- [Gr1] Grossberg, Rami. **A Course in Model Theory I**, to be published by Cambridge University Press.
- [Gr2] Grossberg, Rami. “Classification Theory for Non-elementary Classes.” *Contemporary Mathematics*, **302** (2002): 165–204. Print.
- [GrLe] Grossberg, Rami and Olivier Lessmann. “Shelah’s Stability Spectrum and Homogeneity Spectrum.” *Archive for mathematical Logic*, **41.1** (2002): 1–31. Print.
- [GrVa0] Grossberg, Rami and Monica VanDieren. “Shelah’s Categoricity Conjecture from a Successor for Tame Abstract Elementary Classes.” *Journal of Symbolic Logic*. **71.2** (2006): 553–568. Print.
- [GrVa1] Grossberg, Rami and Monica VanDieren. “Galois-stability for Tame Abstract Elementary Classes.” *Journal of Mathematical Logic* **6.1** (2006): 25–49. Print.
- [GrVa2] Grossberg, Rami and Monica VanDieren. “Categoricity from one successor cardinal in Tame Abstract Elementary Classes.” *Journal of Mathematical Logic* **6.2** (2006): 181–201. Print.
- [HyKe] Hyttinen, Tapani and Meeri Kesälä. “Independence in Finitary Abstract Elementary Classes.” *Annals of Pure and Applied Logic* **143** (2006): 103–138. Print.
- [Jo] Jónsson, Bjarni. “Homogeneous universal systems.” *Math. Scand.* **8** (1960): 137–142. Print.
- [KoSh] Kolman, Oren and Saharon Shelah. “Categoricity of Theories in  $L_{\kappa, \omega}$  when  $\kappa$  is a Measurable Cardinal. Part I.” *Fundamentae Mathematicae* **151** (1996): 209–240. Print.
- [Les05] Lessmann, Olivier. “Upward Categoricity from a Successor Cardinal for an Abstract Elementary Class with Amalgamation.” *Journal of Symbolic Logic* **70** (2005): 639–661. Print.
- [Po] Poizat, Bruno. **A course in model theory**. Springer. (2000). Print.
- [Sh e] Shelah, Saharon. **Classification Theory**. Revised edition. North Holland (1990). Print.
- [Sh i] —. **Classification Theory for Abstract Elementary Classes**. (Studies in Logic: Mathematical Logic and Foundations). College Publications (2009). Print.
- [Sh 3] —. “Finite Diagrams Stable in Power.” *Ann. Math. Logic* **2**, (1970): 69–118. Print.
- [Sh 87b] —. “Classification Theory for Nonelementary Classes. I. The Number of Uncountable Models of  $\psi \in L_{\omega_1, \omega}$ . Part B, *Israel Journal of Mathematics* **46** (1983): 241–273. Print.
- [Sh 394] —. “Categoricity of Abstract Classes with Amalgamation.” *Annals of Pure and Applied Logic* **98** (1999): 261–294. Print.

- [Sh 472] —. “Categoricity of Theories in  $L_{\kappa^*, \omega}$  when  $\kappa^*$  is a Measurable Cardinal. Part II.” Dedicated to the memory of Jerzy Los. *Fundamenta Mathematica* **170** (2001): 165–196. Print.
- [Sh 576] —. “Categoricity of an Abstract Elementary Class in Two Successive Cardinals.” *Israel Journal of Mathematics* **126** (2001): 29–128. Print.
- [Sh 600] —. Categoricity in abstract elementary classes: going up inductive step. Preprint. arXiv:math.LO/0011215 (November 2000 version), 82 pages.
- [Sh 705] —. Toward Classification Theory of Good  $\lambda$  Frames and Abstract Elementary Classes. Preprint.
- [ShVi] Shelah, Saharon and Andrés Villaveces. “Categoricity in abstract elementary classes with no maximal models.” *Annals of Pure and Applied Logic*, **97**.1-3 (1999): 1–25. Print.
- [Va] VanDieren, Monica. “Categoricity in Abstract Elementary Classes with No Maximal Models.” *Annals of Pure and Applied Logic* **141** (2006): 108–147. Print.
- [Za] Zambrano, Pedro. Cats, docilidad, y la propiedad de amalgamación disyunta. Preprint.

*E-mail address*, Rami Grossberg: `rami@cmu.edu`

DEPARTMENT OF MATHEMATICS, CARNEGIE MELLON UNIVERSITY, PITTSBURGH PA 15213

*E-mail address*, Monica VanDieren: `vandieren@rmu.edu`

DEPARTMENT OF MATHEMATICS, ROBERT MORRIS UNIVERSITY, MOON TOWNSHIP PA 15108

*E-mail address*, Andrés Villaveces: `avillavecesn@unal.edu.co`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL DE COLOMBIA, BOGOTÁ, COLOMBIA