

On the game chromatic number of sparse random graphs

Alan Frieze* and Simcha Haber† and Mikhail Lavrov‡

Department of Mathematical Sciences,
Carnegie Mellon University,
Pittsburgh PA, 15213,
USA.

June 8, 2018

Abstract

Given a graph G and an integer k , two players take turns coloring the vertices of G one by one using k colors so that neighboring vertices get different colors. The first player wins iff at the end of the game all the vertices of G are colored. The game chromatic number $\chi_g(G)$ is the minimum k for which the first player has a winning strategy. The paper [6] began the analysis of the asymptotic behavior of this parameter for a random graph $G_{n,p}$. This paper provides some further analysis for graphs with constant average degree i.e. $np = O(1)$ and for random regular graphs. We show that w.h.p. $c_1\chi(G_{n,p}) \leq \chi_g(G_{n,p}) \leq c_2\chi(G_{n,p})$ for some absolute constants $1 < c_1 < c_2$. We also prove that if $G_{n,3}$ denotes a random n -vertex cubic graph then w.h.p. $\chi_g(G_{n,3}) = 4$.

1 Introduction

Let $G = (V, E)$ be a graph and let k be a positive integer. Consider the following game in which two players A(lice) and B(ob) take turns in coloring the vertices of G with k colors. Each move consists of choosing an uncolored vertex of the graph and assigning to it a color from $\{1, \dots, k\}$ so that the resulting coloring is *proper*, i.e., adjacent vertices get different colors. A wins if all the vertices of G are eventually colored. B wins if at some point in the game the current partial coloring cannot be extended to a complete coloring of G , i.e., there is an uncolored vertex such that each of the k colors appears at least once in its neighborhood.

*Research supported in part by NSF Grant CCF1013110. Email: alan@random.math.cmu.edu.

†Email: simi@andrew.cmu.edu.

‡Email: mlavrov@andrew.cmu.edu.

We assume that A goes first (our results will not be sensitive to this choice). The *game chromatic number* $\chi_g(G)$ is the least integer k for which A has a winning strategy.

This parameter is well defined, since it is easy to see that A always wins if the number of colors is larger than the maximum degree of G . Clearly, $\chi_g(G)$ is at least as large as the ordinary chromatic number $\chi(G)$, but it can be considerably more. The game was first considered by Brams about 25 years ago in the context of coloring planar graphs and was described in Martin Gardner's column [12] in Scientific American in 1981. The game remained unnoticed by the graph-theoretic community until Bodlaender [5] re-invented it. For a survey see Bartnicki, Grytczuk, Kierstead and Zhu [4].

In this paper, we study the game chromatic number of the random graph $G_{n,p}$ and the random d -regular graph $G_{n,d}$. Define $b = \frac{1}{1-p}$. The following estimates were proved in Bohman, Frieze and Sudakov [6].

Theorem 1.1.

(a) *There exists $K > 0$ such that for $\varepsilon > 0$ and $p \geq (\ln n)^{K\varepsilon^{-3}}/n$ we have that w.h.p.¹*

$$\chi_g(G_{n,p}) \geq (1 - \varepsilon) \frac{n}{\log_b np}.$$

(b) *If $\alpha > 2$ is a constant, $K = \max\{\frac{2\alpha}{\alpha-1}, \frac{\alpha}{\alpha-2}\}$ and $p \geq (\ln n)^K/n$ then w.h.p.*

$$\chi_g(G_{n,p}) \leq \alpha \frac{n}{\log_b np}.$$

In this paper we complement these results by considering the case where $p = \frac{d}{n}$ where d is at least some sufficiently large constant. We will assume that $d \leq n^{1/4}$ since Theorem 1.1 covers larger d .

Theorem 1.2. *Let $p = \frac{d}{n}$ where d is larger than some absolute constant and $d \leq n^{1/4}$.*

(a) *If $\alpha < \frac{4}{7}$ is a constant then w.h.p.*

$$\chi_g(G_{n,p}) \geq \frac{\alpha d}{\ln d}.$$

(b) *If α is a sufficiently large constant then w.h.p.*

$$\chi_g(G_{n,p}) \leq \frac{\alpha d}{\ln d}.$$

Note that when $p = o(1)$ we have $\frac{n}{\log_b np} \sim \frac{d}{\ln d}$. Note also that the bounds in Theorem 1.1 are stronger than those in Theorem 1.2, whenever both results are applicable.

¹A sequence of events \mathcal{E}_n occurs *with high probability* (w.h.p.) if $\lim_{n \rightarrow \infty} \mathbb{P}(\mathcal{E}_n) = 1$

It is natural to compare our bounds with the asymptotic behavior of the ordinary chromatic number of random graph. It is known by the results of Bollobás [7] and Łuczak [16] that when $p = o(1)$, $\chi(G_{n,p}) = (1 + o(1))\frac{d}{2\ln d}$ w.h.p. (Of course a stronger result is now known, see Achlioptas and Naor [2]). Thus Theorem 1.2 shows that the game chromatic number of $G_{n,p}$ is at most (roughly) twelve times and at least (roughly) 8/7 times its chromatic number.

Having proved Theorem 1.2, we extend the results to the random d -regular graph $G_{n,d}$.

Theorem 1.3. *Let $\varepsilon > 0$ be an arbitrary constant.*

(a) *If α is a constant satisfying the conditions of Theorem 1.1 or Theorem 1.2 where appropriate and d is sufficiently large and $d \leq n^{1/3-\varepsilon}$ then w.h.p.*

$$\chi_g(G_{n,d}) \geq \frac{\alpha d}{\ln d}.$$

(b) *If α is a constant satisfying the conditions of Theorem 1.1 or Theorem 1.2 where appropriate and d is sufficiently large and $d \leq n^{1/3-\varepsilon}$ then w.h.p.*

$$\chi_g(G_{n,d}) \leq \frac{\alpha d}{\ln d}.$$

It is known by the result of Frieze and Łuczak [11] that w.h.p. $\chi(G_{n,d}) = (1 + o(1))\frac{d}{2\ln d}$. (Of course stronger results are now known, see Achlioptas and Moore [1] and Kemkes, Pères-Giménez and Wormald [14]).

Theorem 1.3 says nothing about $\chi_g(G_{n,d})$ when d is small. We have been able to prove

Theorem 1.4. *If $d = 3$ then w.h.p. $\chi(G_{n,d}) = 4$.*

It is easy to see via Brooks' theorem that w.h.p. the chromatic number of a random cubic graph is three and so Theorem 1.4 separates χ and χ_g in this context.

We often refer to the following Chernoff-type bounds for the tails of binomial distributions (see, e.g., [3] or [13]). Let $X = \sum_{i=1}^n X_i$ be a sum of independent indicator random variables such that $\mathbb{P}(X_i = 1) = p_i$ and let $p = (p_1 + \dots + p_n)/n$. Then

$$\mathbb{P}(X \leq (1 - \varepsilon)np) \leq e^{-\varepsilon^2 np/2}, \tag{1.1}$$

$$\mathbb{P}(X \geq (1 + \varepsilon)np) \leq e^{-\varepsilon^2 np/3}, \quad \varepsilon \leq 1, \tag{1.2}$$

$$\mathbb{P}(X \geq \mu np) \leq (e/\mu)^{\mu np}. \tag{1.3}$$

1.1 Outline of the paper

Section 2 is devoted to the proof of Theorem 1.2. In Section 2.1, we prove a lower bound on $\chi_g(G_{n,p})$ by giving a strategy for player B. Basically, B's strategy is to follow A coloring a vertex with color i by coloring a random vertex v with color i . Of course we mean here

that v is randomly chosen from vertices outside of the neighborhood of the set of vertices of color i . Why does this work? Well, it is known that choosing an independent set via a greedy algorithm will w.h.p. find an independent set that is about one half the size of the largest independent set. What we show is that choosing randomly half the time also has a deleterious effect on the size of the independent set (color class) selected. This leads to the game chromatic number being significantly larger than the chromatic number.

In Section 2.2, we prove an upper bound on $\chi_g(G_{n,p})$ by giving a strategy for player A. Here A follows the same strategy used in the proof of Theorem 1.1(b), up until close to the end. We then let A follow a more sophisticated strategy. A's initial strategy is to choose a vertex with as few "available" colors and color it with any available color i.e. one that does not conflict with its colored neighbors. At a certain point there are few uncolored vertices and they all have a substantial number of available colors. We show that the edges of the graph induced by these vertices can be partitioned into a forest F plus a low degree subgraph. Using the tree coloring strategy described in [10] we see that the low degree subgraph does not prevent G from being colored.

Having proved Theorem 1.2 we transfer the results to random d -regular graphs ($d \leq n^{1/4}$) by showing that the underlying structural lemmas remain true or trivially modified. This is done in Section 3.

In Section 3 we show how to convert Theorem 1.1 into a random regular graph setting. The two ranges $d_0 \leq d \leq n^{1/4}$ and $n^{1/4} < d \leq n^{1/3-\varepsilon}$ are treated separately. The lower range is treated in Section 3.1 and the upper range is treated in Section 3.2 using the "Sandwiching Theorem" of Kim and Vu [15].

In Section 4 we provide a strategy for B showing that w.h.p. $\chi_g(G_{n,3}) = 4$. This proves Theorem 1.4. B's strategy is based on his ability to force A into playing on a small set of vertices. B will then make a sequence of such forcing moves along a cycle to create a double threat and win the game.

2 Theorem 1.2: $G_{n,p}, p = d/n$

2.1 The lower bound

Let $D = \frac{d}{\ln d}$ and suppose that there are $k = \alpha D$ colors. At any stage, let C_i be the set of vertices that have been colored i and let $C = \bigcup_{i=1}^k C_i$. Let $U = [n] \setminus C$ be the set of uncolored vertices and let $U_i = U \setminus N(C_i)$. Note that $[n] = \{1, 2, \dots, n\}$ is the vertex set of $G_{n,p}$.

B's strategy will be to choose the same color i that A just chose and then to assign color i to a random vertex in U_i . The idea being that making random choices when constructing an independent set (color class) tends to only get one of half the maximum size. A could be making better choices and so we do not manage to prove that we need twice as many colors

as the chromatic number.

Suppose that we run this process for θn rounds and that $|C_i| = 2\beta n/D$ where we will later take $\theta = 7\alpha/8 < 1/2$ and $\beta = 1/2$. Let S_i be the set of $\beta n/D$ vertices in C_i that were colored by B. We consider the probability that there exists a set T of size $\gamma n/D$ such that $C_i \cup T$ is independent. For expressions X, Y we use the notation $X \leq_b Y$ in place of $X = O(Y)$ when the bracketing is “ugly”.

$$\mathbb{P}(\exists C_i, T) \leq_b k \binom{n}{\beta n/D} \binom{n}{\gamma n/D} \sum_{|S|=\beta n/D} \mathbb{P}(S_i = S) (1-p)^{(2\beta+\gamma)^2 n^2 / 2D^2} \quad (2.1)$$

$$\leq k \binom{n}{\beta n/D} \binom{n}{\gamma n/D} \sum_{|S|=\beta n/D} (\beta n/D)! \prod_{j=1}^{\beta n/D} \frac{7}{(1-p)^{2j} (1-2\theta)n} (1-p)^{(2\beta+\gamma)^2 n^2 / 2D^2} \quad (2.2)$$

$$\begin{aligned} &\leq k \binom{n}{\beta n/D}^2 \binom{n}{\gamma n/D} \frac{(\beta n/D)!}{((1-2\theta)n)^{\beta n/D}} \frac{7^{\beta n/D}}{(1-p)^{\beta^2 n^2 / D^2}} (1-p)^{(2\beta+\gamma)^2 n^2 / 2D^2} \\ &\leq k \left(\left(\frac{eD}{\beta} \right)^\beta \cdot \left(\frac{7}{1-2\theta} \right)^\beta \cdot \left(\frac{eD}{\gamma} \right)^\gamma \cdot \exp \{ (2\beta^2 - (2\beta + \gamma)^2) d / 2D \} \right)^{n/D} \\ &= k \exp \{ (\beta + \gamma + \beta^2 - (2\beta + \gamma)^2 / 2 + o_d(1)) d^{-1} n \ln^2 d \} \\ &= o(1) \end{aligned} \quad (2.3)$$

if $(2\beta + \gamma)^2 > 2(\beta + \beta^2 + \gamma)$. This is satisfied when $\beta = 1/2$ and $\gamma = 3/4$. We will justify (2.1) and (2.2) momentarily.

If the event $\{\exists C_i, T\}$ does not occur then because no color class has size greater than $(2\beta + \gamma)n/D$ the number ℓ of colors i for which $|S_i| \geq \beta n/D$ by this time satisfies

$$\frac{\ell(2\beta + \gamma)}{D} + \frac{2(k - \ell)\beta}{D} \geq 2\theta.$$

We choose $\theta = 7\alpha/8$. Since $k \geq \ell$, this implies that

$$\frac{k}{D} \geq \frac{2\theta}{2\beta + \gamma} = \alpha.$$

This completes the proof of Part (a) of Theorem 1.2.

Justifying (2.1): Here we are taking the union bound over all $\binom{n}{\beta n/D} \binom{n}{\gamma n/D}$ possible choices of $C_i \setminus S_i$ and T . In some sense we are allowing player A to simultaneously choose all possible sets of size $\beta n/D$ for $C_i \setminus S_i$. The union bound shows that w.h.p. all choices fail. We do not sum over orderings of $C_i \setminus S_i$. We instead compute an upper bound on $\mathbb{P}(S_i = S)$ that holds regardless of the order in which A plays. We consider the situation after θn rounds. That is, we think of the following random process: pick a graph $G \sim G(n, p)$, let Alice play the coloring game on G with k colors against a player who randomly chooses an available vertex to be colored by the same color as Alice. Stop after θn moves. At this point Alice played

with color i and there are $\beta n/D$ vertices that were colored i by Alice and the same number that were colored i by Bob. We bound the probability that at this point there are $\gamma n/D$ vertices that form an independent set with the i 'th color class. We take a union bound over all the possible sets for Alice's vertices and for the vertices in T . The probability of Bob choosing a certain set is computed below.

Justifying (2.2): For this we first consider a sequence of random variables

$$X_1 = N = (1 - 2\theta)n, X_j = \text{Bin}(X_{j-1}, q) \text{ where } q = (1 - p)^2 \text{ and } 1 \leq j \leq t.$$

X_j is a lower bound for the number of vertices Bob can color i . The probability that a vertex was i -available at time $j - 1$ and is still i -available now is $(1 - p)^2$. This is because two more vertices have been colored i . Also, we take $X_1 = N$ as a lower bound on the number of choices at the start of the process. Then we estimate $\mathbb{E}(Y_t)$ where

$$Y_t = \begin{cases} 0 & X_t = 0 \\ \frac{1}{X_1 X_2 \dots X_t} & X_t > 0 \end{cases}$$

We use $Y_{\beta D n}$ as an upper bound on the probability that B's sequence of choices is $x_1, x_2, \dots, x_{\beta n/D}$ where $S = \{x_1, x_2, \dots, x_{\beta n/D}\}$. The term X_j lower bounds the number of choices that B has and so $1/X_j$ upper bounds the probability that B chooses x_j . We take the expectation of the product of these bounds over $G_{n,p}$.

Now if $B = \text{Bin}(\nu, q)$ and we take $\prod_{i=1}^k \frac{1}{B+i-1} = 0$ when $B = 0$ then

$$\begin{aligned} \mathbb{E} \left(\prod_{i=1}^k \frac{1}{B+i-1} \right) &= \sum_{\ell=1}^{\nu} \prod_{i=1}^k \frac{1}{\ell+i-1} \binom{\nu}{\ell} q^{\ell} (1-q)^{\nu-\ell} \\ &= \frac{1}{q^k} \prod_{i=1}^k \frac{1}{\nu+i} \sum_{\ell=1}^{\nu} \frac{\ell+k}{\ell} \binom{\nu+k}{\ell+k} q^{\ell+k} (1-q)^{\nu-\ell} \\ &\leq \left(\frac{1}{q^k} \prod_{i=1}^k \frac{1}{\nu+i} \right) \left(1 + \sum_{\ell=1}^{\nu} \frac{k}{\ell} \binom{\nu+k}{\ell+k} q^{\ell+k} (1-q)^{\nu-\ell} \right). \end{aligned} \quad (2.4)$$

Suppose now that $q = 1 - o(1)$. Then

$$\begin{aligned} \sum_{\ell=1}^{\nu} \frac{k}{\ell} \binom{\nu+k}{\ell+k} q^{\ell+k} (1-q)^{\nu-\ell} &\leq \sum_{\ell=1}^{k/2} \frac{k}{\ell} \binom{\nu+k}{\ell+k} q^{\ell+k} (1-q)^{\nu-\ell} + 2 \sum_{\ell=1}^{\nu} \binom{\nu+k}{\ell+k} q^{\ell+k} (1-q)^{\nu-\ell} \\ &\leq k e^{-(\nu+k)/10} + 2 \\ &\leq 6. \end{aligned}$$

Going back to (2.4) we see that

$$\mathbb{E} \left(\prod_{i=1}^k \frac{1}{B+i-1} \right) \leq \frac{7}{q^k} \prod_{i=1}^k \frac{1}{\nu+i}.$$

It follows that

$$\begin{aligned}
& \mathbb{E} \left(\frac{1}{X_1 \cdots X_t} \right) \\
& \leq \mathbb{E} \left(\frac{7}{X_1 \cdots X_{t-1} (X_{t-1} + 1) q} \right) \\
& \leq \mathbb{E} \left(\frac{7^2}{X_1 \cdots X_{t-2} (X_{t-2} + 1) (X_{t-2} + 2) q^{1+2}} \right) \\
& \quad \vdots \\
& \leq \frac{7^t}{N(N+1) \cdots (N+t) q^{1+2+\cdots+t}}.
\end{aligned}$$

2.2 The upper bound

We begin by proving some simple structural properties of $G_{n,p}$.

Lemma 2.1. *If $\theta > 1$ and*

$$\left(\frac{\sigma ed}{2\theta} \right)^\theta \leq \frac{\sigma}{2e} \tag{2.5}$$

then w.h.p. there does not exist $S \subseteq [n]$, $|S| \leq \sigma n$ such that $e(S) \geq \theta|S|$.

Proof

$$\mathbb{P}(\exists S : |S| \leq \sigma n \text{ and } e(S) \geq \theta|S|) \leq \sum_{s=2\theta}^{\sigma n} \binom{n}{s} \binom{\binom{s}{2}}{\theta s} \left(\frac{d}{n} \right)^{\theta s} \tag{2.6}$$

$$\begin{aligned}
& \leq \sum_{s=2\theta}^{\sigma n} \left(\frac{ne}{s} \left(\frac{eds}{2\theta n} \right)^\theta \right)^s \\
& = \sum_{s=2\theta}^{\sigma n} \left(e \left(\frac{s}{n} \right)^{\theta-1} \left(\frac{ed}{2\theta} \right)^\theta \right)^s \\
& = O \left(\frac{d^\theta}{n^{\theta-1}} \right) = o(1)
\end{aligned} \tag{2.7}$$

provided $d = o(n^{1-1/\theta})$. □

We will apply this lemma with $\theta \geq 2 - \varepsilon$ for $\varepsilon \ll 1$ and this fits with our bound on d .

Lemma 2.2. *Let σ, θ be as in Lemma 2.1. If $(\Delta - 2\theta)\tau > 1$ and*

$$\left(\frac{\sigma ed}{(\Delta - 2\theta)\tau} \right)^{(\Delta - 2\theta)\tau} \leq \frac{\sigma}{4e}$$

then w.h.p. there do not exist $S \supseteq T$ such that $|S| = s \leq \sigma n$, $|T| \geq \tau s$ and $d_S(v) \geq \Delta$ for every $v \in T$.

Proof In the light of Lemma 2.1, the assumptions imply that w.h.p. $|e(T : S \setminus T)| \geq (\Delta - 2\theta)\tau s$. In which case,

$$\begin{aligned} & \mathbb{P}(\exists S \supseteq T, |S| \leq \sigma n, |T| \geq \tau s : |e(T : S \setminus T)| \geq (\Delta - 2\theta)\tau s) \\ & \leq \sum_{s=2\theta}^{\sigma n} \sum_{t=\tau s}^s \binom{n}{s} \binom{s}{t} \left(\frac{edt}{(\Delta - 2\theta)\tau n} \right)^{(\Delta - 2\theta)\tau s} \end{aligned} \quad (2.8)$$

$$\begin{aligned} & \leq \sum_{s=2\theta}^{\sigma n} \sum_{t=\tau s}^s \left(\frac{ne}{s} \right)^s \cdot 2^s \cdot \left(\frac{eds}{(\Delta - 2\theta)\tau n} \right)^{(\Delta - 2\theta)\tau s} \\ & = \sum_{s=2\theta}^{\sigma n} \sum_{t=\tau s}^s \left(\frac{2ne}{s} \cdot \left(\frac{eds}{(\Delta - 2\theta)\tau n} \right)^{(\Delta - 2\theta)\tau} \right)^s \\ & = \sum_{s=2\theta}^{\sigma n} \sum_{t=\tau s}^s \left(2e \left(\frac{s}{n} \right)^{(\Delta - 2\theta)\tau - 1} \cdot \left(\frac{ed}{(\Delta - 2\theta)\tau} \right)^{(\Delta - 2\theta)\tau} \right)^s \\ & = O\left(\frac{d^{(\Delta - 2\theta)\tau}}{n^{(\Delta - 2\theta)\tau - 1}} \right) = o(1). \end{aligned} \quad (2.9)$$

□

We will apply this lemma with $(\Delta - 2\theta)\tau \geq 2$ and this fits with our bound on d .

Fix $\alpha > 12$ and let

$$k = \frac{\alpha d}{\ln d} \text{ and } \beta = \frac{\alpha d^{1-1/\alpha}}{\ln d} \text{ and } \gamma = \frac{16 \ln^2 d}{\alpha d^{1-1/\alpha}}.$$

We will now argue that w.h.p. A can win the game if k colors are available.

A's initial strategy will be the same as that described in [6]. Let $\mathcal{C} = (C_1, C_2, \dots, C_k)$ be a collection of pairwise disjoint subsets of $[n]$, i.e. a (partial) coloring. Let $\bigcup \mathcal{C}$ denote $\bigcup_{i=1}^k C_i$. For a vertex v let

$$A(v, \mathcal{C}) = \{i \in [k] : v \text{ is not adjacent to any vertex of } C_i\},$$

and set

$$a(v, \mathcal{C}) = |A(v, \mathcal{C})|.$$

Note that $A(v, \mathcal{C})$ is the set of colors that are available at vertex v when the partial coloring is given by the sets in \mathcal{C} and $v \notin \bigcup \mathcal{C}$. A's initial strategy can now be easily defined. Given the current color classes \mathcal{C} , A chooses an uncolored vertex v with the smallest value of $a(v, \mathcal{C})$ and colors it by any available color.

As the game evolves, we let u denote the number of uncolored vertices in the graph. So, we think of u as running "backward" from n to 0.

We show next that w.h.p. every k -coloring (proper or improper) of the full vertex set has the property that there are at most γn vertices with less than $\beta/2$ available colors. Let

$$B(\mathcal{C}) = \{v : a(v, \mathcal{C}) < 3\beta/4\}.$$

Lemma 2.3. *W.h.p., for all collections \mathcal{C} ,*

$$|B(\mathcal{C})| \leq \gamma n.$$

Proof We first note that if $|S| = \gamma n$ then w.h.p. S contains at most $4\gamma^2 dn$ edges. This follows from Lemma 2.1 with $\sigma = \gamma$ and $\theta = 4\gamma d$. It follows that for any $\varepsilon > 0$ that there is a set $S_1 \subseteq S$ of size at least $(1 - \varepsilon)\gamma n$ such that if $v \in S_1$ then its degree $d_S(v)$ in S is at most $8\varepsilon^{-1}\gamma d$.

Fix \mathcal{C} and suppose that $v \in S_1$. Let

$$b(v, \mathcal{C}) = |\{i \in [k] : v \text{ is not adjacent to any vertex of } C_i \setminus S\}|.$$

Thus $a(v, \mathcal{C}) \geq b(v, \mathcal{C}) - 8\varepsilon^{-1}\gamma d$. $b(v, \mathcal{C})$ is the sum of independent indicator variables X_i , where $X_i = 1$ if v has no neighbors in $C_i \setminus S$ in $G_{n,p}$. Then $\mathbb{P}(X_i = 1) \geq (1 - p)^{|C_i|}$ and since $(1 - p)^t$ is a convex function of t we have

$$\begin{aligned} \mathbb{E}(b(v, \mathcal{C})) &\geq \sum_{i=1}^k (1 - p)^{|C_i|} \\ &\geq k(1 - p)^{(|C_1| + \dots + |C_k|)/k} \\ &\geq k(1 - p)^{n/k} \\ &= \beta - o(\beta). \end{aligned}$$

It follows from the Chernoff bound (1.1) that

$$\mathbb{P}(b(v, \mathcal{C}) \leq 0.51\beta) \leq e^{-\beta/33}.$$

Now, when \mathcal{C} is fixed, the events $\{b(v, \mathcal{C}) \leq 0.51\beta\}, v \in S_1$ are independent. Thus, because $a(v, \mathcal{C}) \leq \beta/2$ implies that $b(v, \mathcal{C}) \leq 0.51\beta$ we have

$$\begin{aligned} &\mathbb{P}(\exists \mathcal{C} : |B(\mathcal{C})| \geq \gamma n) \\ &\leq k^n \binom{n}{(1 - \varepsilon)\gamma n} e^{-(1 - \varepsilon)\gamma\beta n/33} \\ &\leq d^n \left(\frac{e}{(1 - \varepsilon)\gamma} \exp \left\{ -\frac{\alpha d^{1-1/\alpha}}{33 \ln d} \right\} \right)^{(1 - \varepsilon)\gamma n} \tag{2.10} \\ &= \exp \left\{ n \left(\ln d + (1 - \varepsilon)\gamma \left(\ln \left(\frac{1}{1 - \varepsilon} \right) + \ln \left(\frac{\alpha}{16} \right) + (1 - 1/\alpha) \ln d - 2 \ln \ln d - \frac{\alpha d^{1-1/\alpha}}{33 \ln d} \right) \right) \right\} \\ &= o(1), \end{aligned}$$

for large d and small enough ε . □

Let u_0 to be the last time for which A colors a vertex with at least $\beta/2$ available colors, i.e.,

$$u_0 = \min \left\{ u : a(v, \mathcal{C}_u) \geq 3\beta/4, \text{ for all } v \notin \bigcup \mathcal{C}_u \right\},$$

where \mathcal{C}_u denotes the collection of color classes when u vertices remain uncolored.

If u_0 does not exist then A will win.

It follows from Lemma 2.3 that w.h.p. $u_0 \leq 2\gamma n$ and that at time u_0 , every vertex still has at least $\beta/2$ available colors. Indeed, consider the final coloring \mathcal{C}^* in the game that would be achieved if A follows her current strategy, even if she has to improperly color an edge. Let $U = \{v \notin \mathcal{C}_{u_0} : a(v, \mathcal{C}^*) < \beta/2\}$. Now we can assume that $|U| \leq \gamma n$. Because the number of colors available to a vertex decreases as vertices get colored, from u_0 onwards, every vertex colored by A is in U . Therefore $u_0 \leq 2\gamma n$.

Now let u_1 be the first time that there are at most $2\gamma n$ uncolored vertices and $a(v, \mathcal{C}_u) \geq \beta/2$, for all $v \notin \bigcup \mathcal{C}_u$. By the above, w.h.p. $u_1 \leq u_0$, so in particular w.h.p. u_1 exists. A can determine u_1 but not u_0 , as u_0 depends on the future.

A will follow a more sophisticated strategy from u_1 onwards. We will show next that we can find a sequence $U = U_0 \supseteq U_1 \supseteq \dots \supseteq U_\ell$ with the following properties: The edges of $U_i : (U_{i-1} \setminus U_i)$ between U_i and $U_{i-1} \setminus U_i$ will be divided into two classes, *heavy* and *light*. Vertex w is a heavy (resp. light) neighbor of vertex v if the edge (v, w) is heavy (resp. light).

(P1) Each vertex of $U_i \setminus U_{i+1}$ has at most one light neighbor in U_{i+1} , for $0 \leq i < \ell$.

(P2) All $U_i : (U_{i-1} \setminus U_i)$ edges are light for $i \geq 2$.

(P3) Each vertex of U_1 either has degree at most $\beta/3$ in U_0 or it has at most $\beta/20$ heavy neighbors in $U_0 \setminus U_1$.

(P4) $d_{U_i}(v) \leq \beta/3$ for $v \in U_i \setminus U_{i+1}$.

(P5) U_ℓ contains at most one cycle.

From this, we can deduce that the edges of U_0 can be divided up into the heavy edges E_H , light edges E_L , the edges inside U_ℓ and the rest of the edges. Assume first that U_ℓ does not contain a cycle. $F = (U, E_L)$ is a forest and the strategy in [10] can be applied. When attempting to color a vertex v of F , there are never more than three F -neighbors of v that have been colored. Since there are at most $\beta/3 + \beta/20$ non- F neighbors, A will succeed since she has an initial list of size $\beta/2$.

If U_ℓ contains a cycle C then A can begin by coloring a vertex of C . This puts A one move behind in the tree coloring strategy, in which case we can bound the number of F -neighbors by four.

It only remains to prove that the construction P1–P5 exists w.h.p. Remember that d is sufficiently large here.

We can assume without loss of generality that $|U_0| = 2\gamma n$. This will not decrease the sizes of the sets $a(v, U_0)$.

2.2.1 The verification of P1–P4: Constructing U_1

Applying Lemma 2.2 with

$$\sigma = 2\gamma \text{ and } \theta = d^{1/\alpha} \ln^3 d \text{ and } \Delta = 2\theta + \beta/4 < \beta/3 \text{ and } \tau = \theta/\beta$$

we see that w.h.p.

$$U'_{1,a} = \{v \in U_0 : d_{U_0}(v) \geq 2\theta + \beta/4\} \text{ satisfies } |U'_{1,a}| \leq 2\tau\gamma n = \frac{64d^{3/\alpha} \ln^6 d}{\alpha^2 d^2} n.$$

We then let $U_{1,a} \supseteq U'_{1,a}$ be the subset of U_0 consisting of the vertices with the $2\tau\gamma n$ largest values of d_{U_0} .

We then construct $U_{1,b} \supseteq U_{1,a}$ by repeatedly adding vertices x_1, x_2, \dots, x_r of $U \setminus U_{1,a}$ such that x_j is the lowest numbered vertex not in $X_j = U_{1,a} \cup \{x_1, x_2, \dots, x_{j-1}\}$ having at least three neighbors in X_j . This ends with $r \leq 5|U_{1,a}|$ in order that we do not violate the conclusion of Lemma 2.1 with

$$\sigma = 12\tau\gamma = \frac{192d^{3/\alpha} \ln^6 d}{\alpha^2 d^2} \text{ and } \theta = 5/2$$

which is applicable since

$$\left(\frac{384ed^{3/\alpha} \ln^6 d}{5\alpha^2 d} \right)^{5/2} < \frac{192d^{3/\alpha} \ln^6 d}{2e\alpha^2 d^2}.$$

It follows that

$$|U_{1,b}| \leq 12\tau\gamma n. \tag{2.11}$$

Next let A be the set of vertices in $U \setminus U_{1,b}$ that have two neighbors in $U_{1,b}$ and let B be the set of vertices in $U_{1,b}$ that have more than $\beta/20$ neighbors in A . We argue that w.h.p.

$$|B| \leq \frac{1200d^{3/\alpha}\tau\gamma n}{\beta} = \frac{48000 \ln^4 d}{\alpha^3 d^{3-6/\alpha}} n. \tag{2.12}$$

We first consider the size of A . We first prove that w.h.p.

$$|A| \leq 12d^{3/\alpha}\tau\gamma n. \tag{2.13}$$

For a set S , let $D_2(S)$ denote the set of vertices not in S that have at least two neighbors in S .

Lemma 2.4. *W.h.p. $|D_2(S)| \leq d^{3/\alpha}|S|$ for all $|S| \leq 12\tau\gamma n$.*

Proof For easy reference we note that $\tau\gamma = \frac{40\ln^3 d}{\alpha^2 d^{2-2/\alpha}}$. For $|S| \leq 12\tau\gamma n$ and $K = d^{3/\alpha}$,

we have

$$\begin{aligned}
\mathbb{P}(\exists |S| \leq 12\tau\gamma : |D_2(S)| \geq K|S|) &\leq \sum_{s=2}^{12\tau\gamma n} \binom{n}{s} \binom{n}{Ks} \left(\binom{s}{2} \frac{d^2}{n^2} \right)^{Ks} \\
&\leq \sum_{s=2}^{12\tau\gamma n} \left(\frac{ne}{s} \right)^s \left(\frac{ne}{Ks} \right)^{Ks} \left(\frac{s^2 d^2}{2n^2} \right)^{Ks} \\
&= \sum_{s=2}^{12\tau\gamma n} \left(\left(\frac{sed^2}{2Kn} \right)^{K-1} \cdot \frac{e^2 d^2}{2K} \right)^s \\
&= o(1).
\end{aligned}$$

□

Equation (2.13) follows immediately from (2.11) and Lemma 2.4. To bound the size of B we prove the following lemma.

Lemma 2.5. *W.h.p. there do not exist disjoint sets S, T such that $|T| \leq t_0 = 12d^{3/\alpha}\tau\gamma n$ and $|S| \geq 100|T|/\beta$ such that each $v \in S$ has at least $\beta/20$ neighbors in T .*

Proof We have

$$\begin{aligned}
\mathbb{P}(\exists S, T \text{ denying lemma}) &\leq \sum_{t=\beta/20}^{t_0} \binom{n}{t} \binom{n}{100t/\beta} \left(\binom{t}{\beta/20} \left(\frac{d}{n} \right)^{\beta/20} \right)^{100t/\beta} \\
&\leq \sum_{t=\beta/20}^{t_0} \left(\frac{ne}{t} \right)^t \left(\frac{ne\beta}{100t} \right)^{100t/\beta} \left(\frac{20etd}{\beta n} \right)^{5t} \\
&= \sum_{t=\beta/20}^{t_0} \left(\left(\frac{t}{n} \right)^{4-100/\beta} \frac{e^{100/\beta+6}\beta^{100/\beta-5}d^5}{100^{100/\beta}} \right)^t \\
&= o(1).
\end{aligned}$$

□

Equation (2.12) follows immediately from (2.13) and Lemma 2.5.

Now let A_1 be the set of vertices in A that have two neighbors of B . It follows from Lemma 2.4 that w.h.p.

$$|A_1| \leq d^{3/\alpha}|B| \leq \frac{48000 \ln^4 d}{\alpha^3 d^{3-9/\alpha}} n.$$

Next let B_1 be the set of vertices in B that have at least $\beta/20$ neighbors of A_1 . It follows from Lemma 2.5 that w.h.p.

$$|B_1| \leq \frac{100|A_1|}{\beta} \leq \frac{4800000 \ln^5 d}{\alpha^4 d^{4-10/\alpha}} n.$$

Lemma 2.6. *W.h.p. if $s_0 = e^{-d}n \leq |S| \leq s_1 = \frac{4800000 \ln^5 d}{\alpha^4 d^{4-10/\alpha}} n$ then $|N(S)| \leq d^{1+1/\alpha}|S|$.*

Proof

$$\begin{aligned}
\mathbb{P}(\exists S : \text{denying lemma}) &\leq \sum_{s=s_0}^{s_1} \binom{n}{s} \binom{n}{d^{1+1/\alpha}s} \left(\frac{sd}{n}\right)^{d^{1+1/\alpha}s} \\
&= \sum_{s=s_0}^{s_1} \left(\frac{ne}{s} \left(\frac{e}{d^{1/\alpha}}\right)^{d^{1+1/\alpha}}\right)^s \\
&= o(1).
\end{aligned}$$

□

Now let A_2 be the set of vertices in A_1 that have a neighbor in B_1 . It follows from Lemma 2.6 that w.h.p.

$$|A_2| \leq \frac{4800000 \ln^5 d}{\alpha^4 d^{3-11/\alpha}} n.$$

(Note that if $|S| < e^{-d}n$ then Lemma 2.6 implies that $|N(S)| \leq |S| + d^{1+1/\alpha}e^{-d}n$.)

Next let $U_{1,c} = U_{1,b}$ and $Y_0 = A_2 \cup B_1$. We now construct $U_1 \supseteq U_{1,c}$ by repeatedly adding vertices y_1, y_2, \dots, y_s of $U \setminus U_{1,c}$ such that y_j is the lowest numbered vertex not in $Y_j = Y_0 \cup \{y_1, y_2, \dots, y_{j-1}\}$ that has at least two neighbors in Y_j . Taking $\theta = 3/2$ and using Lemma 2.1, this ends with $s \leq 3|B|$ by the same argument used to show $s \leq 5|U_{1,a}|$ above. Note that

$$|U_1| \leq \gamma_1 n = 13\tau\gamma n = \frac{208 \ln^6 d}{\alpha^2 d^{2-3/\alpha}}.$$

We let $W = U_0 \setminus U_1$ and partition the edges $W : U_1$ into light and heavy edges.

- (A) $W : (Y_s \setminus U_{1,b})$: These edges will be heavy.
- (B) $(W \setminus (A \cup Y_s)) : U_{1,b}$. These edges will be light. Note that each $v \in W \setminus (A \cup Y_s)$ has at most one neighbor in $U_{1,b}$.
- (C) $(A \setminus (A_1 \cup Y_s)) : (U_{1,b} \setminus Y_s)$. If $v \in A \setminus (A_1 \cup Y_s)$ then v has at most two neighbors in $U_{1,b}$. At most one of these can be in B and we make the corresponding edge light. If v has a neighbor in $U_{1,b} \setminus B$ then we make the corresponding edge heavy.
- (D) $(A_1 \setminus Y_s) : (U_{1,b} \setminus Y_s)$. If $v \in A_1 \setminus Y_s$ then v has at most two neighbors in B . None of these can be in B_1 and we make the corresponding edges heavy.

We now have to check that **P1–P4** hold.

First consider the light edges. There is at most one for each $v \in W$ and so **P1** holds.

Now consider the heavy edges. Vertices in B can only have light neighbors in $W \setminus A$ and vertices in $B \setminus B_1$ have at most $\beta/20$ heavy neighbors in A . Vertices in $U_{1,b} \setminus B$ can have at most $\beta/20$ heavy neighbors in W . Vertices in $U_1 \setminus U_{1,b}$ have (heavy) degree at most $\beta/3$ in U_0 . This verifies **P3** and **P4** holds by the definition of $U_{1,a}$.

2.2.2 The verification of P1–P4: Constructing U_2

Applying Lemma 2.2 again, with

$$\sigma = \gamma_1 \text{ and } \theta = 3 \text{ and } \Delta = 2\theta + \beta/3 \text{ and } \tau = 12/\beta$$

we see that w.h.p.

$$U'_2 = \{v \in U_1 : d_{U_1}(v) \geq 6 + \beta/3\} \text{ satisfies } |U'_2| \leq \gamma'_2 = \frac{12\gamma_1}{\beta} \leq \frac{2500 \ln^7 d}{\alpha^3 d^{3-4/\alpha}}. \quad (2.14)$$

We then construct $U_2 \supseteq U'_2$ by repeatedly adding vertices x_1, x_2, \dots, x_r of $U_1 \setminus U'_2$ such that x_i has at least two neighbors in $U'_2 \cup \{x_1, x_2, \dots, x_{i-1}\}$. This ends with $r \leq 7|U'_2|$ in order that we do not violate the conclusion of Lemma 2.1 with

$$\sigma = 8\gamma'_2 \leq \frac{20000 \ln^7 d}{\alpha^2 d^{3-4/\alpha}} \text{ and } \theta = \frac{15}{8} \quad (2.15)$$

which is applicable since

$$\left(\frac{20000 \cdot 4 \cdot e \ln^4 d}{15\alpha^3 d^{2-4/\alpha}} \right)^{15/8} < \frac{20000 \ln^7 d}{2e\alpha^3 d^{3-4/\alpha}}.$$

This verifies P1–P4 with $i = 1$.

2.2.3 The verification of P1–P5: Constructing U_i , $i \geq 3$

We now repeat the argument to create the sequence $U_0 \supseteq U_1 \supseteq \dots \supseteq U_\ell$. The value of θ has decreased to $15/8$ (see (2.15)) and $|U_i| \leq (12\gamma/\beta)|U_{i-1}|$, as in (2.14). We choose ℓ so that $|U_\ell| \leq \ln n$. We can easily prove that w.h.p. S contains at most $|S|$ edges whenever $|S| \leq \ln n$, implying P5.

This completes the proof of Part (b) of Theorem 1.2.

3 Theorem 1.3: $G_{n,d}$

We will not change A or B's strategies. We will simply transfer the relevant structural results from $G_{n,d/n}$ to $G_{n,d}$. Some of the unimportant constants will change, but this will not change the verification of the success of the various strategies. We will first do this using Theorem 1.2 under the assumption that $d \leq n^{1/4}$. For larger d we will use Theorem 1.1 and the “sandwiching theorem” of Kim and Vu [15]. This latter analysis is given in Section 3.2.

3.1 $d_0 \leq d \leq n^{1/4}$

Here we assume that d_0 is a sufficiently large constant. We begin with the configuration model of Bollobás [8]. We have a set W of *points* and this is partitioned into sets W_1, W_2, \dots, W_n of size d . We define $\phi : W \rightarrow [n]$ by $\phi(x) = j$ for all $x \in W_j$. We associate each *pairing* or *configuration* F of W into $|W|/2$ pairs to a multigraph G_F on the vertex set $[n]$. A pair $\{x, y\} \in F$ becomes an edge $(\phi(x), \phi(y))$ of G_F . Now there are $\frac{(dn)!}{(dn/2)!2^{dn/2}}$ pairings and each simple d -regular graph (without loops or multiple edges) arises $(d!)^n$ times as G_F . So for any pair of d -regular graphs G_1, G_2 we have

$$\mathbb{P}(G_F = G_1 \mid G_F \text{ is simple}) = \mathbb{P}(G_F = G_2 \mid G_F \text{ is simple}). \quad (3.1)$$

In order to use this, we need a bound on the probability that G_F is simple.

$$\mathbb{P}(G_F \text{ is simple}) \geq e^{-2d^2}. \quad (3.2)$$

This is the content of Lemma 2 of [9].

It follows from (3.1) and (3.2) that for any graph property \mathcal{A} :

$$\varepsilon^{2d^2} \mathbb{P}(G_F \in \mathcal{A}) = o(1) \text{ implies } \mathbb{P}(G_{n,d} \in \mathcal{A}) = o(1). \quad (3.3)$$

We can use the above to estimate $\rho = \mathbb{P}(G_{n,d/n} \text{ is } d \text{ regular})$. We write this as

$$\rho = \mathbb{P}(G = G_{n,d/n} \text{ is } d \text{ regular} \mid |E(G)| = dn/2) \mathbb{P}(|E(G)| = m = dn/2).$$

It is easy to show, using Stirling's approximation, that

$$\mathbb{P}(|E(G)| = m) = \Omega(m^{-1/2})$$

and so we concentrate on the other factor.

Let $N = \binom{n}{2}$. There are $\binom{N}{m} \leq \left(\frac{Ne}{m}\right)^m$, graphs with vertex set $[n]$ and m edges of which

$$\Omega\left(\frac{e^{-2d^2} (dn)!}{(dn/2)!2^{dn/2}(d!)^n}\right) \text{ are } d\text{-regular.}$$

So, since $d = o(n)$,

$$\begin{aligned} \rho &= \Omega\left(\frac{e^{-2d^2}}{(dn)^{1/2}} \cdot \left(\frac{dn}{e}\right)^{dn/2} \cdot \frac{1}{(d!)^n} \cdot \left(\frac{d}{e(n-1)}\right)^{dn/2}\right) = \\ &= \Omega\left(\frac{d^{dn}}{(dn)^{1/2} e^{dn+2d^2} (d!)^n}\right) = \Omega\left(\left(\frac{1}{10d}\right)^{n/2}\right). \end{aligned} \quad (3.4)$$

We need another crude estimate. We prove a small modification of Lemma 1 from [9].

Lemma 3.1. *Given $\{a_i, b_i\}$, $i = 1, 2, \dots, k \leq n/8d$ then*

$$\mathbb{P}(\{(a_i, b_i) \in E(G_{n,d}), 1 \leq i \leq k\}) \leq \left(\frac{20d}{n}\right)^k.$$

Proof Let \mathcal{G}_d denote the set of d -regular graphs with vertex set $[n]$. For $0 \leq t \leq k$ we let

$$\Omega_t = \{G \in \mathcal{G}_d : \{a_i, b_i\} \in E(G), 1 \leq i \leq t \text{ and } \{a_i, b_i\} \notin E(G), t+1 \leq i \leq k\}.$$

We consider the set X of pairs $(G_1, G_2) \in \Omega_t \times \Omega_{t-1}$ such that G_2 is obtained from G_1 by deleting disjoint edges $\{a_t, b_t\}, \{x_1, y_1\}, \{x_2, y_2\}$ and replacing them by $\{a_t, x_1\}, \{y_1, y_2\}, \{b_t, x_2\}$. Given G_1 , we can choose $\{x_1, y_1\}, \{x_2, y_2\}$ to be any ordered pair of disjoint edges which are not incident with $\{a_1, b_1\}, \dots, \{a_k, b_k\}$ or their neighbours and such that $\{y_1, y_2\}$ is not an edge of G_1 . Thus each $G_1 \in \Omega_1$ is in at least $(D - (2kd^2 + 1))(D - (2kd^2 + 2))$ pairs, where $D = dn/2$. Each $G_2 \in \Omega_{t-1}$ is in at most $2Dd^2$ pairs. The factor of 2 arises because a suitable edge $\{y_1, y_2\}$ of G_2 has an orientation relative to the switching back to G_1 . It follows that

$$\frac{|\Omega_t|}{|\Omega_{t-1}|} \leq \frac{2Dd^2}{(D - (2kd^2 + 1))(D - (2kd^2 + 2))} \leq \frac{20d}{n}.$$

It follows that

$$\frac{|\Omega_k|}{|\Omega_0| + \dots + |\Omega_k|} \leq \left(\frac{20d}{n}\right)^k$$

and this implies the lemma. □

3.1.1 The lower bound

Using (3.2) we can replace (2.3) by

$$e^{2d^2} \exp\{(\beta + \gamma + \beta^2 - (2\beta + \gamma)^2/2 + o_d(1))d^{-1}n \ln^2 d\} = o(1)$$

for $d \leq n^{1/4}$. After this, we can argue as in the case $G_{n,p}$.

3.1.2 The upper bound

We first need to prove the equivalent of Lemmas 2.1 and 2.2.

Lemma 3.2. *If $1 < \theta \leq d^{1/6} \ln^3 d$ and*

$$\left(\frac{10\sigma ed}{\theta}\right)^\theta \leq \frac{\sigma}{2e} \tag{3.5}$$

then w.h.p. there does not exist $S \subseteq [n], |S| \leq \sigma n$ such that $e(S) \geq \theta|S|$.

Proof

$$\mathbb{P}(\exists S : |S| \leq \sigma n \text{ and } e(S) \geq \theta |S|) \leq \sum_{s=2\theta}^{\sigma n} \binom{n}{s} \binom{\binom{s}{2}}{\theta s} \pi_s$$

where

$$\pi_s = \max_{\substack{X \subseteq \binom{[s]}{2} \\ |X| = \theta s}} \mathbb{P}(E(G_{n,d}) \supseteq X).$$

It follows from (3.2) that $\pi_s \leq e^{2d^2} \left(\frac{d}{n}\right)^{\theta s}$. If d is small, say $d \leq \ln^{1/3} n$ then we can see from the proof of Lemma 2.1 that

$$\mathbb{P}(\exists S : |S| \leq \sigma n \text{ and } e(S) \geq \theta |S|) \leq O\left(e^{2\ln^{2/3} n} \cdot \frac{d^\theta}{n^{\theta-1}}\right) = o(1).$$

We can therefore assume that $d \geq \ln^{1/3} n$ and then

$$\begin{aligned} \sum_{s=3d^2}^{\sigma n} \binom{n}{s} \binom{\binom{s}{2}}{\theta s} \pi_s &\leq e^{2d^2} \sum_{s=3d^2}^{\sigma n} \binom{n}{s} \binom{\binom{s}{2}}{\theta s} \left(\frac{d}{n}\right)^{\theta s} \\ &\leq e^{2d^2} \sum_{s=3d^2}^{\sigma n} \left(e \left(\frac{s}{n}\right)^{\theta-1} \left(\frac{ed}{2\theta}\right)^\theta \right)^s \\ &\leq e^{2d^2} \sum_{s=3d^2}^{\sigma n} 2^{-s} \\ &= o(1). \end{aligned}$$

When $s \leq 3d^2$ we use Lemma 3.1. For this we will need to have $\theta s \leq 3\theta d^2 \leq n/8d$. The maximum value of θ is $d^{1/6} \ln^3 d$ and so the lemma can indeed be applied for $d \leq n^{1/4}$. Assuming this, we have

$$\begin{aligned} \sum_{2\theta}^{3d^2} \binom{n}{s} \binom{\binom{s}{2}}{\theta s} \pi_s &\leq \sum_{2\theta}^{3d^2} \binom{n}{s} \binom{\binom{s}{2}}{\theta s} \left(\frac{20d}{n}\right)^{\theta s} \\ &\leq \sum_{2\theta}^{3d^2} \left(e \left(\frac{s}{n}\right)^{\theta-1} \left(\frac{20ed}{2\theta}\right)^\theta \right)^s \\ &= o(1). \end{aligned}$$

□

Lemma 3.3. *Let σ, θ be as in Lemma 3.2. If*

$$\left(\frac{10\sigma ed}{(\Delta - 2\theta)\tau} \right)^{(\Delta - 2\theta)\tau} \leq \frac{\sigma}{4e}$$

then w.h.p. there do not exist $S \supseteq T$ such that $|S| \leq \sigma n$, $|T| \geq \tau s$ and $d_S(v) \geq \Delta$ for $v \in T$.

Proof We first argue that if $d \leq \ln^{1/3} n$ then we prove the lemma by just inflating the failure probability by e^{2d^2} as we did for Lemma 3.2.

We therefore assume that $d \geq \ln^{1/3} n$ and write

$$\begin{aligned} \mathbb{P}(\exists S \supseteq T, |S| \leq \sigma n, |T| \geq \tau s, |e(T : S \setminus T)| \geq (\Delta - 2\theta)\tau s) \\ \leq \sum_{s,t} \binom{n}{s} \binom{s}{t} \binom{t(s-t)}{(\Delta - 2\theta)\tau s} \pi_s \end{aligned}$$

where now we have

$$\pi_s = \max_{\substack{X \subseteq T \times (S \setminus T) \\ |X| = (\Delta - 2\theta)\tau s}} \mathbb{P}(E(G_{n,d}) \supseteq X).$$

Using (3.2) we write

$$\begin{aligned} & \sum_{s=3d^2/\tau}^{\sigma n} \sum_{t=\tau s}^s \binom{n}{s} \binom{s}{t} \binom{t(s-t)}{(\Delta - 2\theta)\tau s} \pi_s \\ & \leq e^{2d^2} \sum_{s=3d^2/\tau}^{\sigma n} \sum_{t=\tau s}^s \binom{n}{s} \binom{s}{t} \left(\frac{edt}{(\Delta - 2\theta)\tau n} \right)^{(\Delta - 2\theta)\tau s} \\ & \leq e^{2d^2} \sum_{s=3d^2/\tau}^{\sigma n} \sum_{t=\tau s}^s \left(2e \left(\frac{s}{n} \right)^{(\Delta - 2\theta)\tau - 1} \cdot \left(\frac{ed}{(\Delta - 2\theta)\tau} \right)^{(\Delta - 2\theta)\tau} \right)^s \\ & \leq e^{2d^2} \sum_{s=3d^2/\tau}^{\sigma n} \sum_{t=\tau s}^s 2^{-s} \\ & = o(1). \end{aligned}$$

When $s \leq 3d^2/\tau$ use Lemma 3.1, with the same caveats on the value of d . So,

$$\begin{aligned} & \sum_{s=2\theta}^{3d^2/\tau} \sum_{t=\tau s}^s \binom{n}{s} \binom{s}{t} \binom{t(s-t)}{(\Delta - 2\theta)\tau s} \pi_s \\ & \leq \sum_{s=2\theta}^{3d^2/\tau} \sum_{t=\tau s}^s \binom{n}{s} \binom{s}{t} \binom{t(s-t)}{(\Delta - 2\theta)\tau s} \left(\frac{20d}{n} \right)^{(\Delta - 2\theta)\tau s} \\ & = O \left(\frac{d^{(\Delta - 2\theta)\tau}}{n^{(\Delta - 2\theta)\tau - 1}} \right) = o(1). \end{aligned}$$

□

Remark 1. We can estimate $\mathbb{P}(\exists \mathcal{C} : |B(\mathcal{C})| \geq \gamma n)$ by multiplying (2.10) by $1/\rho$ and notice that it remains $o(1)$.

After this, the proof will much the same as for $G_{n,p}$, but with a few constants being changed.

3.2 $n^{1/4} \leq d \leq n^{1/3-\varepsilon}$

Our approach in this section is to use the sandwiching technique developed by Kim and Vu in [15] to adapt the proof of Theorem 1.1. In some sense it is pretty clear that given the results of [15], it will be possible to translate the results of [6] to deal with large regular graphs. We will carry out the task, but our proof will be abbreviated and rely on notation from the latter paper.

Without changing the strategy used in obtaining the lower bound, we show that each intermediate result used to prove the theorem in [6] continues to hold for random regular graphs $G_{n,d}$ in the range where these can be approximated sufficiently well by random graphs $G_{n,d/n}$. In order to get the required strength from the Kim-Vu coupling, however, we require $d = np = n^\varepsilon$ for some $\varepsilon \geq \varepsilon_0$, where ε_0 is a small absolute constant.

3.2.1 Notation

We use Theorem 2 in [15] to get a joint distribution on (H_1, G, H_2) : G is d -regular, $H_1 \subseteq G$, $H_1 \subseteq H_2$, and although $G \not\subseteq H_2$, this is almost true in a way we discuss further. The graphs H_1 and H_2 are random graphs with edge probabilities p_1 and p_2 , and by judicious choice of parameters we can set $p_1 = p/(1 + \delta)$ and $p_2 = p(1 + \delta)$, where

$$p = \frac{d}{n} \text{ and } \delta = \Theta \left(\left(\frac{\ln n}{d} \right)^{1/3} \right).$$

Constants defined in [6] are in terms of p and we will make this relationship explicit. Of note is the constant

$$\ell_1(p) = \log_b n - \log_b \log_b np - 10 \log_b \ln n$$

where $b = b(p) = \frac{1}{1-p}$.

3.2.2 Kim-Vu coupling

The construction of the coupling (H_1, G, H_2) in [15] yields $H_1 \subseteq G$ w.h.p., but not $G \subseteq H_2$. As a substitute for such a result, we prove the following lemma.

Lemma 3.4. $\Delta(G \setminus H_2) = O(1)$ w.h.p.

Proof We rely on the bound $\Delta(G \setminus H_2) \leq \Delta(G) - \delta(H_2) + \Delta(H_2 \setminus G)$. Trivially, $\Delta(G) = d$. Part 3 of Theorem 2 in [15] states that w.h.p.

$$\Delta(H_2 \setminus G) \leq \frac{(1 + o(1)) \ln n}{\ln(\delta d / \ln n)} = \frac{(1 + o(1)) \ln n}{\frac{2}{3} \ln d - \frac{2}{3} \ln \ln n + O(1)} = \frac{3 + o(1)}{2\varepsilon}.$$

We prove that w.h.p. $\delta(H_2) \geq d$. For any vertex v , $\deg_{H_2} v$ follows the binomial distribution $B(n-1, p_2)$. By the Chernoff bound,

$$\mathbb{P}[\deg_{H_2} v < d] \leq \mathbb{P}\left[B(n-1, p_2) < \left(1 - \frac{\delta}{2(1+\delta)}\right)(n-1)p_2\right] \leq e^{-\delta^2/(10(1+\delta))}.$$

We can simplify the exponent here to

$$-\frac{\delta^2 d}{10(1+\delta)} = -\frac{\Omega((\ln n)^{2/3} d^{1/3})}{1+\delta} \leq -\Omega(n^{\varepsilon/3}).$$

So $\mathbb{P}[\deg_{H_2} v < d] \leq O(n^{-\Omega(n^{\varepsilon/3})})$ and $\mathbb{P}[\delta(H_2) < d] = o(1)$, completing the proof. \square

3.2.3 Bounds

We first prove a few auxiliary bounds on the relationship between p , p_1 , and p_2 , as well as other constants in terms of these probabilities.

Bound 3.1.

$$1 \geq \frac{\ell_1(p)}{\ell_1(p_1)} \geq 1 - 2\delta, \text{ and } 1 \leq \frac{\ell_1(p_1)}{\ell_1(p)} \leq 1 + 2\delta.$$

Proof We first note that if $np \rightarrow \infty$ then the derivative $(\log_{b(p)} np)' < 0$ and so if we let $x = \frac{n}{\log_{np} np \ln^{10} n}$ then,

$$\begin{aligned} \frac{\ell_1(p_1)}{\ell_1(p)} &= \frac{\log_{b(p_1)} n - \log_{b(p_1)} \log_{b(p_1)} np_1 - 10 \log_{b(p_1)} \ln n}{\log_{b(p)} n - \log_{b(p)} \log_{b(p)} np - 10 \log_{b(p)} \ln n} \\ &\leq \frac{\log_{b(p_1)} n - \log_{b(p_1)} \log_{b(p)} np - 10 \log_{b(p_1)} \ln n}{\log_{b(p)} n - \log_{b(p)} \log_{b(p)} np - 10 \log_{b(p)} \ln n} \\ &= \frac{\log_{b(p_1)}(x)}{\log_{b(p)}(x)} = \frac{\ln b(p)}{\ln b(p_1)} \leq \frac{p(1+p)}{p_1} \leq 1 + 2\delta. \end{aligned}$$

This proves the second inequality. For the first, we take the reciprocal, and note that $(1+2\delta)^{-1} > 1-2\delta$. \square

Bound 3.2.

$$1 \geq \frac{\ell_1(p_2)}{\ell_1(p)} \geq 1 - 2\delta, \text{ and } 1 \leq \frac{\ell_1(p)}{\ell_1(p_2)} \leq 1 + 2\delta.$$

Proof Apply Bound 3.1 with p_2 in place of p and p in place of p_1 , since their relationships are the same. \square

Note now that if $d = n^\theta$ where $\theta = \Theta(1)$ then

$$\frac{\log_b \log_b np}{\log_b n} = \frac{\ln \log_b np}{\ln n} \approx 1 - \theta$$

which implies that

$$\ell_1(p) = (\theta + o(1)) \log_b np. \tag{3.6}$$

Bound 3.3.

$$(1 - p_1)^{\ell_1(p)} = \frac{\ell_1(p)(\ln n)^{10}}{(\theta + o(1))n} \text{ and } (1 - p_2)^{\ell_1(p)} = \frac{\ell_1(p)(\ln n)^{10}}{(\theta + o(1))n}.$$

Proof. It follows from Bound 3.1 that

$$(1 - p_1)^{\ell_1(p)} = (1 + o(1))(1 - p)^{\ell_1(p)} = (1 + o(1)) \frac{(\log_b np)(\log^{10} n)}{n}.$$

Now use (3.6). The proof for p_2 is similar. \square

3.2.4 Lemmas used for the lower bound in [6]

The strategy used in [6] to prove the lower bound relies on probabilistic assumptions labeled there as Lemmas 2.1 through 2.4. By assuming that those lemmas hold for random graphs (and occasionally referencing the proofs of the original lemmas), we prove that they hold in the random regular case as well. It follows that the lower bound of Theorem 1.1 is valid in the case of $G_{n,d}$ as well, provided our assumption that $d = n^\varepsilon$ holds.

Lemma 3.5 (Lemma 2.1 of [6]). *For every $S \subseteq [n]$ with $|S| = \ell_1(p)$, w.h.p.*

$$\ell_1(p)(\ln n)^9 \leq |\overline{N}(S)| \leq \ell_1(p)(\ln n)^{11}.$$

Proof. For an S as above, $\overline{N}(S) = \overline{N}_G(S) \subseteq \overline{N}_{H_1}(S)$. The distribution of $\overline{N}_{H_1}(S)$ is binomial with mean $n(1 - p_1)^{\ell_1(p)}$, which is at most $O(\ell_1(p)(\ln n)^{10})$ by Bound 3.3. We can use Chernoff bounds to get $|\overline{N}_{H_1}(S)| \leq \ell_1(p)(\ln n)^{11}$, which implies the same for $|\overline{N}(S)|$.

The proof of the lower bound is similar, except that we don't have the strict containment $\overline{N}_{H_2}(S) \subseteq \overline{N}_G(S)$. However, by Lemma 3.4, any vertex in S has $O(1)$ neighbors in G that it does not have in H_2 . Therefore $|\overline{N}_G(S)| \leq |\overline{N}_{H_2}(S)| + O(|S|)$. Because $|S| = \ell_1(p)$, and the Chernoff bound gives $|N_{H_2}| = \Omega(\ell_1(p)(\ln n)^{10})$ w.h.p., this difference will be absorbed in the $(1 + o(1))$ asymptotic factor. \square

Lemma 3.6 (Lemma 2.2 of [6]). *W.h.p. there do not exist $S, A, B \subseteq n$ such that (conditions omitted) and every $x \in B$ has fewer than $ap/2$ neighbors in A (where $a = |A|$).*

Proof. The proof of the corresponding lemma in [6] relies on the distribution to say that the number of neighbors any $x \in B$ has in A is distributed according to the binomial distribution $B(a, p)$, and uses the Chernoff bound $\mathbb{P}[B(a, p) \leq ap/2]^{b_1} \leq e^{-ab_1p/8}$.

The number of edges between x and A is bounded below by the number of such edges in the graph H_1 , which is distributed according to $B(a, p_1)$. So we replace the bound above by

$$\mathbb{P}[B(a, p_1) \leq ap/2] = \mathbb{P}\left[B(a, p_1) \leq \frac{ap_1(1 + \delta)}{2}\right] \leq \mathbb{P}\left[B(a, p_1) \leq ap_1\left(1 - \frac{1 - \delta}{2}\right)\right].$$

By the Chernoff bound, this is at most $e^{-ap_1(1 - \delta)^2/8} \leq e^{-(1 - o(1))ab_1p/8}$, and the argument of [6] still goes through. \square

Lemma 3.7 (Lemma 2.3 of [6]). *Let $a_1 = 2000\varepsilon^{-2}$ where ε is a small positive constant. W.h.p. there do not exist sets of vertices S, T_1, \dots, T_{a_1} such that (conditions omitted) and $N(S) \cap T_i = \emptyset$ for $i = 1, \dots, a$.*

Proof. If such sets exist in the graph G , then they will still exist when we lose some edges in passing to the graph H_1 . Examining the proof in [6] we see that all it requires is to consider the following factor which inflates the $o(1)$ probability estimate they use by:

$$\left(\frac{1-p_1}{1-p}\right)^{a_1 \binom{\varepsilon \ell_1(p)/21}{2}} \leq \left(\frac{1-p_1}{1-p}\right)^{3\ell_1(p)^2} = \left(1 + \frac{p\delta}{(1+\delta)(1-p)}\right)^{3\ell_1(p)^2} = 1 + o(1).$$

□

Lemma 3.8 (Lemma 2.4 of [6]). *Let $t = \frac{n}{\ell_1(p)(\ln n)^7}$. W.h.p. there do not exist pairwise disjoint sets of vertices S_1, \dots, S_t, U , such that (conditions omitted) and $|U \cap \overline{N}(S_i)| \leq \ell_1(p)(\ln n)^8$ for $i = 1, \dots, t$.*

Proof. Suppose such sets exist in the graph G . By Lemma 3.4 each vertex of S_i has at most $O(1)$ neighbors in G that are not in H_2 ; therefore in passing to the graph H_2 , $|U \cap \overline{N}(S_i)|$ will be at most $\ell_1(p)(\ln n)^8 + O(\ell_0(p))$ where each $|S_i| = \ell_0(p) = \ell_1(p) + C/p$ for some constant C (a fact we will use again). Since $\ell_1(p) = \frac{\ln n}{(\theta + o(1))p}$, the size of $|U \cap \overline{N}(S_i)|$ in H_2 is also $(1 + o(1))\ell_1(p)(\ln n)^8$. There is room in the argument of [6] to prove this lemma for intersections of this size as well. Therefore we will proceed by arguing that w.h.p. sets such as S_1, \dots, S_t, U do not exist in H_2 .

The proof in [6] hinges upon the claim that $(1-p)^{\ell_0(p)} = \Omega((1-p)^{\ell_1(p)})$. So it suffices to prove that $(1-p_2)^{\ell_0(p)} = \Omega((1-p)^{\ell_1(p)})$. We split $(1-p_2)^{\ell_0(p)}$ into factors $(1-p_2)^{\ell_1(p)}$ and $(1-p_2)^{C/p}$. By Bound 3.3, the first factor is $\Omega((1-p)^{\ell_1(p)})$. The second factor is no less than $(1-p_2)^{C/p_2}$, which stays in $(e^{-C}, 4^{-C})$ as p_2 ranges over $(0, 1/2]$, so it is effectively a constant. □

3.2.5 Lemmas used for the upper bound in [6]

As in the lower bound, the strategy used in [6] to prove the upper bound relies on some properties that hold w.h.p. in $G_{n,p}$. Again, we apply the Kim-Vu Sandwich Theorem [15, Theorem 2] to show that the same properties hold in $G_{n,d}$ as well.

The first player's strategy described in [6] is simple — she chooses an uncolored vertex with minimal number of available colors and then colors it with an arbitrary (available) color. We present some notation used in [6] during the analysis of the strategy. Given a (partial) coloring \mathcal{C} and a vertex v let $\alpha(v, \mathcal{C})$ be the number of available colors for v in \mathcal{C} . For a constant $\alpha > 3$ define

$$\beta_G = \alpha \frac{n(np)^{-1/\alpha}}{\log_b np}, \quad \gamma_G = \frac{10n \ln n}{\beta_G}$$

and

$$\mathcal{B}(\mathcal{C}) = \{v \mid \alpha(v) \leq \beta_G/2\}.$$

The first lemma states that there are not many vertices with few available colors. We show that the same is also true in $G_{n,d}$.

Lemma 3.9 (Lemma 3.1 of [6]). *W.h.p. for all collections \mathcal{C} ,*

$$|\mathcal{B}(\mathcal{C})| \leq \gamma.$$

Proof Here we can just use Remark 1. □

Lemma 3.10 (Lemma 3.2 of [6]). *W.h.p. every subset S of $G_{n,p}$ of size s spans at most $\phi = \phi(s) = (5ps + \ln n)s$ edges.*

Proof. The proof in [6] actually gives the result for $\phi_2 = (4.5ps + 0.9 \ln n)s$. Thus, applying Lemma 3.2 of [6] to H_2 gives that w.h.p. every set S of size s spans at most $(4.5p_2s + 0.9 \ln n)s$ edges. Since w.h.p. every vertex of G touches at most $O(1)$ edges not in H_2 , we have that w.h.p. the number of edges spanned by S in G is bounded by

$$(4.5p_2s + 0.9 \ln n + O(1))s = (4.5ps(1 + o(1)) + 0.9 \ln n + O(1))s \leq (5ps + \ln n)s$$

as required. □

This completes the proof of Theorem 1.3.

4 Theorem 1.4: Random Cubic Graphs

Consider the coloring game on $G_{n,3}$ with three colors. We describe a strategy for B that wins the game for him w.h.p., so $\chi_g(G_{n,3}) \geq 4$ w.h.p. This proves Theorem 1.4: in general $\chi_g(G) \leq \Delta(G) + 1$ where Δ denotes maximum degree, and in particular $\chi_g(G_{n,3}) \leq 4$.

We proceed in two steps. First, we describe a strategy for B that wins the game, given the existence of a subgraph H in G satisfying certain conditions. Next, we will prove that w.h.p., a random cubic graph contains such a subgraph.

4.1 The winning strategy

We will say that two vertices are *close* if they are connected by a path of length two or less, and that a path is *short* if some vertex on it is close to both endpoints. (This is not the same as being of length at most four). Vertices that are not close are *far apart* and a path that is not short is *long*. The motivation for this terminology is that coloring a vertex can only have an effect on vertices that are close to it; we will make this precise later on.

We first assume the existence of a subgraph H with the following properties (see Figure 1):




Figure 1: The subgraph H required for Bob's strategy on Random Cubic Graphs.

1. H consists of two vertices, v and w , together with three (internally disjoint) paths from one to the other.
2. Each of the paths consists of an even number of edges.
3. No two vertices in H are connected by a short path outside of H (in particular, H is induced).
4. The three paths themselves are all long.

In addition,

Property F: if A goes first, then the vertex colored by A on her first move is far from H .

B first plays on the vertex v . Provided A's next move is not on the vertex w , or on the neighbors of v or w , it is close to at most one of the three paths which make up H (this follows from Properties 3 and 4). The remaining two paths form a cycle containing v , with no other already colored vertices close to the cycle; by Property 2, the cycle is even. Call the vertices around the cycle $(v, v_1, v_2, \dots, v_{2k-1})$.

Starting from this even cycle, B proceeds as follows. He colors v_2 a different color from v ; this creates the threat that on his next move, he will color the third neighbor of v_1 the remaining color, leaving no way to color v_1 and winning. We will call such a move by B a *forcing move at v_1* . A can counter this threat in several ways:

- By coloring v_1 the only remaining viable color.
- By coloring v_1 's third neighbor the same color as either v or v_2 .

- By coloring that neighbor's other neighbors in the color different from both v and v_2 .

In all cases, A must color some vertex close to v_1 , that does not lie on the cycle.

B continues by making a forcing move at v_3 : coloring v_4 a different color from v_2 . Continuing to play on the even vertices v_{2i} , B makes forcing moves at each odd v_{2i-1} . By Property 3 of H , the set of vertices A must play on to counter each threat are disjoint; thus, A's response to each forcing move does not affect the rest of the strategy. By Property F, A's first play does not affect the strategy either.

When B colors v_{2k-2} , this is a forcing move both at v_{2k-3} and at v_{2k-1} (provided Bob chooses a color different both from v_{2k-4} and v). A cannot counter both threats, therefore B wins.

We now account for the remaining few cases. If A colors a neighbor of v or w on her second move, this vertex will be close to all three possible even cycles. However, we know that all three paths in H have even length. Therefore we can still apply this strategy to the even cycle not containing the vertex A colored. Even though it will be close to v or w , we will never need to force at v or at w , because we only force at odd numbered vertices along the path.

Finally, if A colors w itself, then there is no path we can choose that will avoid the vertex. Instead, B picks any of the paths from v to w , and makes forcing moves down that path. Provided that the path is sufficiently long to do so (which follows from Property 4), the final move will be a forcing move in two ways, winning the game for B once again.

4.2 Proof of the existence of H

It remains to show that the subgraph H exists w.h.p. (even allowing for A's first move). We will assume G is chosen by adding a random perfect matching to a cycle C on n vertices, and find H w.h.p. That this is a contiguous model to $G_{n,3}$ is well known, see [17]. In the following, let c be a constant; we will later see that we need c to be less than 1 for the proof to hold.

We begin by counting *good* segments of length $m = \lfloor c\sqrt{n} \rfloor$ on C , by which we mean those with no internal chords. First of all let X be twice the number of chords that intercept segments of length m or less – these are the only chords that could possibly be internal to a segment of the desired length. X can be written as the sum $X_1 + X_2 + \dots + X_n$, where X_i is the 0-1 indicator for the i -th vertex (call it v_i) to be the endpoint of such a chord. Also, let Y_i denote the length of the smaller of the two segments defined by v_i (this segment stretches from v_i to its partner). Thus

$$\mathbb{P}(Y_i = t) = \begin{cases} \frac{2}{n-1} & 2 \leq t \leq \lfloor (n-1)/2 \rfloor \\ \frac{1}{n-1} & t = n/2, n \text{ even} \end{cases}$$

Clearly $X_i = 1$ if and only if $Y_i \leq m$, and so

$$\mathbb{E}(X_i) = \frac{2m}{n-1} \text{ and } \mathbb{E}(X) = \frac{2mn}{n-1} \approx 2c\sqrt{n}.$$

In addition, $\text{Var}(X_i) \leq \mathbb{E}(X_i)$, and so

$$\text{Var}(X) = \sum_{i=1}^n \text{Var}(X_i) + \sum_{i \neq j} \text{Cov}(X_i, X_j) \leq \mathbb{E}(X) + \sum_{i \neq j} \text{Cov}(X_i, X_j).$$

Now

$$\begin{aligned} \text{Cov}(X_i, X_j) &= -\frac{4m^2}{(n-1)^2} + \sum_{t=2}^m \mathbb{P}(X_i = 1 \mid Y_j = t) \mathbb{P}(Y_j = t) \\ &\leq -\frac{4m^2}{(n-1)^2} + \frac{2m}{n-3} \cdot \frac{2m}{n-1} \\ &= \frac{8m^2}{(n-1)^2(n-3)}. \end{aligned}$$

Thus,

$$\text{Var}(X) \leq \mathbb{E}(X) + \frac{8m^2n}{(n-1)(n-3)} \approx \mathbb{E}(X).$$

By Chebyshev's inequality,

$$\mathbb{P}(|X - \mathbb{E}(X)| \leq \lambda \mathbb{E}(X)) \leq \frac{\text{Var}(X)}{\lambda^2 \mathbb{E}(X)^2} \leq \frac{2}{\lambda^2 c \sqrt{n}}.$$

Putting $\lambda = n^{-1/5}$ we see that w.h.p. $X \sim 2c\sqrt{n}$.

Consider the n different segments of length m on C . Each chord counted by X eliminates at most m of these segments as being good, which leaves $(1 - c^2)n$ segments remaining. We will want non-overlapping good segments; each good segment overlaps at most $2m$ other good segments and so we can assume that we can find $2n_1 \sim (c^{-1} - c)\sqrt{n}/2$ non-overlapping good segments w.h.p. Here the segments are $\sigma_1, \sigma_2, \dots, \sigma_{2n_1}$ are in clockwise order around C . We pair them together $P_i = (\sigma_i, \sigma_{n_1+i}), i = 1, 2, \dots, n_1$.

Pick any pair P_j . If there are exactly 3 chords from one segment to the other, as in Figure 2, then we will construct H as follows (assuming a_i and b_i are the endpoints of the chords, as labeled in Figure 2):

- Set v and w to be a_2 and b_2 , respectively.
- The first path from v to w is $(a_2, \dots, a_1, b_3, \dots, b_2)$, where the vertices in the ellipses are chosen along C .
- The second path from v to w is (a_2, b_1, \dots, b_2) .




Figure 2: A typical example of the subgraph H found in the Hamiltonian cycle model.

- The third path from v to w is (a_2, \dots, a_3, b_2) .

The paths given above require that the three chords are (a_1, b_3) , (a_2, b_1) , and (a_3, b_2) . In order for H to satisfy Properties 2 and 4, we impose conditions on the lengths of the paths (a_1, \dots, a_2) , (a_2, \dots, a_3) , (b_1, \dots, b_2) , and (b_2, \dots, b_3) : they must not be too small, and must have the right parity so that the three paths from v to w have even length. However, these conditions (and the condition that (a_2, b_2) must not be a chord) eliminate only a constant fraction of the possible chords; therefore there are $\Omega(m^6)$ ways to choose the chords.

The probability, then, that a subgraph H can be found between two given good segments, is at least

$$\Omega\left(\frac{m^6}{n^3}\right) \cdot \left(1 - \frac{m}{n - 2m}\right)^{2m} \quad (4.1)$$

where the last factor bounds the probability that there are no extra chords between the two segments. This tends to a constant ζ_1 that does not depend on n . Thus the expected number of j for which P_j has Properties 1,2 and 4 are satisfied is at least $\zeta_1 n_1$.

We now consider the number of pairs of good segments in which we can hope to find this structure. In order to ensure that, should a subgraph H be found, it satisfies Property F, we eliminate all pairs which contain a vertex close to the vertex A chooses on her first move – a constant number of pairs.

To ensure Property 3 we eliminate all pairs P_j in which two vertices have chords whose other endpoints are 1 or 2 edges apart. This happens with probability $O(1/n)$ for any two vertices, regardless of the disposition of the other chords incident with the segments in P_j . The pair P_j s contains $\binom{2m}{2} \leq 2c^2n$ pairs of vertices. Therefore with probability at least $(1 - O(1/n))^{2c^2n}$, which tends to a constant, ζ_2 say, a pair P_j satisfies Property 3.

Thus the expected number of j for which the pair P_j give rise to a copy of H satisfying all required properties is at least ζn_1 where $\zeta = \zeta_1 \zeta_2$. To prove concentration for the number of j we can simply use the Chebyshev inequality. This will work, because exposing the chords incident with a particular pair P_j will only have a small effect on the probability that any other P_j has the required properties. \square

References

- [1] D. Achlioptas and C. Moore, The chromatic number of random regular graphs, In *Proceedings of the 7th International Workshop on Approximation Algorithms for Combinatorial Optimization Problems and 8th International Workshop on Randomization and Computation, volume 3122 of Lecture Notes in Computer Science*, Springer (2004) 219–228.
- [2] D. Achlioptas and A. Naor, The two possible values of the chromatic number of a random graph, *Annals of Mathematics* 162 (2005) 1333–1349.
- [3] N. Alon and J. H. Spencer, **The probabilistic method**, 3rd Ed., Wiley, New York, 2008.
- [4] T. Bartnicki, J.A. Grytczuk, H.A. Kierstead and X. Zhu, The map coloring game, to appear in *American Mathematical Monthly*.
- [5] H.L. Bodlaender, On the complexity of some coloring games, *Internat. J. Found. Comput. Sci.* 2 (1991), 133–147.
- [6] T. Bohman, A.M. Frieze and B. Sudakov, The game chromatic number of random graphs, *Random Structures and Algorithms* 32 (2008) 223–235.
- [7] B. Bollobás, The chromatic number of random graphs, *Combinatorica* 8 (1988), 49–55.
- [8] B. Bollobás, A probabilistic proof of an asymptotic formula for the number of labelled regular graphs, *European Journal on Combinatorics* 1 (1980) 311–316.
- [9] C. Cooper, A.M. Frieze and B. Reed, Random regular graphs of non-constant degree: connectivity and Hamilton cycles, *Combinatorics, Probability and Computing* 11 (2002) 249–262.

- [10] U. Faigle, U. Kern, H. Kierstead and W.T. Trotter, On the Game Chromatic Number of some Classes of Graphs, *Ars Combinatoria* 35 (1993) 143–150.
- [11] A.M. Frieze and T. Łuczak, On the independence and chromatic numbers of random regular graphs, *Journal of Combinatorial Theory B* 54 (1992) 123–132.
- [12] M. Gardner, Mathematical Games, *Scientific American* 244 (1981), no. 4, 18–26.
- [13] S. Janson, T. Łuczak and A. Ruciński, **Random Graphs**, John Wiley and Sons, New York, 2000.
- [14] G. Kemkes, X. Pérez-Giménez and N. Wormald, On the chromatic number of random d -regular graphs, *Advances in Mathematics* 223 (2010), 300–328.
- [15] J.H. Kim and V.H. Vu. Sandwiching random graphs: universality between random graph models. *Advances in Mathematics* 188 (2004): 444–469.
- [16] T. Łuczak, The chromatic number of random graphs, *Combinatorica* 11 (1991), 45–54.
- [17] N. Wormald, Models of random regular graphs, *Surveys in Combinatorics*, J.D. Lamb and D.A. Preece, eds., London Mathematical Society Lecture Note Series, 276 (1999) 239–298. Cambridge University Press, Cambridge, 1999