

Simplex slicing: an asymptotically-sharp lower bound


Colin Tang

cstang@andrew.cmu.edu

April 12, 2024


Our goal

Let Δ_n denote the regular n -simplex.



Our goal

Let Δ_n denote the regular n -simplex.



Main question

How may we choose a 1-codimensional hyperplane H passing through the center of Δ_n , so that the volume of the intersection $\text{vol}_{n-1}(\Delta_n \cap H)$ is *minimized*?

Motivation

If K is a convex body, we call a set of the form $K \cap H$ (where H is a 1-codimensional hyperplane) a *section* of K .

Motivation

If K is a convex body, we call a set of the form $K \cap H$ (where H is a 1-codimensional hyperplane) a *section* of K . If H passes through the barycenter of K , we call it a *central section*.

Motivation

If K is a convex body, we call a set of the form $K \cap H$ (where H is a 1-codimensional hyperplane) a *section* of K . If H passes through the barycenter of K , we call it a *central section*.

Bourgain's slicing problem

Does every convex body K of volume 1 admit a section whose volume is at least some universal constant, independent of the dimension n ?

Motivation

If K is a convex body, we call a set of the form $K \cap H$ (where H is a 1-codimensional hyperplane) a *section* of K . If H passes through the barycenter of K , we call it a *central section*.

Bourgain's slicing problem

Does every convex body K of volume 1 admit a section whose volume is at least some universal constant, independent of the dimension n ?

- ▶ Open problem

Motivation

If K is a convex body, we call a set of the form $K \cap H$ (where H is a 1-codimensional hyperplane) a *section* of K . If H passes through the barycenter of K , we call it a *central section*.

Bourgain's slicing problem

Does every convex body K of volume 1 admit a section whose volume is at least some universal constant, independent of the dimension n ?

- ▶ Open problem
- ▶ Key to understanding the uniform distribution on a high-dimensional convex body

Motivation

If K is a convex body, we call a set of the form $K \cap H$ (where H is a 1-codimensional hyperplane) a *section* of K . If H passes through the barycenter of K , we call it a *central section*.

Bourgain's slicing problem

Does every convex body K of volume 1 admit a section whose volume is at least some universal constant, independent of the dimension n ?

- ▶ Open problem
- ▶ Key to understanding the uniform distribution on a high-dimensional convex body
- ▶ Connections to isoperimetry in high dimensions (cf. *KLS conjecture*)

Previous work

A general type of question

Given a specific convex body K , can we identify its minimum central section?

¹Hugo Hadwiger. "Gitterperiodische Punktmengen und Isoperimetrie". In: *Monatshefte für Mathematik* 76.5 (1972), pp. 410–418.

²Douglas Hensley. "Slicing the Cube in \mathbb{R}^n and Probability (Bounds for the Measure of a Central Cube Slice in \mathbb{R}^n by Probability Methods)". In: *Proceedings of the American Mathematical Society* 73.1 (1979), pp. 95–100.

³Keith Ball. "Cube slicing in \mathbb{R}^n ". In: *Proceedings of the American Mathematical Society* 97.3 (1986), pp. 465–473.

⁴Simon Webb. "Central slices of the regular simplex". In: *Geometriae Dedicata* 61.1 (1996), pp. 19–28.

Previous work

A general type of question

Given a specific convex body K , can we identify its minimum central section? Maximum central section?

¹Hadwiger, "Gitterperiodische Punktmengen und Isoperimetrie".

²Hensley, "Slicing the Cube in \mathbb{R}^n and Probability (Bounds for the Measure of a Central Cube Slice in \mathbb{R}^n by Probability Methods)".

³Ball, "Cube slicing in \mathbb{R}^n ".

⁴Webb, "Central slices of the regular simplex".

Previous work

A general type of question

Given a specific convex body K , can we identify its minimum central section? Maximum central section?

- ▶ $K = Q_n$ (n -dimensional hypercube), minimal central section identified in [Hadwiger 1972¹, Hensley 1979²]

¹Hadwiger, "Gitterperiodische Punktmengen und Isoperimetrie".

²Hensley, "Slicing the Cube in \mathbb{R}^n and Probability (Bounds for the Measure of a Central Cube Slice in \mathbb{R}^n by Probability Methods)".

³Ball, "Cube slicing in \mathbb{R}^n ".

⁴Webb, "Central slices of the regular simplex".

Previous work

A general type of question

Given a specific convex body K , can we identify its minimum central section? Maximum central section?

- ▶ $K = Q_n$ (n -dimensional hypercube), minimal central section identified in [Hadwiger 1972¹, Hensley 1979²]
- ▶ $K = Q_n$, maximal central section identified in [Ball 1986³]

¹Hadwiger, "Gitterperiodische Punktmengen und Isoperimetrie".

²Hensley, "Slicing the Cube in \mathbb{R}^n and Probability (Bounds for the Measure of a Central Cube Slice in \mathbb{R}^n by Probability Methods)".

³Ball, "Cube slicing in \mathbb{R}^n ".

⁴Webb, "Central slices of the regular simplex".

Previous work

A general type of question

Given a specific convex body K , can we identify its minimum central section? Maximum central section?

- ▶ $K = Q_n$ (n -dimensional hypercube), minimal central section identified in [Hadwiger 1972¹, Hensley 1979²]
- ▶ $K = Q_n$, maximal central section identified in [Ball 1986³]
- ▶ $K = \Delta_n$ (n -dimensional regular simplex), maximal central section identified in [Webb 1996⁴]

¹Hadwiger, "Gitterperiodische Punktmengen und Isoperimetrie".

²Hensley, "Slicing the Cube in \mathbb{R}^n and Probability (Bounds for the Measure of a Central Cube Slice in \mathbb{R}^n by Probability Methods)".

³Ball, "Cube slicing in \mathbb{R}^n ".

⁴Webb, "Central slices of the regular simplex".

Simplex minimum

This leaves open the question from the beginning:

Simplex minimum

What is the minimum central section of the regular simplex?

⁵Patryk Brzezinski. "Volume estimates for sections of certain convex bodies". In: *Mathematische Nachrichten* 286.17-18 (2013), pp. 1726–1743.

Simplex minimum

This leaves open the question from the beginning:

Simplex minimum

What is the minimum central section of the regular simplex?

Conjecture

The minimum central section is the central section $\Delta_n \cap H_{\text{facet}}$ that's parallel to a facet.

⁵Brzezinski, "Volume estimates for sections of certain convex bodies".

Simplex minimum

This leaves open the question from the beginning:

Simplex minimum

What is the minimum central section of the regular simplex?

Conjecture

The minimum central section is the central section $\Delta_n \cap H_{\text{facet}}$ that's parallel to a facet.

Previous best bound [Brzezinski 2013⁵]

The central section $\Delta_n \cap H_{\text{facet}}$ is within a factor of $\frac{2\sqrt{3}}{e} \approx 1.27$ of the minimum.

⁵Brzezinski, "Volume estimates for sections of certain convex bodies".

Main result

Conjecture is true up to a $1 - o(1)$ factor [T. 2024+⁶]

The central section $\Delta_n \cap H_{\text{facet}}$ is within a factor of $1 - o(1)$ of the minimum. (Little o is with respect to the dimension n .)

⁶Colin Tang. *Simplex slicing: an asymptotically-sharp lower bound*. 2024. arXiv: 2403.13224 [math.MG].

Main result

Conjecture is true up to a $1 - o(1)$ factor [T. 2024+⁶]

The central section $\Delta_n \cap H_{\text{facet}}$ is within a factor of $1 - o(1)$ of the minimum. (Little o is with respect to the dimension n .)

Tools used:

⁶Tang, *Simplex slicing: an asymptotically-sharp lower bound*.

Main result

Conjecture is true up to a $1 - o(1)$ factor [T. 2024+⁶]

The central section $\Delta_n \cap H_{\text{facet}}$ is within a factor of $1 - o(1)$ of the minimum. (Little o is with respect to the dimension n .)

Tools used:

- ▶ ***probability distributions***

⁶Tang, *Simplex slicing: an asymptotically-sharp lower bound*.

Main result

Conjecture is true up to a $1 - o(1)$ factor [T. 2024+⁶]

The central section $\Delta_n \cap H_{\text{facet}}$ is within a factor of $1 - o(1)$ of the minimum. (Little o is with respect to the dimension n .)

Tools used:

- ▶ ***probability distributions***
- ▶ ***Fourier analysis***

⁶Tang, *Simplex slicing: an asymptotically-sharp lower bound*.

Main result

Conjecture is true up to a $1 - o(1)$ factor [T. 2024+⁶]

The central section $\Delta_n \cap H_{\text{facet}}$ is within a factor of $1 - o(1)$ of the minimum. (Little o is with respect to the dimension n .)

Tools used:

- ▶ ***probability distributions***
- ▶ ***Fourier analysis***
- ▶ New: ***moving the contour of integration of a meromorphic function***

⁶Tang, *Simplex slicing: an asymptotically-sharp lower bound*.

Main result

Conjecture is true up to a $1 - o(1)$ factor [T. 2024+⁶]

The central section $\Delta_n \cap H_{\text{facet}}$ is within a factor of $1 - o(1)$ of the minimum. (Little o is with respect to the dimension n .)

Tools used:

- ▶ ***probability distributions***
- ▶ ***Fourier analysis***
- ▶ New: ***moving the contour of integration of a meromorphic function***

We'll prove this result in the remainder of the presentation.

⁶Tang, *Simplex slicing: an asymptotically-sharp lower bound*.

Tool: probability distributions

Embed Δ_n into \mathbb{R}^{n+1} via

$$\Delta_n = \left\{ (x_1, x_2, \dots, x_{n+1}) \in \mathbb{R}^{n+1} \mid \begin{array}{l} x_1 + x_2 + \dots + x_{n+1} = 1 \\ x_i \geq 0 \text{ for each } i \end{array} \right\}.$$

Tool: probability distributions

Embed Δ_n into \mathbb{R}^{n+1} via

$$\Delta_n = \left\{ (x_1, x_2, \dots, x_{n+1}) \in \mathbb{R}^{n+1} \mid \begin{array}{l} x_1 + x_2 + \dots + x_{n+1} = 1 \\ x_i \geq 0 \text{ for each } i \end{array} \right\}.$$

Central sections $\Delta_n \cap H$ correspond to a choice of vector a with

$$\begin{cases} a_1 + a_2 + \dots + a_{n+1} = 0 \\ a_1^2 + a_2^2 + \dots + a_{n+1}^2 = 1 \end{cases}$$

where a is the normal vector to H .

Tool: probability distributions

Embed Δ_n into \mathbb{R}^{n+1} via

$$\Delta_n = \left\{ (x_1, x_2, \dots, x_{n+1}) \in \mathbb{R}^{n+1} \mid \begin{array}{l} x_1 + x_2 + \dots + x_{n+1} = 1 \\ x_i \geq 0 \text{ for each } i \end{array} \right\}.$$

Central sections $\Delta_n \cap H$ correspond to a choice of vector a with

$$\begin{cases} a_1 + a_2 + \dots + a_{n+1} = 0 \\ a_1^2 + a_2^2 + \dots + a_{n+1}^2 = 1 \end{cases}$$

where a is the normal vector to H .

Idea: Instead of Δ_n , consider the density

$$\Phi(x_1, x_2, \dots, x_{n+1}) = \begin{cases} e^{-x_1 - x_2 - \dots - x_{n+1}} & \text{if each } x_i \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

Tool: probability distributions

Embed Δ_n into \mathbb{R}^{n+1} via

$$\Delta_n = \left\{ (x_1, x_2, \dots, x_{n+1}) \in \mathbb{R}^{n+1} \mid \begin{array}{l} x_1 + x_2 + \dots + x_{n+1} = 1 \\ x_i \geq 0 \text{ for each } i \end{array} \right\}.$$

Central sections $\Delta_n \cap H$ correspond to a choice of vector a with

$$\begin{cases} a_1 + a_2 + \dots + a_{n+1} = 0 \\ a_1^2 + a_2^2 + \dots + a_{n+1}^2 = 1 \end{cases}$$

where a is the normal vector to H .

Idea: Instead of Δ_n , consider the density

$$\Phi(x_1, x_2, \dots, x_{n+1}) = \begin{cases} e^{-x_1 - x_2 - \dots - x_{n+1}} & \text{if each } x_i \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

Then $\int_{a^\perp} \Phi d\mathcal{H}^n$ is proportional to the volume of the section.

Tool: probability distributions

Embed Δ_n into \mathbb{R}^{n+1} via

$$\Delta_n = \left\{ (x_1, x_2, \dots, x_{n+1}) \in \mathbb{R}^{n+1} \mid \begin{array}{l} x_1 + x_2 + \dots + x_{n+1} = 1 \\ x_i \geq 0 \text{ for each } i \end{array} \right\}.$$

Central sections $\Delta_n \cap H$ correspond to a choice of vector a with

$$\begin{cases} a_1 + a_2 + \dots + a_{n+1} = 0 \\ a_1^2 + a_2^2 + \dots + a_{n+1}^2 = 1 \end{cases}$$

where a is the normal vector to H .

Idea: Instead of Δ_n , consider the density

$$\Phi(x_1, x_2, \dots, x_{n+1}) = \begin{cases} e^{-x_1 - x_2 - \dots - x_{n+1}} & \text{if each } x_i \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

Then $\int_{a^\perp} \Phi d\mathcal{H}^n$ is proportional to the volume of the section.

Minimum central sections correspond to minimizing $\int_{a^\perp} \Phi d\mathcal{H}^n$.

Tool: probability distributions

But Φ is a product measure, so $\int_{a^\perp} \Phi d\mathcal{H}^n$ is the density at 0 of the random variable

$$Z_a := a_1 Y_1 + a_2 Y_2 + \cdots + a_{n+1} Y_{n+1}$$


(where the Y_i are i.i.d. standard exponentials (mean 1)).

Tool: probability distributions

But Φ is a product measure, so $\int_{a^\perp} \Phi d\mathcal{H}^n$ is the density at 0 of the random variable

$$Z_a := a_1 Y_1 + a_2 Y_2 + \cdots + a_{n+1} Y_{n+1}$$

(where the Y_i are i.i.d. standard exponentials (mean 1)).




Tool: probability distributions

But Φ is a product measure, so $\int_{a^\perp} \Phi d\mathcal{H}^n$ is the density at 0 of the random variable

$$Z_a := a_1 Y_1 + a_2 Y_2 + \cdots + a_{n+1} Y_{n+1}$$

(where the Y_i are i.i.d. standard exponentials (mean 1)).



Let $G_a(x)$ denote the density of Z_a , so what we said above is $\int_{a^\perp} \Phi d\mathcal{H}^n = G_a(0)$.

Tool: probability distributions

Reduction

The minimum central section corresponds to a choice of vector a minimizing $G_a(0)$.

Tool: probability distributions

Reduction

The minimum central section corresponds to a choice of vector a minimizing $G_a(0)$. Conjectured minimizer a_{facet} satisfies

$$G_{a_{\text{facet}}}(0) = \sqrt{\frac{n}{n+1}} \left(\frac{n}{n+1}\right)^{n-1} \approx \frac{1}{e}.$$

Tool: probability distributions

Reduction

The minimum central section corresponds to a choice of vector a minimizing $G_a(0)$. Conjectured minimizer a_{facet} satisfies

$$G_{a_{\text{facet}}}(0) = \sqrt{\frac{n}{n+1}} \left(\frac{n}{n+1}\right)^{n-1} \approx \frac{1}{e}.$$

It's hard to optimize over the set $\mathcal{S}^n \cap \mathbf{1}^\perp$ (the feasible region of a).

Tool: probability distributions

Reduction

The minimum central section corresponds to a choice of vector a minimizing $G_a(0)$. Conjectured minimizer a_{facet} satisfies

$$G_{a_{\text{facet}}}(0) = \sqrt{\frac{n}{n+1}} \left(\frac{n}{n+1}\right)^{n-1} \approx \frac{1}{e}.$$

It's hard to optimize over the set $\mathcal{S}^n \cap \mathbf{1}^\perp$ (the feasible region of a). Expand the feasible region:

Tool: probability distributions

Reduction

The minimum central section corresponds to a choice of vector a minimizing $G_a(0)$. Conjectured minimizer a_{facet} satisfies

$$G_{a_{\text{facet}}}(0) = \sqrt{\frac{n}{n+1}} \left(\frac{n}{n+1}\right)^{n-1} \approx \frac{1}{e}.$$

It's hard to optimize over the set $\mathcal{S}^n \cap \mathbf{1}^\perp$ (the feasible region of a). Expand the feasible region:

- ▶ Let $u \in \mathcal{S}^n$ be arbitrary (the feasible region of u has one fewer constraint than that of a !).

Tool: probability distributions

Reduction

The minimum central section corresponds to a choice of vector a minimizing $G_a(0)$. Conjectured minimizer a_{facet} satisfies

$$G_{a_{\text{facet}}}(0) = \sqrt{\frac{n}{n+1}} \left(\frac{n}{n+1}\right)^{n-1} \approx \frac{1}{e}.$$

It's hard to optimize over the set $\mathcal{S}^n \cap \mathbf{1}^\perp$ (the feasible region of a). Expand the feasible region:

- ▶ Let $u \in \mathcal{S}^n$ be arbitrary (the feasible region of u has one fewer constraint than that of a !).
- ▶ Define $Z_u := u_1(Y_1 - 1) + u_2(Y_2 - 1) + \cdots + u_{n+1}(Y_{n+1} - 1)$.

Tool: probability distributions

Reduction

The minimum central section corresponds to a choice of vector a minimizing $G_a(0)$. Conjectured minimizer a_{facet} satisfies

$$G_{a_{\text{facet}}}(0) = \sqrt{\frac{n}{n+1}} \left(\frac{n}{n+1}\right)^{n-1} \approx \frac{1}{e}.$$

It's hard to optimize over the set $\mathcal{S}^n \cap \mathbf{1}^\perp$ (the feasible region of a). Expand the feasible region:

- ▶ Let $u \in \mathcal{S}^n$ be arbitrary (the feasible region of u has one fewer constraint than that of a !).
- ▶ Define $Z_u := u_1(Y_1 - 1) + u_2(Y_2 - 1) + \cdots + u_{n+1}(Y_{n+1} - 1)$.
- ▶ This extends the earlier definition of Z_a since

$$\begin{aligned} & a_1(Y_1 - 1) + a_2(Y_2 - 1) + \cdots + a_{n+1}(Y_{n+1} - 1) \\ &= a_1 Y_1 + a_2 Y_2 + \cdots + a_{n+1} Y_{n+1} - (a_1 + a_2 + \cdots + a_{n+1}) \\ &= a_1 Y_1 + a_2 Y_2 + \cdots + a_{n+1} Y_{n+1}. \end{aligned}$$

Tool: probability distributions

Question

What's the minimum possible value that $G_u(0)$ can attain, as u varies in \mathcal{S}^n ?

Tool: probability distributions

Question

What's the minimum possible value that $G_u(0)$ can attain, as u varies in \mathcal{S}^n ?

Our result

$G_u(0) \geq \frac{1}{e}$ for each $u \in \mathcal{S}^n$. Equality achieved if $u = (1) \in \mathcal{S}^0$.

Tool: probability distributions

Question

What's the minimum possible value that $G_u(0)$ can attain, as u varies in \mathcal{S}^n ?

Our result

$G_u(0) \geq \frac{1}{e}$ for each $u \in \mathcal{S}^n$. Equality achieved if $u = (\mathbf{1}) \in \mathcal{S}^0$.

We lost a bit by expanding the feasible region from $\mathcal{S}^n \cap \mathbf{1}^\perp \ni a$ to $\mathcal{S}^n \ni u$.

Tool: probability distributions

Question

What's the minimum possible value that $G_u(0)$ can attain, as u varies in \mathcal{S}^n ?

Our result

$G_u(0) \geq \frac{1}{e}$ for each $u \in \mathcal{S}^n$. Equality achieved if $u = (1) \in \mathcal{S}^0$.

We lost a bit by expanding the feasible region from $\mathcal{S}^n \cap \mathbf{1}^\perp \ni a$ to $\mathcal{S}^n \ni u$. Indeed, the minimum over u of $G_u(0)$ is exactly $\frac{1}{e}$, but we think the minimum over a of $G_a(0)$ is given by

$$G_{a_{\text{facet}}}(0) = \sqrt{\frac{n}{n+1}} \left(\frac{n}{n+1}\right)^{n-1}.$$

Tool: probability distributions

Question

What's the minimum possible value that $G_u(0)$ can attain, as u varies in \mathcal{S}^n ?

Our result

$G_u(0) \geq \frac{1}{e}$ for each $u \in \mathcal{S}^n$. Equality achieved if $u = (1) \in \mathcal{S}^0$.

We lost a bit by expanding the feasible region from $\mathcal{S}^n \cap \mathbf{1}^\perp \ni a$ to $\mathcal{S}^n \ni u$. Indeed, the minimum over u of $G_u(0)$ is exactly $\frac{1}{e}$, but we think the minimum over a of $G_a(0)$ is given by

$$G_{a_{\text{facet}}}(0) = \sqrt{\frac{n}{n+1}} \left(\frac{n}{n+1}\right)^{n-1}.$$

But certainly

$$\frac{1}{e} = \min_u G_u(0) \leq \min_a G_a(0) \leq G_{a_{\text{facet}}}(0),$$

and since $G_{a_{\text{facet}}}(0) = \frac{1}{e}(1 + o(1))$, we lost at most a $1 + o(1)$ factor by expanding the feasible region.

Tool: Fourier analysis

$G_u(x)$ is the density of a sum of independent centered exponentials $u_j(Y_j - 1)$, so G_u is a convolution $f_1 * f_2 * \cdots * f_{n+1}$.

Tool: Fourier analysis

$G_u(x)$ is the density of a sum of independent centered exponentials $u_j(Y_j - 1)$, so G_u is a convolution $f_1 * f_2 * \cdots * f_{n+1}$.

Here, $f_j(x)$ is the density of $u_j(Y_j - 1)$. It's given by

$f_j(x) = \frac{1}{|u_j|} f\left(\frac{x}{u_j} + 1\right)$ where f is the density of the standard (uncentered) exponential with mean 1:

$$f(x) = \begin{cases} e^{-x} & \text{if } x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

Tool: Fourier analysis

Take the Fourier transform. Convolution becomes pointwise multiplication.

Tool: Fourier analysis

Take the Fourier transform. Convolution becomes pointwise multiplication.

$$\hat{f}(t) = \frac{1}{1 + it}$$

Tool: Fourier analysis

Take the Fourier transform. Convolution becomes pointwise multiplication.

$$\hat{f}(t) = \frac{1}{1 + it}$$

$$\hat{f}_j(t) = \frac{e^{iu_j t}}{1 + iu_j t}$$

Tool: Fourier analysis

Take the Fourier transform. Convolution becomes pointwise multiplication.

$$\hat{f}(t) = \frac{1}{1 + it}$$

$$\hat{f}_j(t) = \frac{e^{iu_j t}}{1 + iu_j t}$$

$$\widehat{G}_u(t) = \prod_{j=1}^{n+1} \hat{f}_j(t) = \prod_{j=1}^{n+1} \frac{e^{iu_j t}}{1 + iu_j t}$$

Tool: Fourier analysis

Take the Fourier transform. Convolution becomes pointwise multiplication.

$$\hat{f}(t) = \frac{1}{1 + it}$$

$$\hat{f}_j(t) = \frac{e^{iu_j t}}{1 + iu_j t}$$

$$\widehat{G}_u(t) = \prod_{j=1}^{n+1} \hat{f}_j(t) = \prod_{j=1}^{n+1} \frac{e^{iu_j t}}{1 + iu_j t}$$

Fourier inversion formula, valid if u has at least two nonzero entries:

$$G_u(0) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \widehat{G}_u(t) dt = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \prod_{j=1}^{n+1} \frac{e^{iu_j t}}{1 + iu_j t} dt$$

Tool: Fourier analysis

We wanted to show $G_u(0) \geq \frac{1}{e}$, and this is equivalent to

$$\frac{1}{2\pi} \int_{-\infty}^{+\infty} \prod_{j=1}^{n+1} \frac{e^{iu_j t}}{1 + iu_j t} dt \geq \frac{1}{e}.$$

Tool: Fourier analysis

We wanted to show $G_u(0) \geq \frac{1}{e}$, and this is equivalent to

$$\frac{1}{2\pi} \int_{-\infty}^{+\infty} \prod_{j=1}^{n+1} \frac{e^{iu_j t}}{1 + iu_j t} dt \geq \frac{1}{e}.$$

Letting $F_u(t) := \prod_{j=1}^{n+1} \frac{e^{iu_j t}}{1 + iu_j t}$, we just want to show

$$\frac{1}{2\pi} \int_{-\infty}^{+\infty} F_u(t) dt \geq \frac{1}{e}.$$

Some complex analysis

Thus far, all the techniques have been known.

Some complex analysis

Thus far, all the techniques have been known.

The main difficulty now is estimating the highly oscillatory integral

$$\int_{-\infty}^{+\infty} F_u(t) dt.$$

Some complex analysis

Thus far, all the techniques have been known.

The main difficulty now is estimating the highly oscillatory integral

$$\int_{-\infty}^{+\infty} F_u(t) dt.$$

I'll spare you the pictures from my first attempt. It really wasn't great.

Tool: moving the contour of integration

New idea: *moving the contour of integration.*

Tool: moving the contour of integration

New idea: ***moving the contour of integration.***

- ▶ Recall from complex analysis that the integral of a meromorphic function doesn't depend on the path taken (with some caveats).

Tool: moving the contour of integration

New idea: ***moving the contour of integration.***

- ▶ Recall from complex analysis that the integral of a meromorphic function doesn't depend on the path taken (with some caveats).
- ▶ If we want to estimate the integral $\int_{-\infty}^{+\infty} F_u(t) dt$, we can change the contour of integration, from the real line, to a special curve γ_u .


Tool: moving the contour of integration

New idea: ***moving the contour of integration.***

- ▶ Recall from complex analysis that the integral of a meromorphic function doesn't depend on the path taken (with some caveats).
- ▶ If we want to estimate the integral $\int_{-\infty}^{+\infty} F_u(t) dt$, we can change the contour of integration, from the real line, to a special curve γ_u .
- ▶ We will choose γ_u to have the property that F_u is always a positive real number along γ_u .


Tool: moving the contour of integration

Here's a plot of $F_u(t)$ with $u = (\sqrt{0.42}, \sqrt{0.38}, \sqrt{0.20})$:



Tool: moving the contour of integration

Here's a plot of $F_u(t)$ with $u = (\sqrt{0.42}, \sqrt{0.38}, \sqrt{0.20})$:



The color denotes the argument of $F_u(t)$. Red means real. Follow the red color, trace out a curve γ_u .

Tool: moving the contour of integration

Black box (basically just the Implicit Function Theorem)

We can always find such a curve γ_u , along which F_u takes positive real values, such that γ_u is \mathcal{C}^∞ and passes through the origin.

Moreover, γ_u can be viewed as the graph of an even function $y_u(x)$ in the xy -plane (identified with the complex plane in the usual manner).


Tool: moving the contour of integration

Black box (basically just the Implicit Function Theorem)

We can always find such a curve γ_u , along which F_u takes positive real values, such that γ_u is C^∞ and passes through the origin.

Moreover, γ_u can be viewed as the graph of an even function $y_u(x)$ in the xy -plane (identified with the complex plane in the usual manner).

Here's a plot of γ_u with the same u ($u = (\sqrt{0.42}, \sqrt{0.38}, \sqrt{0.20})$):



Tool: moving the contour of integration

Black box (some crude tail bounds)

As long as u has at least two nonzero entries, we have that the integral $\int_{-\infty}^{+\infty} F_u(t) dt$ exists and equals $\int_{\gamma_u} F_u(t) dt$. Moreover, the integrand $F_u(t)$ is always a positive real number if t is on γ_u .

Tool: moving the contour of integration

Black box (some crude tail bounds)

As long as u has at least two nonzero entries, we have that the integral $\int_{-\infty}^{+\infty} F_u(t) dt$ exists and equals $\int_{\gamma_u} F_u(t) dt$. Moreover, the integrand $F_u(t)$ is always a positive real number if t is on γ_u .

This is the part when we actually ***move the contour of integration***.

Tool: moving the contour of integration

Black box (some crude tail bounds)

As long as u has at least two nonzero entries, we have that the integral $\int_{-\infty}^{+\infty} F_u(t) dt$ exists and equals $\int_{\gamma_u} F_u(t) dt$. Moreover, the integrand $F_u(t)$ is always a positive real number if t is on γ_u .

This is the part when we actually ***move the contour of integration***.

So we just need to estimate $\int_{\gamma_u} F_u(t) dt$.

Differential equations

Recall that $y_u(x)$ is the function whose graph is γ_u .

Differential equations

Recall that $y_u(x)$ is the function whose graph is γ_u . Defining $\tilde{F}_u(x) := F_u(x + iy_u(x))$, we can compute that $\int_{\gamma_u} F_u(t) dt = \int_{\mathbb{R}} \tilde{F}_u(x) dx$ since \tilde{F}_u is an even function of x .

Differential equations

Recall that $y_u(x)$ is the function whose graph is γ_u . Defining $\tilde{F}_u(x) := F_u(x + iy_u(x))$, we can compute that $\int_{\gamma_u} F_u(t) dt = \int_{\mathbb{R}} \tilde{F}_u(x) dx$ since \tilde{F}_u is an even function of x . (We changed dt to dx .)

Differential equations

Recall that $y_u(x)$ is the function whose graph is γ_u . Defining $\tilde{F}_u(x) := F_u(x + iy_u(x))$, we can compute that $\int_{\gamma_u} F_u(t) dt = \int_{\mathbb{R}} \tilde{F}_u(x) dx$ since \tilde{F}_u is an even function of x . (We changed dt to dx .)

So we just need to show $\frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_u(x) dx \geq \frac{1}{e}$.

Differential equations

Compute that equality holds if $u = (1) \in \mathcal{S}^0$; i.e.

$$\frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_{(1)}(x) dx = \frac{1}{e}.$$

Differential equations

Compute that equality holds if $u = (1) \in \mathcal{S}^0$; i.e.

$$\frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_{(1)}(x) dx = \frac{1}{e}.$$

If we could show $\tilde{F}_u(x) \geq \tilde{F}_{(1)}(x)$ for each x , then we would automatically get

$$\frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_u(x) dx \geq \frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_{(1)}(x) dx = \frac{1}{e}$$

as desired.

Differential equations

Compute that equality holds if $u = (1) \in \mathcal{S}^0$; i.e.

$$\frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_{(1)}(x) dx = \frac{1}{e}.$$

If we could show $\tilde{F}_u(x) \geq \tilde{F}_{(1)}(x)$ for each x , then we would automatically get

$$\frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_u(x) dx \geq \frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_{(1)}(x) dx = \frac{1}{e}$$

as desired.

Let's show the boxed statement.

Differential equations

Compute that equality holds if $u = (1) \in \mathcal{S}^0$; i.e.

$$\frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_{(1)}(x) dx = \frac{1}{e}.$$

If we could show $\tilde{F}_u(x) \geq \tilde{F}_{(1)}(x)$ for each x , then we would automatically get

$$\frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_u(x) dx \geq \frac{1}{2\pi} \int_{\mathbb{R}} \tilde{F}_{(1)}(x) dx = \frac{1}{e}$$

as desired.

Let's show the boxed statement. From now on, assume $x > 0$.

Differential equations

Defining property of y_u

$$y'_u = \frac{\sum_{j=1}^{n+1} \frac{-y_u + u_j(x^2 + y_u^2)}{x^2 + (\frac{1}{u_j} - y_u)^2}}{\sum_{j=1}^{n+1} \frac{x}{x^2 + (\frac{1}{u_j} - y_u)^2}}$$

Differential equations

Defining property of y_u

$$y'_u = \frac{\sum_{j=1}^{n+1} \frac{-y_u + u_j(x^2 + y_u^2)}{x^2 + (\frac{1}{u_j} - y_u)^2}}{\sum_{j=1}^{n+1} \frac{x}{x^2 + (\frac{1}{u_j} - y_u)^2}}$$

Corollary

$$y'_u \leq \frac{-y_u + x^2 + y_u^2}{x}$$

Differential equations

Defining property of y_u

$$y'_u = \frac{\sum_{j=1}^{n+1} \frac{-y_u + u_j(x^2 + y_u^2)}{x^2 + (\frac{1}{u_j} - y_u)^2}}{\sum_{j=1}^{n+1} \frac{x}{x^2 + (\frac{1}{u_j} - y_u)^2}}$$

Corollary

$$y'_u \leq \frac{-y_u + x^2 + y_u^2}{x}$$

Using this, we can prove

Black box

$$-y_{(1)} \leq y_u \leq y_{(1)} \text{ for all } x > 0. \quad (*)$$

Differential equations

Compute

$$\frac{d}{dx} \log \tilde{F}_u(x) = - \frac{\left(\sum_{j=1}^{n+1} \frac{x}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2} \right)^2 + \left(\sum_{j=1}^{n+1} \frac{-y_u + u_j(x^2 + y_u^2)}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2} \right)^2}{\sum_{j=1}^{n+1} \frac{x}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2}}$$

Differential equations

Compute

$$\frac{d}{dx} \log \tilde{F}_u(x) = - \frac{\left(\sum_{j=1}^{n+1} \frac{x}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2} \right)^2 + \left(\sum_{j=1}^{n+1} \frac{-y_u + u_j(x^2 + y_u^2)}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2} \right)^2}{\sum_{j=1}^{n+1} \frac{x}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2}}$$

Substituting $u = (1)$ yields

$$\frac{d}{dx} \log \tilde{F}_{(1)}(x) = - \frac{x^2 + y_{(1)}^2}{x}.$$

Differential equations: Two curious inequalities

Use Cauchy-Schwarz:

$$\begin{aligned} \left(\sum_{j=1}^{n+1} \frac{x}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2} \right)^2 &= \left(\sum_{j=1}^{n+1} \frac{(x/u_j) \cdot u_j}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2} \right)^2 \\ &\leq \left(\sum_{j=1}^{n+1} \frac{(x/u_j)^2}{\left(x^2 + \left(\frac{1}{u_j} - y_u\right)^2\right)^2} \right) \left(\sum_{j=1}^{n+1} u_j^2 \right) \\ &= \sum_{j=1}^{n+1} \frac{(x/u_j)^2}{\left(x^2 + \left(\frac{1}{u_j} - y_u\right)^2\right)^2} \end{aligned}$$

Differential equations: Two curious inequalities

Use Cauchy-Schwarz again:

$$\begin{aligned} \left(\sum_{j=1}^{n+1} \frac{-y_u + u_j(x^2 + y_u^2)}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2} \right)^2 &= \left(\sum_{j=1}^{n+1} \frac{(-y_u/u_j + x^2 + y_u^2) \cdot u_j}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2} \right)^2 \\ &\leq \left(\sum_{j=1}^{n+1} \frac{(-y_u/u_j + x^2 + y_u^2)^2}{\left(x^2 + \left(\frac{1}{u_j} - y_u\right)^2\right)^2} \right) \left(\sum_{j=1}^{n+1} u_j^2 \right) \\ &= \sum_{j=1}^{n+1} \frac{(-y_u/u_j + x^2 + y_u^2)^2}{\left(x^2 + \left(\frac{1}{u_j} - y_u\right)^2\right)^2} \end{aligned}$$

Differential equations

Putting it together:

$$\begin{aligned}\frac{d}{dx} \log \tilde{F}_u(x) &\geq -\frac{\sum_{j=1}^{n+1} \frac{x^2 + y_u^2}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2}}{\sum_{j=1}^{n+1} \frac{x}{x^2 + \left(\frac{1}{u_j} - y_u\right)^2}} \\ &= -\frac{x^2 + y_u^2}{x} \\ &\stackrel{(*)}{\geq} -\frac{x^2 + y_{(1)}^2}{x} \\ &= \frac{d}{dx} \log \tilde{F}_{(1)}(x)\end{aligned}$$

which is sufficient to imply $\tilde{F}_u(x) \geq \tilde{F}_{(1)}(x)$, as desired.

Thanks