MATH STUDIES HOMEWORK 5

JAMES CUMMINGS

Submit the LATEX file containing your solutions to the email address algebrahw@legba.math.cmu.edu by 1159 pm on Friday 25 March.

A point of clarification: when we say that an operator K in B(V) is invertible we mean that there is L in B(V) such that KL = LK = Iwhere as usual I is the identity.

As we see in class soon, if a map in B(V) is bijective then its inverse is bounded. So a non-invertible K in B(V) fails to be bijective, hence either it's not into or not onto. If V is FD then not-into equals not-onto but they are distinct in l_2 ; consider the maps taking $(x_0, x_1, x_2...)$ to $(0, x_0, x_1, ...)$ and $(x_1, x_2, ldots)$.

- (1) Let H be the subset of $l_2(\mathbb{C})$ consisting of those sequences with $|x_n| \leq 1/(n+1)$. Show that H is compact. Hint: sequential compactness.
- (2) Let K be the bounded operator on $l_2(\mathbb{C})$ which maps (x_n) to $(x_n/(n+1))$. Show that K is compact. (Trickier) For which sequences of complex numbers (α_n) is the map taking (x_n) to $(\alpha_n x_n)$ compact?
- (3) Let K be as in the previous question. Find the eigenvalues of K and show that 0 is in the spectrum but not an eigenvalue. Is 0 the only point in the spectrum which is not an eigenvalue?
- (4) Let $V = C^n$ and let T be a linear operator on V. Show that TT^{\dagger} is self-adjoint. Are all self-adjoint operators of this form?
- (5) Let V and W be complex Hilbert spaces. Show how to give $V \oplus W$ a Hilbert space structure so that $V \oplus \{0\}$ is a closed subspace IMic to V and $\{0\} \oplus W$ is a closed subspace IMIc to W. Can you find a reasonable Hilbert space structure for $V \otimes W$?