

Review Problems for Test 2

1. Consider the function f defined by $f(x, y, z) = xy + y^2z$.
 - a) Find $\vec{\nabla} f$.
 - b) Find the directional derivative of f at the point $(2, 3, -1)$ in the direction of the vector $\vec{v} = \langle 2, 4, -3 \rangle$.
2. Suppose that f is a smooth function with the following property: At the point $(0, 0, 0)$, f decreases most rapidly in the direction of $\langle 6, -1, 2 \rangle$ and the rate of change of f in this direction is -5 . Find $\vec{\nabla} f(0, 0, 0)$.
3. Find the tangent plane and normal line to the surface $z = x^2 + 4y^2$ at the point $P_0 = (-1, 2, 17)$.
4. Let $\vec{u} = \frac{1}{\sqrt{2}} \langle 1, 1 \rangle$ and $\vec{v} = \frac{1}{\sqrt{2}} \langle 1, -1 \rangle$.
Assume that f_x and f_y are continuous. If $D_{\vec{u}}f(2, 1) = 2\sqrt{2}$ and $D_{\vec{v}}f(2, 1) = -8\sqrt{2}$, find $\vec{\nabla} f(2, 1)$.
5. Find all points on the surface $x^2 + 4y^2 - z^2 = 4$ at which the tangent plane is parallel to the plane $x + 2y + z = 6$.
6. Suppose that $w = x^3y + y^2 - 4z^3$, $x = rs$, $y = (r^2 - s^2)^3$, and $z = 2r + s$.
Use the chain rule to compute $\frac{\partial w}{\partial s}$. (Express your answer in terms of r and s alone. Algebraic simplification is not necessary.)
7. Suppose that $w = f\left(\frac{y}{y-x}\right)$ where f is continuously differentiable.
Show that
$$x \frac{\partial w}{\partial x} + y \frac{\partial w}{\partial y} = 0, \quad x \neq y.$$

8. Assume that u and v have continuous second order partial derivatives and that

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = \frac{-\partial v}{\partial x}.$$

Show that

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \text{and} \quad \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = 0.$$

9. Assume that $w = f(u)$, where $u = x^2 + y^2$ and f is a smooth function. Find an expression for $\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2}$ in terms of u , $\frac{dw}{du}$, and $\frac{d^2w}{du^2}$.
10. Find all local maxima, local minima, and saddle points for the function f defined by

a) $f(x, y) = x^2 + xy + y^2 + x - y + 2,$

b) $f(x, y) = 3x^2 - 3xy^2 + y^3 + 3y^2.$

11. Let R be region described by

$$0 \leq y \leq \sqrt{16 - x^2}, \quad 0 \leq x \leq 4$$

Find the absolute maximum and the absolute minimum values of

$$f(x, y) = x^2 + y^2 - 4x - 2y + 1$$

on R .

12. Use Lagrange multipliers to find all possible maxima and minima of f subject to the given constraint.

a) $f(x, y, z) = \frac{2}{3}x^3 + y - 3z;$
 $x^2 + y^2 + z^2 = 10$

b) $f(x, y) = xy; \quad 9x^2 + y^2 = 4$

13. Write an equivalent double integral with the order of integration reversed.

a) $\int_0^1 \int_y^{\sqrt{y}} f(x, y) dx dy$

b) $\int_0^2 \int_1^{e^{2x}} f(x, y) dy dx$

c) $\int_0^2 \int_{2x}^4 f(x, y) dy dx$

14. Evaluate each of the following.

a) $\int_0^1 \int_0^{x^3} 4x^2 y dy dx$

b) $\int_0^1 \int_x^1 3x\sqrt{x^2 + y^2} dy dx$