

MEASURABLE REALIZATIONS OF ABSTRACT SYSTEMS OF CONGRUENCES

CLINTON T. CONLEY, ANDREW S. MARKS, AND SPENCER T. UNGER

ABSTRACT. An abstract system of congruences describes a way of partitioning a space into finitely many pieces satisfying certain congruence relations. Examples of abstract systems of congruences include paradoxical decompositions and n -divisibility of actions. We consider the general question of when there are realizations of abstract systems of congruences satisfying various measurability constraints. We completely characterize which abstract systems of congruences can be realized by nonmeager Baire measurable pieces of the sphere under the action of rotations on the 2-sphere. This answers a question of Wagon. We also construct Borel realizations of abstract systems of congruences for the action of $\mathrm{PSL}_2(\mathbb{Z})$ on $\mathbb{P}^1(\mathbb{R})$. The combinatorial underpinnings of our proof are certain types of decomposition of Borel graphs into paths. We also use these decompositions to obtain some results about measurable unfriendly colorings.

1. INTRODUCTION

Recently, several results have been proved about the extent to which realizations of geometrical paradoxes can be found with sets having measurability properties such as being Borel, Lebesgue measurable, or Baire measurable (see for instance [DF][GMP16][GMP17][MU16][MU17]). This is a growing area of study at the interface of descriptive set theory, combinatorics, and ergodic theory. This paper is a contribution to this study. One of the earliest results in this vein is the theorem of Dougherty and Foreman [DF] that the Banach-Tarski paradox can be realized using Baire measurable pieces. In contrast to the classical Banach-Tarski paradox which uses five pieces, Dougherty and Foreman's Baire measurable solution uses six pieces. A result of Wehrung [Weh] implies that this is optimal; there is no Baire measurable realization of the Banach-Tarski paradox with five pieces. This suggests a subtle difference between the classical and Baire measurable contexts.

In this paper, we consider a refined framework—abstract systems of congruence—for describing when an action can be partitioned into finitely many pieces satisfying certain congruence relations. As one application, we give an exact characterization of which abstract systems of congruences can be realized in the 2-sphere with arbitrary pieces versus nonmeager Baire measurable pieces. This refines the dual results of Wehrung [Weh], and Dougherty and Foreman [DF].

We formally define abstract systems of congruences as follows. Given a set S , its **proper powerset** $\mathcal{P}_{\mathrm{pr}}(S)$ is $\mathcal{P}_{\mathrm{pr}}(S) = \{R \subseteq S : R \neq \emptyset \wedge R \neq S\}$. Following Wagon [W, Definition 4.10], an **abstract system of congruences** on $n = \{0, \dots, n-1\}$

Date: March 11, 2019.

The first author was partially supported by DMS-1500906. The second author was partially supported by NSF grant DMS-1500974. The third author was partially supported by NSF grant DMS-1700425.

is an equivalence relation E on $\mathcal{P}_{\text{pr}}(n)$, so that if $U E V$, then $U^c E V^c$. Here U^c denotes the complement of U . Suppose $a: \Gamma \curvearrowright X$ is an action of a group on a set X . Then we say that $A, B \subseteq X$ are a -congruent if there is a group element $\gamma \in \Gamma$ such that $\gamma \cdot A = B$. An a -**realization** of an abstract system E of congruences is a partition $\{A_0, \dots, A_{n-1}\}$ of X such that for all $U, V \in \mathcal{P}_{\text{pr}}(n)$ with $U E V$, we have that $\bigcup_{i \in U} A_i$ and $\bigcup_{i \in V} A_i$ are a -congruent. The definition of an abstract system of congruences reflects that fact that congruence is an equivalence relation, and that if $A, B \subseteq X$ are congruent, then A^c and B^c are also congruent.

An important example of an abstract system of congruences is the smallest abstract system of congruences E on $\mathcal{P}_{\text{pr}}(4)$ containing the relations $\{0\} E \{0, 1, 2\}$ and $\{1\} E \{0, 1, 3\}$. A realization of this system gives a paradoxical decomposition, since $\{0, 3\}$ and $\{1, 2\}$ partition $\{0, 1, 2, 3\}$. The translation action of the free group on two generators \mathbb{F}_2 on itself is an example of an action realizing this system of congruences [W, Theorem 4.2]. Another important example of an abstract system of congruences is the smallest abstract system of congruences E on $\mathcal{P}_{\text{pr}}(n)$ where $\{i\} E \{j\}$ for every $i, j \in n$. An action is said to be n -divisible if it satisfies this system of congruences (that it, is can be partitioned into n congruent pieces). For example, it is easy to see that the action of the rotation group SO_3 on the 2-sphere is not 2-divisible by considering the ‘‘poles’’ of the rotation. However, this action is n -divisible for $n \geq 3$ (see [W, Corollary 4.14]).

Wagon has characterized which abstract systems of congruences can be realized in the action of the group SO_3 of rotations on the 2-sphere. We say that an abstract system of congruences E on n is **non-complementing** if there is no set $X \in \mathcal{P}_{\text{pr}}(n)$ such that $X E X^c$.

Theorem 1.1 ([W, Corollary 4.12]). *Suppose E is an abstract system of congruences. E can be realized in the action of SO_3 on the 2-sphere if and only if E is non-complementing.*

We show that in order to realize an abstract system of congruences with Baire measurable pieces in the sphere, we need one additional property. Say that an abstract system of congruences E on $\mathcal{P}_{\text{pr}}(n)$ is **non-expanding** if there do not exist sequences of sets $(V_i)_{i \leq k}$ and $(W_i)_{i \leq k}$ where $V_i E W_i$ for every $i \leq k$ and $W_i \subseteq V_{i+1}$ for every $i < k$, such that $W_k \subsetneq V_0$. Hence,

$$V_0 E W_0 \subseteq V_1 E W_1 \subseteq \dots V_k E W_k \subsetneq V_0.$$

Theorem 1.2. *Suppose E is an abstract system of congruences. Then E can be realized in action of SO_3 on the 2-sphere with Baire measurable pieces each of which is nonmeager if and only if E is non-complementing and non-expanding.*

This theorem positively answers Wagon’s question [W, Page 47] of whether the 2-sphere is n -divisible with Baire measurable pieces for $n \geq 3$. Wagon has also asked whether the 2-sphere is n -divisible into Lebesgue measurable pieces. This remains an open problem.

Let \mathbb{F}_n be the free group on n generators. Our proof of Theorem 1.2 shows more generally that if $n \geq 2$, then any free Borel action of \mathbb{F}_n on a Polish space X can realize an abstract system of congruences that is non-expanding and non-complementing using Baire measurable pieces. (See Lemma 3.6).

Our main tool for proving Theorem 1.2 is a decomposition lemma for acyclic Borel graphs into sets of paths with a property concerning how the paths from different sets may overlap.

Definition 1.3. Suppose G is a graph and G_0, G_1, \dots is a sequence of subgraphs of G . Then we say G_0, G_1, \dots is **end-ordered** if for all vertices x in G , if x is a vertex in G_i and G_j where $i < j$, then x is a leaf in G_j . Similarly, if S_0, S_1, \dots are sets of subgraphs of G , then we say that S_0, S_1, \dots is **end-ordered** if for all vertices x in G , if x is vertex in $H \in S_i$ and a vertex in $H' \in S_j$ where $i < j$, then x is a leaf in H' .

Definition 1.4. Suppose G is an acyclic Borel graph. Then a path decomposition of G is a sequence P_0, P_1, \dots of sets of paths in G such that P_0, P_1, \dots is end ordered, every P_i consists of vertex disjoint paths, and for every edge e in G , there exists exactly one P_i so that e appears in a path in P_i . We say that a path decomposition is Borel if each set P_i is Borel, and the path decomposition has length at least n if every path has length at least n .

Roughly, a path decomposition is a way of covering the graph with sets of paths P_0, P_1, \dots so that all the paths in P_j have interiors that are disjoint from the paths in P_i , for $i < j$.

One of our main lemmas (Lemma 3.4) says that if G is a locally finite acyclic Borel graph, then for all n , there is a comeager set on which G has a Borel path decomposition of length at least n .

A different case in which we have Borel path decompositions is when we have Borel end selections. Recall that if G is a graph on X , a **ray** is an infinite simple path in G , and that two rays $(x_i)_{i \in \mathbb{N}}$ and $(y_i)_{i \in \mathbb{N}}$ are **end-equivalent** if for every finite set $S \subseteq X$, the rays (x_i) and (y_i) eventually lie in the same connected component of $G \upharpoonright (X \setminus S)$. An **end** of G is an end-equivalence class of G . If G is a Borel graph on X , we say that G **has a Borel selection of finitely many ends in each connected component** if and only if there is a Borel function sending each $x \in X$ to a ray $r(x)$ in the connected component of x so that the image of every connected component of G under r contains representatives of only finitely many end equivalence classes.

We show that if G is an acyclic bounded degree Borel graph on X such that there is a Borel selection of finitely many ends in every connected component of G , then for every n we can find a Borel path decompositions of G of length at least n (see Lemma 4.2). We construct explicit realizations of abstract systems of congruences for the action of $\mathrm{PSL}_2(\mathbb{Z})$ on $P^1(\mathbb{R})$, by combining this lemma with an explicit end selection defined using continued fraction expansions.

Theorem 1.5. *Suppose E is an abstract system of congruences which is non-complementing and non-expanding. Then E can be realized in the action of $\mathrm{PSL}_2(\mathbb{Z})$ on $P^1(\mathbb{R})$ by Borel pieces.*

For example, this action is n -divisible using Borel pieces for $n \geq 3$.

Open Problem 1.6. *Characterize the abstract systems of congruences which can be realized in the action of $\mathrm{PSL}_2(\mathbb{Z})$ on $P^1(\mathbb{R})$ by Borel pieces.*

It is a theorem of Adams [JKL, Lemma 3.21] that if G is a locally finite graph on a standard probability space (X, μ) and G is μ -hyperfinite, then G admits a μ -measurable selection of finitely many ends. Using Adams's theorem, we also show that any μ -hyperfinite action of \mathbb{F}_2 on a standard probability space (X, μ) has a μ -measurable realization of any abstract system of congruences E , if E is non-complementing and non-expanding. (See Theorem 4.3).

Our decomposition lemmas have some other applications in Borel combinatorics. Simon Thomas has asked whether every locally finite Borel graph has an unfriendly Borel coloring, where an **unfriendly coloring** of a graph G on X is a function $f: X \rightarrow 2$ such that for every x ,

$$|\{y \in N(x): c(x) \neq c(y)\}| \geq |\{y \in N(x): c(x) = c(y)\}|.$$

Thomas's question is partially motivated by the open problem in classical combinatorics of whether every countable graph admits an unfriendly coloring. If G is a graph on X , say that a function $f: X \rightarrow 2$ is **strongly unfriendly** if for every x , $|\{y \in N(x): c(x) = c(y)\}| \leq 1$.

We use our decomposition lemma to prove the following result:

Theorem 1.7. *Suppose G is a locally finite acyclic Borel graph on a Polish space X that admits a Borel path decomposition of length at least 5. Then G has a Borel strongly unfriendly coloring. Hence, if G is a locally finite acyclic Borel graphs of degree at least 2, then G admits a Baire measurable strongly unfriendly colorings, and G admits a μ -measurable strongly unfriendly colorings for every Borel probability measure on X rendering G μ -hyperfinite.*

In Section 6 we also discuss some further applications of our decompositions, such as new proofs of Baire measurable and μ -measurable edge-coloring and matchings.

Acknowledgments. The authors would like to thank Alekos Kechris and Anton Bernshteyn for helpful discussions about the material of the paper.

2. PRELIMINARIES

Our notation for graph theory is standard, see [D]. We recall a few notions. By a graph on X we mean a simple undirected graph with vertex set X . The **degree** of a vertex is its number of neighbors. Two vertices are **adjacent** if there is an edge between them. A vertex is a **leaf** if it has degree 1, and is a **splitting vertex** if it has degree at least 3. By a **path** we mean a simple path of finite length x_0, \dots, x_n . The endpoints of the path are x_0 and x_n , and the remaining vertices are interior vertices of the path. By a **ray**, we mean a simple infinite path $(x_i)_{i \in \mathbb{N}}$.

If G is a graph, we say a set of vertices is **independent** if it does not contain two adjacent vertices. We say that a set A is **k -independent** if for all distinct $x, y \in A$, we have $d(x, y) > k$.

Suppose G is a graph on X . A **subgraph** H of G is a graph on a subset of X so that every edge in H is an edge in G . If $Y \subseteq X$, the **restriction of G to Y** or **induced subgraph on Y** , denoted $G \upharpoonright Y$, is the graph on Y where the edges of $G \upharpoonright Y$ are all edges in G with vertices in Y .

A Borel graph is a graph on a Polish space X whose edge relation is Borel. For background on Borel graphs see [KM]. An important example of a Borel graph arises from Borel group actions. If a is a Borel action of a countable group Γ on a Polish space X and $S \subseteq \Gamma$ is a symmetric set of group elements, then we let $G(a, S)$ be the graph on X where x, y are adjacent if there exists some $\gamma \in S$ such that $\gamma \cdot x = y$.

If G is a Borel graph on X , the set of all paths of G is a Borel subset of $\bigcup_n X^n$, and hence a standard Borel space. Hence we may speak about a set of paths in G being Borel.

We note that in contrast to Lemmas 3.4 and 4.2 there exist Borel graphs which do not admit Borel path decompositions of length at least 3.

Theorem 2.1. *Suppose that G is Borel graph of degree at least 3 on a Polish space X that admits an invariant measure μ . Then G does not admit a Borel path decomposition on any μ -conull Borel set.*

Proof. Let P_0, P_1, \dots be a Borel path decomposition. Note that every vertex x must be the endpoint vertex of some unique path $p(x) \in P_i$ since G has degree at least 3. Define a Borel function $f: X \rightarrow X$ where $f(x)$ is the vertex adjacent to x in $p(x)$. Then f is a compression function contradicting μ being measure preserving. [N] \square

We will use the following lemma giving a criterion for the existence of abstract systems of congruences without any measurability properties.

Definition 2.2. Suppose E is an abstract system of congruences. Say that a relation R on a set X **generates** the equivalence relation E on X if the smallest abstract system of congruences containing R is equal to E . Say that a generating set R for E is **good** if R contains all pairs $(U, V) \in E$ such that $U = V^c$. Finally, a minimal good generating set of E is a good generating set R so that there is no proper subset of R that is a good generating set for E .

Lemma 2.3 (See also [W, Section 4]). *Suppose that E is an abstract system of congruences on n , and $R = \{(S_1, T_1), \dots, (S_k, T_k)\}$ is a minimal good generating set of E . Suppose $a: \Gamma \curvearrowright X$ is an action of*

$$\Gamma = \langle \gamma_1 \dots \gamma_k \mid \{\gamma_i^2 = 1: T_i = S_i^c\} \rangle.$$

Suppose finally that for every $x \in X$, $\text{Stab}(\{x\})$ is abelian. Then there is an a -realization $\{A_0, \dots, A_{n-1}\}$ of E , witnessed by

$$\gamma_i \cdot \bigcup_{j \in S_i} A_j = \bigcup_{j \in T_i} A_j.$$

Proof. This Lemma is proved in [W, Section 4] when E is non-complementing.

Using the axiom of choice, it suffices to prove the lemma when the action has the single orbit. By the Kurosh subgroup theorem, it is clear that the stabilizer of every point must be cyclic. Hence, the graph $G(a, \{\gamma_i: i \leq k\})$ is acyclic. From this it is easy to construct a realization of the desired abstract system of congruence. \square

Throughout we will be working with actions of such groups Γ that are free products of copies of \mathbb{Z} and $\mathbb{Z}/2\mathbb{Z}$, and where the generators of Γ of order 2 will witness congruences of the form $U E U^c$.

3. BAIRE MEASURABLE REALIZATIONS

In this section we prove Theorem 1.2. We begin with a decomposition lemma for acyclic locally finite Borel graphs (Lemma 3.4). As an intermediate step towards this lemma, we consider decompositions into certain types of trees that themselves have suitable decompositions into paths. Recall that a **tree** is a connected acyclic graph, a **leaf** of a tree is a vertex of degree 1, and a **splitting vertex** is a vertex of degree at least 3. Say that a tree T is **n -spindly** if there is at most one leaf l of T so that for all distinct leaves x, y , if $l \notin \{x, y\}$, then $d(x, y) > 2n$, and if $l \in \{x, y\}$, then $d(x, y) \geq n$. The utility of spindly trees is the following lemma:

Lemma 3.1. *Every finite n -spindly tree T can be written as a union of edge-disjoint paths p_0, p_1, \dots each having length at least n , and which are end-ordered.*

Proof. We construct p_0, p_1, \dots, p_k by induction. Let p_0 be a path from one leaf to another leaf, having minimal length among such paths between leaves. Let the endpoints of p_0 be x and y . We may assume $x \neq l$ for the distinguished leaf l if it exists.

For each vertex z not in p_0 , let V_z be the set of w such that there is a path p from z to w such that no interior vertex of p is in p_0 . Since T is a tree, there is exactly one vertex in $T \upharpoonright V_z$ which is contained in p_0 . Let this vertex be l_z , which is a leaf in $T \upharpoonright V_z$. For any leaf w in $T \upharpoonright V_z$, the distance $d(x, l_z) \leq d(w, l_z)$. Otherwise if $d(w, l_z) < d(x, l_z)$, the path from w to y would have smaller length than p_0 , but p_0 has minimal length. Hence, $d(w, l_z) \geq n$ since otherwise $d(x, l_z) \leq d(w, l_z) < n$ which implies that $d(x, w) < 2n$ contradicting T being n -spindly, since neither x nor w are equal to l . It follows that $T \upharpoonright V_z$ is n -spindly witnessed by l_z .

The lemma follows by inductively applying the lemma to all these n -spindly subgraphs of the form $T \upharpoonright V_y$. \square

Remark 3.2. *Every locally finite n -spindly tree T can be written as a union of edge-disjoint paths p_0, p_1, \dots that are of length at least n and which are end-ordered. That is, Lemma 3.1 remains true for infinite n -spindly trees. This is by an infinite iteration of the same process in the proof of Lemma 3.1 (or by a compactness argument).*

As an intermediate step towards our path decomposition, we prove a lemma decomposing into n -spindly trees.

Lemma 3.3. *Suppose G is a locally finite acyclic graph on a Polish space X , and $n \geq 1$. Then there are a G -invariant comeager Borel set D and edge-disjoint Borel subgraphs G_0, G_1, \dots such that $\bigcup_i G_i = G \upharpoonright D$, every connected component of G_i is a finite n -spindly tree, and the G_0, \dots is end-ordered.*

Proof. We give a construction in countably many steps. Let $d(i) = 3n6^i$. By [MU16, Lemma 3.1], let $(A_i)_{i \in \mathbb{N}}$ be subsets of X such that the elements of A_i are pairwise of distance greater than $3d(i)$, and $D = \bigcup_i A_i$ is comeager and G -invariant. Before step s we will have constructed edge-disjoint Borel subgraphs G_0, G_1, \dots, G_{s-1} . Let $H_i = \bigcup_{j \leq i} G_j$. At step i , if C is a connected component of H_i , then we call C a **k -component in H_i** if k is least such that C is also a connected component of H_k . Our induction hypotheses are as follows:

- (1) For every $i < s$ and $x \in A_i$, there is an edge incident to x in H_i .
- (2) For every $k \leq s-1$, the diameter of any k -component in H_{s-1} is at most $d(k)$.
- (3) For every $k \leq s-1$, the distance between any two k -components in H_{s-1} is at least $2d(k)$.
- (4) The distance between any two connected components of H_{s-1} is greater than $2n$.

Note that these hypotheses imply that every edge in $G \upharpoonright D$ will appear in some G_i . To see this suppose x, x' are adjacent where $x \in A_s$ and $x' \in A_{s'}$. Then x and x' must both be in connected components of $H_{\max(s, s')}$ by (1), and hence the same connected component by (4). Thus the edge (x, x') must be in $H_{\max(s, s')}$ since G is acyclic.

Below we inductively define G_s . Note that to satisfy part (1) of the induction hypothesis, we need to add an edge incident to each $x \in A_s$ to G_s if there is not one already one in H_{s-1} . However, simply adding such edges by themselves may violate induction hypothesis (4). So we will need to inductively define G_s to include paths to all nearby k -components for $k < s$ so they all become the same connected component in H_s . Hypotheses (2) and (3) give us control over this process so we can satisfy (4).

To begin, let $G_{s,0}$ be the graph consisting of all vertices in A_s (and no edges). Inductively, for $0 < i \leq s$, let $G_{s,i} \supseteq G_{s,i-1}$ be the union of $G_{s,i-1}$ with all paths of length at most $d(s-i)$ from vertices in $G_{s,i-1}$ to $(s-i)$ -components in H_{s-1} . Since elements of A_s have pairwise distance at least $3d(s)$ it is clear by induction that components of $G_{s,i}$ have diameter at most $2d(s-1) + \dots + 2d(s-i)$, and hence components of $G_{s,s}$ have diameter at most $2d(s-1) + \dots + 2d(0) \leq d(s) - 2d(s-1)$. Similarly, the components of $G_{s,s}$ are pairwise of distance at least $3d(s) - 2d(s-1) - \dots - 2d(0) \geq 2d(s) + 2d(s-1)$. Each connected component of $G_{s,s}$ is n -spindly by construction, since any $x \in A_s$ can be of distance less than n from at most one connected component of H_{s-1} by (4).

Let A'_s be the set of $x \in A_s$ that are not incident to any edge of H_{s-1} or $G_{s,s}$. (Hence, every vertex of A'_s is at least distance $d(0) > 3n$ from any i -component in H_{s-1} for $i < s$). For each $x \in A'_s$, let $p(x)$ be the lex-least path of length n in G starting at x . Let $J_s = \{p(x) : x \in A'_s\}$. Let $G_s = G_{s,s} \cup J_s$. Hence H_s satisfies (1). By the above paragraph, it is clear that connected components of G_s are pairwise of distance at least $2d(s) - 2d(s-1)$ and are of diameter at most $d(s) - 2d(s-1)$. Hence, since each connected component of G_s is connected, and connected components of H_{s-1} have diameter at most $d(s-1)$ by (2), we see that s -components in H_s are of diameter at most $d(s)$ and pairwise of distance at least $d(s-1)$. Hence H_s satisfies (2) and (3).

Now we show that part (4) of the induction hypothesis holds. Suppose C is an s -component of H_s . We want to show that distance from C to any other k -component C' is greater than $2n$. Let x be the closest vertex in C to C' . We may assume that $x \in G_{s,s}$. Let x' be the closest element of $G_{s,s-k} \upharpoonright C$ to C' . Hence $d(x, x') \leq d(k-1) + \dots + d(0)$. Since C' is distance at most $2n$ from x , C' is distance at most $2n + d(k-1) + \dots + d(0) < d(k)$ from x' . First suppose x' is also a vertex in $G_{s,s-k-1}$. Then x' would be an element of $G_{s,s-k-1}$ of distance $< d(k)$ from a k -component, and so in the definition of $G_{s,s-k}$ there should have been a path added from $G_{s,s-k-1}$ to C' in $G_{s,s-k}$. If x' is not a vertex in $G_{s,s-k-1}$, then x' must be part of a path added in $G_{s,s-k}$ from an element of $G_{s,s-k-1}$ to some k -component C'' . Since this path is of distance at most $d(k)$, this would imply that C' and C'' are of distance $< 2d(k)$ which contradicts part (3) of the induction hypothesis. \square

We are now ready to prove our path decomposition lemma.

Lemma 3.4. *Suppose G is a locally finite acyclic graph on a Polish space and $n \geq 1$. Then there is a comeager Borel set D such that $G \upharpoonright D$ has a path decomposition of length at least n .*

Proof. We prove this lemma by combining Lemma 3.3 and Lemma 3.1 with the obvious derivative process to obtain sets of paths.

Suppose $(D_i)_{i \in \mathbb{N}}$ is such that each $S \in D_i$ is a finite sequence of paths in G that are end-ordered, and $(D_i)_{i \in \mathbb{N}}$ is end-ordered. Let $<_{(D_i)}$ (suppressing the indexing

for clarity) be the partial order on the paths appearing in the elements of the D_i where $p <_{(D_i)} p'$ if p, p' share some vertex, and either p, p' both appear in some sequence $S \in D_i$ where p appears before p' , or p is in an element of D_i and p' is in an element of D_j for $i < j$.

We begin by applying Lemma 3.3 to obtain a n -spindly decomposition G_0, G_1, \dots of G restricted to some comeager G -invariant Borel set. If C is an n -spindly connected component of some G_i , let $P(C)$ be the lexicographically least decomposition (p_0, \dots, p_k) of C satisfying the conclusion of Lemma 3.1. Letting $D_{i,0} = \{P(C) : C \text{ is a connected component of } G_i\}$, we obtain a sequence $(D_{i,0})_{i \in \mathbb{N}}$ of sets of finite sequences of paths in G , and the associated partial order $<_{(D_{i,0})}$ defined in the previous paragraph.

Inductively, for $j \geq 0$, let P_j be the set of p appearing in some element of $D_{i,j}$ such that there is no $p' <_{(D_{i,j})} p$. Then let $D_{i,j+1}$ be the set of all sequence in $D_{i,j}$ with all elements of P_j removed. These P_j are our desired set of paths. Every path p' in each $S \in D_i$ must eventually appear in some P_j since there are only finitely many p such that $p <_{(D_{i,0})} p'$. \square

A useful observation is that if G is a graph with a path decomposition, the decomposition may be assumed to consist of paths of bounded length. This follows the fact that the intersection graph on paths has a countable Borel coloring, and a derivative operation analogous to that of Lemma 3.4.

Lemma 3.5. *Suppose G is a locally finite Borel graph with a path decomposition P_0, P_1, \dots of length at least n . Then G admits a Borel path decomposition P'_0, P'_1, \dots of length at least n such that every path $p \in P'_i$ has length at most $2n$.*

Proof. Every path of length greater than $2n$ can clearly be written as a finite union of paths of length between n and $2n$. Hence, we may replace any path $p \in P_i$ of length greater than $2n$ by the lex-least finite set of paths of length between n and $2n$ whose union is p . This gives a sequence P_0, P_1, \dots having every property of being a Borel path decomposition with the exception that the P_i may not consist of vertex disjoint paths (but with the property that every path in every P_i has length at most $2n$).

Let H be the graph on the paths $\bigcup_i P_i$ where distinct $p, p' \in \bigcup_i P_i$ are adjacent in H if they share some vertex. Then H is a locally finite Borel graph and hence has a countable Borel coloring $c: \bigcup_i P_i \rightarrow \mathbb{N}$ by [KST, Proposition 4.10].

Inductively, let $D_{i,0} = P_i$. For a fixed j , we can order the paths in $\bigcup_i D_{i,j}$ by $p <_{(D_{i,j})} p'$ if $p \in D_{i,j}$ and $p' \in D_{i',j}$ where either $i < i'$, or $i = i'$ and $c(p) < c(p')$. Now a construction identical to the last paragraph of the proof of Lemma 3.4 gives our desired Borel path decomposition. \square

Lemma 3.6. *Suppose that E is an abstract system of congruences on n which is non-expanding, and $R = \{(S_1, T_1), \dots, (S_k, T_k)\}$ is a minimal good generating set of E . Suppose also a is a free Borel action of the group*

$$\Gamma = \langle \gamma_1 \dots \gamma_k \mid \{\gamma_i^2 = 1 : T_i = S_i^c\} \rangle$$

on a Polish space X . If $G(a, \{\gamma_1, \dots, \gamma_k\})$ has a Borel path decomposition of length at least r for sufficiently large r (depending on E), then there is an a -realization of E with Borel pieces witnessed by

$$(*) \quad \gamma_i \cdot \bigcup_{j \in S_i} A_j = \bigcup_{j \in T_i} A_j.$$

Proof. Let G be the graph $G = G(a, \{\gamma_1, \dots, \gamma_k\})$. The idea of our proof is as follows. We first argue that there is a sufficiently large length r so that given any path p of length at least r in G , if we have already assigned the endpoints of p to be in elements of A_0, \dots, A_{n-1} , then there is some way of consistently assigning the interior points of the path to elements of A_0, \dots, A_{n-1} so as to obey the congruences required in (*). Then we use a path decomposition of length at least r for G to inductively construct a realization of this system of congruences.

Suppose that $g = g_l \dots g_0$ is a reduced word in Γ , where $g_i \in \{\gamma_1^\pm, \dots, \gamma_k^\pm\}$ are generators. If we begin at some $x \in X$, then such a reduced word of length $l + 1$ gives a path of length $l + 1$ in G : the path $x, g_0 \cdot x, \dots, g_l \dots g_0 \cdot x$. We give a definition concerning what elements of A_0, \dots, A_{n-1} the elements of this path can belong to. Define functions X and Y on generators as follows: $X(\gamma_j) = S_j$, $Y(\gamma_j) = T_j$, $X(\gamma_j^{-1}) = T_j$, and $Y(\gamma_j^{-1}) = S_j$. Say that n_0, \dots, n_{l+1} is a **labeling of $g = g_l \dots g_0$** if for all i , we have $n_i \in X(g_i)$ if and only if $n_{i+1} \in Y(g_i)$. So labelings correspond to acceptable assignments of the points $x, g_0 \cdot x, \dots, g_l \dots g_0 \cdot x$ to the sets A_0, \dots, A_{n-1} .

We are interested in the ways labelings of g may start and end. If $k, m \in n$, say a reduced word g is (k, m) -**bad** if there is no labeling n_0, \dots, n_{l+1} of g with $n_0 = m$ and $n_{l+1} = k$. Say that g is **bad** if there is some $k, m \in n$ such that g is (k, m) -bad. We will use a pigeonhole principle argument to show there is a bound on the length of bad words.

To begin, note that if $g = g_l \dots g_0$ is bad, then $g_l \dots g_1$ and $g_{l-1} \dots g_0$ are also bad. That is, initial segments and final segments of bad words are bad.

Suppose $g = g_l \dots g_0$ is (k, m) -bad. It must be that either $m \in Y(g_l)$ in which case g is (k, m') -bad for every $m' \in Y(g_l)$, or $m \in Y(g_l)^c$ in which case g is (k, m') -bad for every $m' \in Y(g_l)^c$. Note that exactly one of the above holds since there is always a labeling n_0, \dots, n_{l+1} of g with $n_0 = m$ for any given m .

Fix a (k, m) -bad word g . Define a pair of associated sequences $V_{g,k}(i)$ and $W_{g,k}(i)$ where $(V_{g,k}(i), W_{g,k}(i)) = (X(g_i)^c, Y(g_i)^c)$ if $g_i \dots g_0$ is (k, m') -bad for every $m' \in Y(g_i)^c$ and $(V_{g,k}(i), W_{g,k}(i)) = (X(g_i), Y(g_i))$ otherwise. By identical reasoning to the previous paragraph, for every i , there are labelings n_0, \dots, n_{l+1} of g where $n_i \in V_{g,k}(i)$ and $n_{i+1} \in W_{g,k}(i)$ for every i . Indeed, we have that $V_{g,k}(i) E W_{g,k}(i)$ by definition, and $W_{g,k}(i) \subseteq V_{g,k}(i+1)$ for all $i \leq l$ or else g is not a bad word.

Suppose for a contradiction that there are infinitely many bad words. We break into two cases

Case 1: suppose that there are arbitrarily long (k, m) -bad words g such that $W_{g,k}(i) = V_{g,k}(i+1)$ for all $i < l$. Hence $V_{g,k}(0) E V_{g,k}(1) E \dots E V_{g,k}(l)$. By the pigeonhole principle, and since initial segments and final segments of bad words are bad, we can find some (k, m) -bad word g such that

$$(**) \quad g \text{ has length at least } 2 \text{ and } V_{g,k}(0) = W_{g,k}(l) \text{ or } V_{g,k}(0) = W_{g,k}(l)^c.$$

We claim that this implies that either the word g is not reduced, or the generating set of E is not a minimal good generating set.

First, we may assume that g has minimal length among bad words with property (**), and so no proper subword of g has property (**).

If $V_{g,k}(0) = W_{g,k}(l)$, then the minimal length of g among words with (**) implies that $g_0 \neq g_i^{\pm 1}$ for any $i > 0$. This implies that the generating set R is not a minimal good generating set; the fact that $V_{g,k}(0) E V_{g,m}(1)$ follows from

$V_{g,k}(1) E W_{g,k}(1) = V_{g,k}(2) E \dots E V_{g,k}(l) E W_{g,k}(l)$ and $W_{g,k}(l) = V_{g,k}(0)$. In particular, removing the pair (S_j, T_j) where $g_0 = \gamma_j$ would still generate E . Hence the generating set is not minimal.

In the case that $V_{g,k}(0) = W_{g,k}(l)^c$, we can also remove the pair (S_j, T_j) where $g_0 = \gamma_j$, since there must be a generator witnessing $V_{g,k}(0) E V_{g,k}(0)^c = W_{g,k}(l)$ by our definition of a good generating set (see Definition 2.2). In particular, a good generating set must contain every relation of the form (S, S^c) where $S E S^c$.

Case 2: suppose case 1 does not hold. Then by the pigeonhole principle, and since initial segments and final segments of bad words are bad, we can find some (k, m) -bad word g such that $V_{g,k}(0) = W_{g,k}(l+1)$, and $W_{g,k}(i) \subsetneq V_{g,k}(i+1)$ for some $i \leq l$. Then we can obtain a contradiction to the non-expansion of E by cyclically permuting the sequences to bring $V_{g,k}(i+1)$ to the 0 position and $W_{g,k}(i)$ to the l th position.

This finishes the proof that there are only finitely many bad words.

Now let r be sufficiently large so that there are no bad words of length r , and let P_0, P_1, \dots be a Borel path decomposition of G of length at least r . We may assume that this path decomposition satisfies the conclusion of Lemma 3.5. Now we inductively construct a Borel a -realization A_0, \dots, A_{n-1} of E in countably many steps. After step i we will have assigned each vertex appearing in the paths in P_j for $j \leq i$ to some A_0, \dots, A_{n-1} .

At step i we will consider the paths $p \in P_i$. For each such path p , we assign the vertices of p to be the lex-least assignment to A_0, \dots, A_{n-1} that is consistent with the requirement (*) in the statement of the lemma. There is guaranteed to be such an assignment since we will have assigned at most the start and end node of the path to A_0, \dots, A_{n-1} and since the path has length at least r , the group element corresponding to it is not bad.

At the end of this construction we will have assigned every element of X to some A_0, \dots, A_{n-1} . Since every edge in G appears in some path p , this ensures that the requirement (*) is satisfied at the end of the construction. \square

Remark 3.7. *In Lemma 3.6, if the space X is assumed to be perfect, then the sets A_1, \dots, A_k can be chosen so each is nonmeager. To see this, notice first that it suffices to have a path decomposition where the first set P_0 of paths has a set of endpoints D that is nonmeager. This is because we may partition D into k many nonmeager sets since X is perfect, and in our construction in Lemma 3.6, we may assign the elements of D to the partition witnessing the abstract system of congruence arbitrarily.*

So we need to show that we can construct a path decomposition where the set of endpoints of paths in P_0 is nonmeager. To see this, observe that in our proof of Lemma 3.3 given the subsets $(A_i)_{i \in \mathbb{N}}$ of X such that the elements of A_i are pairwise of distance greater than $d(i)$, note that all the elements of the set A_0 become endpoints of paths in P_0 in the final path decomposition. Hence, it suffices to show that A_0 can be chosen to be nonmeager in [MU16, Lemma 3.1]. To see this, note first that we can find a Borel nonmeager k -independent set. This is because $G^{\leq k}$ has a countable Borel coloring [KST, Proposition 4.10] and one of the color sets must therefore be a nonmeager k -independent Borel set A_0 . Now apply [MU16, Lemma 3.1] to the graph $G \setminus A_0$ and the function $f(n) = d(n+1)$.

We are now ready to prove Theorem 1.2.

Proof of Theorem 1.2. We begin with the forward direction of Theorem 1.2. Suppose A_0, \dots, A_{n-1} is a Baire measurable realization of an abstract system of congruence E on n where every A_i is nonmeager. By Theorem 1.1 it suffices to show that E is non-expanding. For a contradiction, suppose there are sequences of sets $(V_i)_{i \leq k}$ and $(W_i)_{i \leq k}$ with $V_i, W_i \in \mathcal{P}_{\text{pr}}(m)$ such that $V_i E W_i$ for every $i \leq k$, $W_i \subseteq V_{i+1}$ for every $i < k$ and $V_0 \supsetneq W_k$. Let $A = \bigcup_{i \in V_0} A_i$ and $B = \bigcup_{i \in W_k} A_i$. Let γ be the product of the group elements witnessing $V_i E W_i$ taken in increasing order for $i \leq k$. It follows that $\gamma \cdot A \subseteq B$. Clearly if $x \in A \setminus B$, then for all n , $\gamma^n \cdot x \notin B \setminus A$.

Now there are two cases. First, if the rotation given by γ is rational (i.e. periodic), this implies that $A \setminus B$ is not in any orbit of γ except possibly the poles of the rotation. This contradicts the fact that $A \setminus B$ is nonmeager.

Second, suppose the rotation of γ is aperiodic. Then $A \setminus B$ meets each orbit of γ in at most one point which contradicts $A \setminus B$ being nonmeager as follows. If $A \setminus B$ was nonmeager, there would be an open set U in which $A \setminus B$ is comeager. But since γ is an irrational rotation, we can find some $n > 0$ rendering γ^n arbitrarily close to the identity, and hence some n for which $\gamma^n U \cap U \neq \emptyset$. Since γ is a homeomorphism, this implies that both $A \setminus B$ and $\gamma^n \cdot (A \setminus B)$ are comeager in $\gamma^n U \cap U$. But then there is some x so that $x \in A \setminus B$ and $\gamma^n \cdot x \in A \setminus B$ which is a contradiction. This finishes the proof of the forward implications.

To prove the reverse implication, suppose that E is non-complementing and non-expanding. Choose some $R = \{(S_1 T_1), \dots, (S_k T_k)\}$ which minimally generates E , and let $\langle \gamma_1 \dots \gamma_k \rangle$ be rotations of the 2-sphere which generate a copy of \mathbb{F}_k .

Now let r be sufficiently large (so as to satisfy the hypothesis of Lemma 3.6). By Lemma 3.4, we can find a comeager G -invariant Borel set D so that there is a Borel path decomposition of length at least r of $G \upharpoonright D$. Let a' be the restriction of the action of $\langle \gamma_1, \dots, \gamma_k \rangle$ to D . Then by Lemma 3.4 we can find a Borel a' -realization A'_0, \dots, A'_{n-1} of E . By Remark 3.7, we can assume each of A'_0, \dots, A'_{n-1} to be nonmeager.

By Lemma 2.3, there is some realization A''_0, \dots, A''_{n-1} of E on the 2-sphere witnessed using (*). To finish our proof, replace A''_i with A'_i on D to obtain a Baire measurable realization of E on the 2-sphere. That is, set $A_i = (A''_i \cap D^c) \cup (A'_i \cap D)$. \square

4. BOREL PATH DECOMPOSITIONS FROM BOREL END SELECTIONS

Suppose $f: X \rightarrow X$. Say that f is **aperiodic** if for all $x \in X$ and $n \geq 1$, we have $f^n(x) \neq x$. Let G_f be the graph induced by f where distinct $x_0, x_1 \in X$ are G_f -adjacent if $f(x_0) = x_1$ or $f(x_1) = x_0$. Suppose $A \subseteq X$. Say that A is **forward recurrent** (with respect to f) if for every $x \in X$ there exists some $n \geq 0$ such that $f^n(x) \in A$.

We have the following lemma showing that bounded-to-one Borel functions admit forward recurrent r -independent sets.

Lemma 4.1. *Suppose X is a standard Borel space and $f: X \rightarrow X$ is an aperiodic bounded-to-one Borel function. Then for every $r \geq 1$ there exists a Borel set $A \subseteq X$ that is forward recurrent and r -independent.*

Proof. Let $G_f^{\leq r}$ be the graph on X where distinct $x, y \in X$ are $G_f^{\leq r}$ -adjacent if $d(x, y) \leq r$. Since G_f has bounded degree, $G_f^{\leq r}$ also has bounded degree. Hence,

by [KST, Theorem 4.6], there is a Borel coloring c of $G_f^{\leq r}$ with finitely many colors. Let A be the set of $x \in X$ such that $c(x)$ is equal to the least number appearing infinitely often in the sequence $c(x), c(f(x)), c(f^2(x)) \dots$. Then for each x , all the elements of A in the (G^-) -connected component of x have the same color, and hence A is r -independent, since c is a coloring of $G_f^{\leq r}$. A is forward recurrent by construction. \square

Now we show that we can obtain Borel path decompositions from Borel end selections

Lemma 4.2. *Suppose G is a bounded degree Borel graph on X such that there is a Borel selection of finitely many ends in every connected component of G . Then for every $n > 0$, G admits a Borel path decomposition of length at least n .*

Proof of Lemma 4.2. We are given a bounded degree acyclic Borel graph G on a standard Borel space X where every vertex has degree at least 2. First, by [HM, Theorem C] which builds on methods from [M], if there is a Borel function selecting finitely many ends from every connected component of G , then there is a Borel function selecting one or two ends in every connected component of G . Hence, we can partition X into two G -invariant Borel sets C_1, C_2 so that $G \upharpoonright C_1$ is generated by a bounded-to-one Borel function $f: C_1 \rightarrow C_1$, and there is a Borel set $B_2 \subseteq C_2$ so that every connected component of $G \upharpoonright B_2$ is 2-regular and every connected component of $G \upharpoonright C_2$ contains exactly one connected component of $G \upharpoonright B_2$.

By Lemma 4.1, we can find a forward recurrent Borel set $A \subseteq C_1$ such that A is $2n$ -independent in G . Let P_0^1 be the set of lex-least paths of length n which begin at some vertex of A , and let B_1 be the set of vertices contained in some element of P_0^1 . If $x \in C_1 \setminus B_1$, let $[x]$ be the set of vertices y for which there is a path p from x to y for which no interior vertex of p is in B_1 . The forward recurrence of A implies that for every $x \in C_1$, there is a unique forward-most element of $[x]$ under f . It is also clear that $G \upharpoonright [x]$ satisfies the hypothesis of Remark 3.2. For each x , the space of n -spindly decompositions is a compact space in the natural topology on all such decompositions. Hence, by compact uniformization [S, Theorem 5.7.1], see also [K, Theorem 18.18], there is a Borel way of selecting a unique path decomposition of length at least n for $G \upharpoonright [x]$ for each $x \in C_1 \setminus B_1$. Hence, we can extend P_0^1 to a Borel path decomposition of length at least n for $G \upharpoonright C_1$.

On $G \upharpoonright C_2$, we can first partition $G \upharpoonright B_2$ into a Borel set P_0^2 of finite paths of length at least n . If $x \in C_2 \setminus B_2$, let $[x]$ be the set of $y \in X$ such that there is a path p from x to y for which no interior vertex of p is in B_2 . Once again, $G \upharpoonright [x]$ is n -spindly. Hence by Remark 3.2 we can extend P_0^2 to a Borel path decomposition of length at least n for $G \upharpoonright C_2$. \square

Using Adams end selection, we can use this lemma to show that μ -hyperfinite free actions of \mathbb{F}_n have μ -measurable realizations of abstract systems of congruences that are non-complementing and non-expanding.

Theorem 4.3. *Suppose that $n \geq 2$, and a is a free Borel action of \mathbb{F}_n on a standard probability space (X, μ) that is μ -hyperfinite. Then there is a μ -measurable a -realization of every abstract system of congruences E that is non-complementing and non-expanding.*

Proof. Let R minimally generate E . Pass to a free subgroup $\mathbb{F}_k \leq \mathbb{F}_n$ where $k = |R|$. Let S be the set of generators of S . By a theorem of Adams [JKL, Lemma 3.21],

on a conull set there is a Borel function selecting either one or two ends from each connected component of $G(a, S)$. Hence, the theorem follows from Lemmas 4.2 and 3.6. \square

When we apply Lemma 4.2, it will be useful to know that end selections pass between finite index subgroups.

Lemma 4.4. *Suppose a is a free Borel action of a finitely generated group Γ on X . Let $\Delta \leq \Gamma$ be a finitely generated finite index subgroup of Γ , and b be the restriction of the action of a to Δ . Then if $S \subseteq \Gamma$ and $R \subseteq \Delta$ are finite symmetric generating sets, then $G(a, S)$ has a Borel selection of finitely many ends if and only if $G(b, R)$ has a Borel selection of finitely many ends.*

Proof. Since Δ is finite index in Γ , each $G(a, S)$ connected component contains finitely many components of $G(b, R)$, and each connected component of $G(b, R)$ is bounded distance from every point in the connected component of $G(a, S)$ it is contained in. Hence, there is an effectively defined bijection between ends in a connected component of $G(a, S)$, and ends in each $G(b, R)$ -component that it contains.

More precisely, suppose $r = (x_i)_{i \in \mathbb{N}}$ is a ray representing an end in $G(a, S)$, and C is a connected component of $G(b, R)$. We define a ray $f_C(r)$ in $G(b, R) \upharpoonright C$ as follows. To each x_i we associate the nearest point y_i in C , and let $f_C(r)$ be the lex least ray passing through all the points $(y_i)_{i \in \mathbb{N}}$, erasing loops. The map f_C clearly lifts to a map sending a selection of finitely many ends in $G(a, S)$ to a selection of finitely many ends in $G(b, R)$. The reverse implication is similar. \square

5. CONSTRUCTIVE REALIZATIONS OF NON-EXPANDING ABSTRACT SYSTEMS OF CONGRUENCES FOR $\mathrm{PSL}_2(\mathbb{Z})$ ACTING ON $\mathbb{P}^1(\mathbb{R})$

The group $\mathrm{PSL}_2(\mathbb{Z})$ acts on the space $\mathbb{P}^1(\mathbb{R})$ of lines in \mathbb{R}^2 through the origin. By identifying such a line with the x -value $x \in \mathbb{R} \cup \{\infty\}$ of its intersection point with the line $y = 1$, it is easy to see that this action is isomorphic to the action of $\mathrm{PSL}_2(\mathbb{Z})$ on $\mathbb{R} \cup \{\infty\}$ by fractional linear transformations, where $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ acts via $x \mapsto \frac{ax+b}{cx+d}$.

It is a standard fact that $\mathrm{PSL}_2(\mathbb{Z})$ is generated by the two transformations $\alpha(x) = x + 1$ and $\beta(x) = -1/x$.

The group $\mathrm{PGL}_2(\mathbb{Z})$ is index 2 over $\mathrm{PSL}_2(\mathbb{Z})$, and similarly is generated by $\alpha(x) = x + 1$ and $\gamma(x) = 1/x$. Note that $\beta(x) = \alpha^{-1}(\gamma(\alpha(\beta(\alpha^{-1}(x)))))) = (-1 + 1/(1 + 1/(x - 1))) = -1/x$.

Let Irr denote the set of irrational numbers. Each $x \in \mathrm{Irr}$ has a unique continued fraction expansion

$$x = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}$$

Where $a_0 \in \mathbb{Z}$ and $a_1, a_2, \dots \in \mathbb{Z}^+$ are positive integers. We note the continued fraction expansion of x as $(a_0; a_1, \dots)$. The following lemma is standard.

Lemma 5.1. *Let $f: \text{Irr} \rightarrow \text{Irr}$ be the function given by*

$$f(x) = \begin{cases} x - 1 & \text{if } x > 0 \\ 1/x & \text{if } x \in (0, 1) \\ x + 1 & \text{if } x < 0 \end{cases}$$

Then f generates the orbit equivalence relation of $\text{PGL}_2(\mathbb{Z})$ on Irr , and so $x, y \in \text{Irr}$ are in the same orbit if and only if their continued fraction expansions are tail equivalent.

Proof. The equivalence relation generated by f is clearly contained in the orbit equivalence relation, since f is defined piecewise by fractional linear transformations.

Recall that two continued fraction expansions $(a_0; a_1, \dots)$ and $(b_0; b_1, \dots)$ are tail equivalent if there exists some $n, m > 0$ such that $a_{n+i} = b_{m+i}$ for all $i \geq 0$. Since

$$f\left(a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}\right) = \begin{cases} (a_0 - 1) + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}} & \text{if } a_0 > 0 \\ a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}} & \text{if } a_0 = 0 \\ (a_0 + 1) + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}} & \text{if } a_0 < 0 \end{cases}$$

it is clear that if x and y are tail equivalent, then there are in the same equivalence class of the equivalence relation generated by f .

To finish, since $\alpha(x) = x + 1$ and $\gamma(x) = 1/x$ generate $\text{PSL}_2(\mathbb{Z})$, it suffices to show that if $x \in \text{Irr}$, then $x, x + 1$, and $1/x$ are tail equivalent. It is trivial to see that x and $x + 1$ are tail equivalent. That x and $1/x$ are tail equivalent is clear when $x > 0$. When $x < 0$, since one of x and $1/x$ are less than -1 , by swapping x and $1/x$, we may assume the continued fraction expansion of x is $x = a + \frac{1}{b+C}$, where $a \leq -2$, and $b \geq 1$. Then apply the following identity:

$$\frac{1}{a + \frac{1}{b+C}} = -1 + \frac{1}{1 + \frac{1}{(-a-2) + \frac{1}{1 + \frac{1}{(b-1)+C}}}}.$$

Note that $-a - 2 \geq 0$ and $b - 1 \geq 0$. If either of these two terms are equal to zero, this just removes the corresponding term in the continued fraction expansion, since $\frac{1}{0 + \frac{1}{a_n + C}} = a_n + C$. \square

Corollary 5.2. *Let a be the restriction of the action of $\text{PSL}_2(\mathbb{Z})$ to the irrationals. Let $S = \{s, t\}$ where $s, t \in \text{PSL}_2(\mathbb{Z})$ are the generators $x \mapsto 1/x$ and $x \mapsto x + 1$. Then there is a Borel selection of one end in each equivalence class of $G(a, S)$. \square*

The action of $\text{PSL}_2(\mathbb{Z})$ is free modulo a countable set, since if $x = (ax+b)/(cx+d)$, then x is the solution to a quadratic equation with integer coefficients. To finish, we need to analyze the countable set on which the action is nonfree.

Theorem 5.3 ([Ka, Theorem 2.3.2]). *Two non-identity element of $\text{PSL}_2(\mathbb{R})$ commute if and only if they have the same set of fixed points.*

We prove the following lemma.

Lemma 5.4. *If $x \in P^1(\mathbb{R})$ and $\text{Stab}(x)$ is the stabilizer of x in $\text{PSL}_2(\mathbb{Z})$, then $\text{Stab}(x)$ is abelian.*

Proof. Suppose that x is a fixed point of $z \mapsto \frac{az+b}{cz+d}$ and this transformation is not the identity. Assume that $c \neq 0$. Then using the fact that the transformation has determinant 1 we have that x is of the form

$$\frac{a-d \pm \sqrt{(a+d)^2 - 4}}{2c}$$

If x is rational, then we must have that $|a+d| = 2$. Hence each transformation in $\text{Stab}(x)$ only fixes x . Hence $\text{Stab}(x)$ is abelian by Theorem 5.3.

If x is irrational, then for each transformation in $\text{Stab}(x)$ we can represent x as $r+y$ where r is rational and y is irrational (and possibly negative). If we have two such representations, say $x = r+y$ and $x = r'+y'$, then we have $y-y'$ is rational and hence 0. It follows that $r=r'$. Now clearly $r-y = r'-y'$ is the other fixed point of the two transformations and so any two transformations in $\text{Stab}(x)$ have the same fixed points. Hence $\text{Stab}(x)$ is abelian again by Theorem 5.3.

Next assume that $c = 0$. Then the transformation is linear and since it has determinant 1, we must have $ad = 1$. It follows that $a = d = 1$. Since our transformation is not the identity, $b \neq 0$. It follows that x is the point at infinity. The argument above shows that the only transformations fixing the point at infinity are of the form $z \mapsto z+b$ where $b \in \mathbb{Z}$. Clearly any two such transformations commute. \square

We can now prove Theorem 1.5 from the introduction.

Proof of Theorem 1.5. Let R be a minimal relation generating E . Let $k = |R|$. There is a finite index copy of \mathbb{F}_2 in $\text{PSL}_2(\mathbb{Z})$ and hence a finite index copy of \mathbb{F}_k . Let the free generating set of \mathbb{F}_k be S . Let a be the restriction of the action of $\text{PSL}_2(\mathbb{Z})$ to this copy of \mathbb{F}_k . Let $F \subseteq X$ be the subset on which the action of F is nonfree. By Lemma 5.4, this action has abelian stabilizers, and so by Lemma 2.3, there is a realization of E witnessed by letting the generators S of F_k witness the elements of R . Since $X \setminus F$ is a subset of the quadratic irrationals, $X \setminus F$ is countable, and so the sets realization E on $X \setminus F$ are Borel.

Now on F , the graph $G(a \upharpoonright F, S)$ has a Borel selection of finitely many ends by Lemmas 5.2 and 4.4. Hence, by Lemma 4.2 we have a Borel path decomposition and hence by Lemma 3.6 there is a realization of E on $a \upharpoonright F$ once again with the i th generator witnesses the i th congruence in R . The Theorem follows by taking the union of these two realizations. \square

6. APPLICATIONS OF PATH DECOMPOSITIONS IN BOREL COMBINATORICS

If G is a locally finite acyclic Borel graph, then path decompositions for G give a very strong type of unfriendly coloring:

Lemma 6.1. *Suppose G is a locally finite acyclic Borel graph on X where every vertex has degree at least 2. Then if G has a Borel path decomposition of length at least 5, then G admits a Borel unfriendly coloring. Indeed, there is a Borel function $c: X \rightarrow 2$ such that for every x , $|\{y \in N(x): c(x) = c(y)\}| \leq 1$.*

Proof. Suppose P_0, P_1, \dots is the Borel path decomposition of G of length at least 5. We may assume that this path decomposition satisfies the conclusion of Lemma 3.5.

We inductively construct c . At step i we will ensure that every vertex in a path $p \in P_i$ has been colored. For all such paths $p \in P_i$, inductively, the only vertices in p that can have already been colored must be endpoints of p . Hence, there is some

extension of our partial coloring so that every vertex of p has at most one adjacent vertex of the same color, and the endpoint of p have neighbors of the opposite color. (For example, alternate between the two colors along p , possibly breaking parity once in the middle of the path). Since P_i is a path decomposition, each vertex is an interior vertex of at most one path, and every edge is contained in some path. Hence, our final coloring c of X has the desired property that each vertex has at most one neighbor of the same color. \square

By combining this Lemma with Lemma 3.4, we obtain Theorem 1.7 as a Corollary.

Suppose G is an acyclic locally finite Borel graph where every vertex has degree at least 3. Then an almost identical greedy construction shows that if G has a path decomposition of length at least 3, then G has a Borel perfect matching, and if G has maximum degree d , then G has a Borel d -list-coloring for any Borel assignment of lists to edges of G . For example, this gives a new way of proving a Baire measurable version of Vizing's theorem for acyclic bounded degree Borel graphs, and the existence of Baire measurable perfect matchings for acyclic locally finite Borel graphs.

REFERENCES

- [CK] C.T. Conley and A.S. Kechris, Measurable chromatic and independence numbers for ergodic graphs and group actions, *Groups Geom. Dyn.*, **7** (2013), 127–180.
- [D] R. Diestel, *Graph Theory*, Fourth, Graduate Texts in Mathematics, vol. 173, Springer, Heidelberg, (2010).
- [DF] R. Dougherty and M. Foreman, Banach-Tarski decompositions using sets with the property of Baire. *J. Amer. Math. Soc.*, **7**(1) (1994), 75–124.
- [DHK] L. Dubins, M. W. Hirsch, and J. Karush, *Scissor congruence*, Israel J. Math. **1** (1963), 239–247.
- [G] R. J. Gardner, *Convex bodies equidecomposable by locally discrete groups of isometries*, *Mathematika* **32** (1985), 1–9.
- [GJ] S. Gao and S. Jackson, *Countable abelian group actions and hyperfinite equivalence relations*. *Invent. Math.* **201** (2015), no. 1, 309–383.
- [GJKS] S. Gao, S. Jackson, E. Krohne, and B. Seward *Forcing constructions and countable Borel equivalence relations*. Preprint. 2015.
- [GMP16] L. Grabowski, A. Máthé, O. Pikhurko, *Measurable equidecompositions for group actions with an expansion property*, <https://arxiv.org/abs/1601.02958>.
- [GMP17] L. Grabowski, A. Máthé, O. Pikhurko, *Measurable circle squaring*, *Ann. of Math* **185** (2017), 671–710.
- [HM] G. Hjorth and B.D. Miller, *Ends of graphed equivalence relations, II*, *Israel Journal of Mathematics*, **169** (2009), 393–415.
- [JKL] S. Jackson, A.S. Kechris, and A. Louveau, *countable Borel equivalence relations*, *J. Math. Logic*, **2** (2002), 1–80.
- [Ka] S. Katok, *Fuchsian groups*, Chicago Lectures in Mathematics, 1992.
- [K] A.S. Kechris, *Classical descriptive set theory*, Springer, 1995.
- [KM] A.S. Kechris and A.S. Marks, *Descriptive graph combinatorics*. Preprint, 2016.
- [KST] A.S. Kechris, S. Solecki and S. Todorcevic, *Borel chromatic numbers*, *Adv. Math.*, **141** (1999), 1–44.
- [L88] M. Laczkovich, *Closed sets without measurable matching*, *P. Amer. Math. Soc.* **103** (1988), no. 3, 894–896.
- [L90] M. Laczkovich, *Equidecomposability and discrepancy; a solution of Tarski's circle-squaring problem*, *J. Reine Angew. Math.* **404** (1990), 77–117.
- [L92] M. Laczkovich, *Decomposition of sets with small boundary*, *J. Lond. Math. Soc.* **46** (1992), 58–64.

- [LN] R. Lyons and F. Nazarov, Perfect matchings as IID factors of non-amenable groups, *European J. Combin.*, **32** (2011), 1115–1125.
- [MU16] A. Marks and S. Unger, Baire measurable paradoxical decompositions via matchings, *Advances in Math.* **289** (2016), 397–410.
- [MU17] A. Marks and S. Unger, Borel circle squaring, *Ann. of Math.* 186 (2017), 581–605.
- [M] B.D. Miller, *Ends of graphed equivalence relations, I*, Israel Journal of Mathematics, **169(1)** (2009), 375–392.
- [N] M. Nadkarni, *On the existence of a finite invariant measure*. Proc. Indian Acad. Sci. Math. Sci., **100**, (1990) 203–220.
- [SS] S. Schneider and B. Seward, *Locally nilpotent groups and hyperfinite equivalence relations*, Preprint.
- [S] S.M. Srivastava, *A Course on Borel Sets*, Springer-Verlag, New York, (1998).
- [T] A. Tarski, Problème 38, *Fund. Math.* **7** (1925), 381.
- [W] S. Wagon, *The Banach Tarski Paradox*, University Press, Cambridge, 1986.
- [Weh] F. Wehrung, Baire paradoxical decompositions need at least six pieces, *Proc. Amer. Math. Soc.*, **121(2)** (1994), 643–644.

DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE MELLON UNIVERSITY, PITTSBURGH, PA 15213

Email address: `clintonc@andrew.cmu.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT LOS ANGELES

Email address: `marks@math.ucla.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT LOS ANGELES

Email address: `sunger@math.ucla.edu`