

21-720 Measure Theory: Midterm.

December 10th, 2013

- This is a closed book test. No calculators or computational aids are allowed.
- You have 3 hours. The exam has a total of 8 questions and 35 points.
- You may use any result from class or homework **PROVIDED** it is independent of the problem you want to use the result in. (You must also **CLEARLY** state the result you are using.)
- The questions are **ROUGHLY** in order of length / difficulty, and not in the order the material was covered. However, depending on your intuition, you might find a few of the later questions easier. Good luck!

In this exam, we always assume (X, Σ, μ) is a measure space. We use λ to denote the Lebesgue measure on \mathbb{R}^d .

- 5 1. Recall the maximal function of a function f is defined by $Mf(x) = \sup_{r>0} \frac{1}{\lambda(B(x,r))} \int_{B(x,r)} |f| d\lambda$. True or false:

If $f \in L^1$, then Mf is measurable.

Prove it, or find a counter-example.

- 5 2. Consider the function $f(x, y) = \frac{xy}{x^4 + y^4}$ when $(x, y) \neq 0$ and $f(0, 0) = 0$ defined on the rectangle $[-1, 1]^2$.

Consider each of the following integrals:

(a) $\int_{-1}^1 \left(\int_{-1}^1 f(x, y) d\lambda(x) \right) d\lambda(y)$ (b) $\int_{-1}^1 \left(\int_{-1}^1 f(x, y) d\lambda(y) \right) d\lambda(x)$ (c) $\int_{[-1,1]^2} f d\lambda$.

For each of the integrals above, decide if they exist (in the extended sense) or not. If yes, compute it. If not, prove it.

- 5 3. Let ϕ_n be an approximate identity. True or false:

For every $f \in L^1(\mathbb{R}^d)$ such that $\hat{f} \in L^1(\mathbb{R}^d)$ the sequence $(\phi_n * f)^\wedge$ is convergent in $L^1(\mathbb{R}^d)$.

Prove, or find a counter example. [Recall that (ϕ_n) is an approximate identity if $\phi_n \geq 0$, $\int_{\mathbb{R}^d} \phi_n d\lambda = 1$, and for every $\varepsilon > 0$ we have $\lim_{n \rightarrow \infty} \int_{|x|>\varepsilon} \phi_n(x) dx = 0$.]

- 5 4. Let $g \in L^1(X, \mu)$, and $f : X \rightarrow \mathbb{R}$ be measurable. Let (f_n) be a sequence of measurable functions such that $(f_n) \rightarrow f$ in measure, and for all $n \in \mathbb{N}$ we have $|f_n| \leq g$ almost everywhere. True or false:

The sequence (f_n) necessarily has a subsequence (f_{n_k}) such that $(f_{n_k}) \rightarrow f$ almost everywhere.

Prove it, or find a counter example.

- 5 5. Let $f : [0, 1] \rightarrow \mathbb{R}$ be absolutely continuous. Suppose further $f' \in L^2([0, 1])$. True or false:

For every $\varepsilon > 0$ there exists $\delta > 0$ such that $0 < |x - y| < \delta$ implies $|f(x) - f(y)| < \varepsilon \sqrt{|x - y|}$.

Prove it, or find a counter example.

- 5 6. True or false:

For every $E \in \mathcal{L}(\mathbb{R})$ with $0 < \lambda(E) < \infty$, we must have $\lim_{n \rightarrow \infty} \int_E \cos^2(nx) dx$ exists.

Prove it, or find a counter example.

- 5 7. Let $\varphi \in C_c(\mathbb{R}^d)$ be a non-negative radially decreasing function with $\int_{\mathbb{R}^d} \varphi = 1$. True or false:

For all $f \in L^1(\mathbb{R}^d)$, we must have $f * \varphi \leq Mf$.

Prove it, or find a counter example. [RECALL: We say φ is radially decreasing if there exists a decreasing function $h : [0, \infty) \rightarrow \mathbb{R}$ such that $\varphi(x) = h(|x|)$.]

If you've completed the remainder of this exam and have time to spare, here is a fun question. This is for your entertainment only, and *will not influence your grade*.

- 0 8. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is Lebesgue measurable and $f(x + y) = f(x) + f(y)$, then show f is continuous.