

# Putnam C.8

Po-Shen Loh

17 October 2011

## 1 Problems

**Putnam 1991/B1.** For each integer  $n \geq 0$ , let  $S(n) = n - m^2$ , where  $m$  is the greatest integer with  $m^2 \leq n$ . Define a sequence  $(a_k)_{k=0}^{\infty}$  by  $a_0 = A$  and  $a_{k+1} = a_k + S(a_k)$  for  $k \geq 0$ . For what positive integers  $A$  is this sequence eventually constant?

**Putnam 1991/B2.** Suppose  $f$  and  $g$  are non-constant, differentiable, real-valued functions defined on  $(-\infty, \infty)$ . Furthermore, suppose that for each pair of real numbers  $x$  and  $y$ ,

$$\begin{aligned}f(x+y) &= f(x)f(y) - g(x)g(y), \\g(x+y) &= f(x)g(y) + g(x)f(y).\end{aligned}$$

If  $f'(0) = 0$ , prove that  $(f(x))^2 + (g(x))^2 = 1$  for all  $x$ .

**Putnam 1991/B3.** Does there exist a real number  $L$  such that, if  $m$  and  $n$  are integers greater than  $L$ , then an  $m \times n$  rectangle may be expressed as a union of  $4 \times 6$  and  $5 \times 7$  rectangles, any two of which intersect at most along their boundaries?