

Putnam D.4

Po-Shen Loh

18 September 2011

1 Problems

Putnam 2002/A4. In Determinant Tic-Tac-Toe, Player 1 enters a 1 in an empty 3×3 matrix. Player 0 counters with a 0 in a vacant position, and play continues in turn until the 3×3 matrix is completed with five 1's and four 0's. Player 0 wins if the determinant is 0 and player 1 wins otherwise. Assuming both players pursue optimal strategies, who will win and how?

Putnam 2002/A5. Define a sequence by $a_0 = 1$, together with the rules $a_{2n+1} = a_n$ and $a_{2n+2} = a_n + a_{n+1}$ for each integer $n \geq 0$. Prove that every positive rational number appears in the set

$$\left\{ \frac{a_{n-1}}{a_n} : n \geq 1 \right\} = \left\{ \frac{1}{1}, \frac{1}{2}, \frac{2}{1}, \frac{1}{3}, \frac{3}{2}, \dots \right\}$$

Putnam 2002/A6. Fix an integer $b \geq 2$. Let $f(1) = 1$, $f(2) = 2$, and for each $n \geq 3$, define $f(n) = nf(d)$, where d is the number of base- b digits of n . For which values of b does

$$\sum_{n=1}^{\infty} \frac{1}{f(n)}$$

converge?