

Putnam D.13

Po-Shen Loh

20 November 2011

1 Problems

Putnam 1997/A1. A rectangle, $HOMF$, has sides $HO = 11$ and $OM = 5$. A triangle ABC has H as the intersection of the altitudes, O the center of the circumscribed circle, M the midpoint of BC , and F the foot of the altitude from A . What is the length of BC ?

Putnam 1997/A2. Players $1, 2, 3, \dots, n$ are seated around a table, and each has a single penny. Player 1 passes a penny to player 2, who then passes two pennies to player 3. Player 3 then passes one penny to Player 4, who passes two pennies to Player 5, and so on, players alternately passing one penny or two to the next player who still has some pennies. A player who runs out of pennies drops out of the game and leaves the table. Find an infinite set of numbers n for which some player ends up with all n pennies.

Putnam 1997/A3. Evaluate

$$\int_0^\infty \left(x - \frac{x^3}{2} + \frac{x^5}{2 \cdot 4} - \frac{x^7}{2 \cdot 4 \cdot 6} + \dots \right) \left(1 + \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} + \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots \right) dx.$$

Putnam 1997/A4. Let G be a group with identity e and $\phi : G \rightarrow G$ a function such that

$$\phi(g_1)\phi(g_2)\phi(g_3) = \phi(h_1)\phi(h_2)\phi(h_3)$$

whenever $g_1g_2g_3 = e = h_1h_2h_3$. Prove that there exists an element $a \in G$ such that $\psi(x) = a\phi(x)$ is a homomorphism (i.e. $\psi(xy) = \psi(x)\psi(y)$ for all $x, y \in G$).

Putnam 1997/A5. Let N_n denote the number of ordered n -tuples of positive integers (a_1, a_2, \dots, a_n) such that $1/a_1 + 1/a_2 + \dots + 1/a_n = 1$. Determine whether N_{10} is even or odd.

Putnam 1997/A6. For a positive integer n and any real number c , define x_k recursively by $x_0 = 0$, $x_1 = 1$, and for $k \geq 0$,

$$x_{k+2} = \frac{cx_{k+1} - (n-k)x_k}{k+1}.$$

Fix n and then take c to be the largest value for which $x_{n+1} = 0$. Find x_k in terms of n and k , $1 \leq k \leq n$.