Putnam $\Sigma.14$

Po-Shen Loh

23 November 2014

1 Problems

Putnam 2008/A4. Define $f: \mathbb{R} \to \mathbb{R}$ by

$$f(x) = \begin{cases} x & \text{if } x \le e \\ xf(\ln x) & \text{if } x > e. \end{cases}$$

Does $\sum_{n=1}^{\infty} \frac{1}{f(n)}$ converge?

Putnam 2008/A5. Let $n \geq 3$ be an integer. Let f(x) and g(x) be polynomials with real coefficients such that the points $(f(1), g(1)), (f(2), g(2)), \ldots, (f(n), g(n))$ in \mathbb{R}^2 are the vertices of a regular n-gon in counterclockwise order. Prove that at least one of f(x) and g(x) has degree greater than or equal to n-1.

Putnam 2008/A6. Prove that there exists a constant c > 0 such that in every nontrivial finite group G there exists a sequence of length at most $c \ln |G|$ with the property that each element of G equals the product of some subsequence. (The elements of G in the sequence are not required to be distinct. A *subsequence* of a sequence is obtained by selecting some of the terms, not necessarily consecutive, without reordering them; for example, 4, 4, 2 is a subsequence of 2, 4, 6, 4, 2, but 2, 2, 4 is not.)