

Putnam $\Sigma.4$

Po-Shen Loh

16 September 2012

1 Problems

Putnam 1995/A4. Suppose we have a necklace of n beads. Each bead is labeled with an integer and the sum of all these labels is $n - 1$. Prove that we can cut the necklace to form a string whose consecutive labels x_1, x_2, \dots, x_n satisfy

$$\sum_{i=1}^k x_i \leq k - 1 \quad \text{for } k = 1, 2, \dots, n.$$

Putnam 1995/A5. Let x_1, x_2, \dots, x_n be differentiable (real-valued) functions of a single variable t which satisfy

$$\begin{aligned} \frac{dx_1}{dt} &= a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ \frac{dx_2}{dt} &= a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ &\vdots \\ \frac{dx_n}{dt} &= a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n \end{aligned}$$

for some constants $a_{ij} > 0$. Suppose that for all i , $x_i(t) \rightarrow 0$ as $t \rightarrow \infty$. Are the functions x_1, x_2, \dots, x_n necessarily linearly dependent?

Putnam 1995/A6. Suppose that each of n people writes down the numbers 1,2,3 in random order in one column of a $3 \times n$ matrix, with all orders equally likely and with the orders for different columns independent of each other. Let the row sums a, b, c of the resulting matrix be rearranged (if necessary) so that $a \leq b \leq c$. Show that for some $n \geq 1995$, it is at least four times as likely that both $b = a + 1$ and $c = a + 2$ as that $a = b = c$.