

Putnam $\Sigma.12$

Po-Shen Loh

10 November 2012

1 Problems

Putnam 1992/B4. Let $p(x)$ be a nonzero polynomial of degree less than 1992 having no nonconstant factor in common with $x^3 - x$. Let

$$\frac{d^{1992}}{dx^{1992}} \left(\frac{p(x)}{x^3 - x} \right) = \frac{f(x)}{g(x)}$$

for polynomials $f(x)$ and $g(x)$. Find the smallest possible degree of $f(x)$.

Putnam 1992/B5. Let D_n denote the value of the $(n-1) \times (n-1)$ determinant

$$\begin{bmatrix} 3 & 1 & 1 & 1 & \cdots & 1 \\ 1 & 4 & 1 & 1 & \cdots & 1 \\ 1 & 1 & 5 & 1 & \cdots & 1 \\ 1 & 1 & 1 & 6 & \cdots & 1 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 1 & 1 & \cdots & n+1 \end{bmatrix}.$$

Is the set $\left\{ \frac{D_n}{n!} \right\}_{n \geq 2}$ bounded?

Putnam 1992/B6. Let \mathcal{M} be a set of real $n \times n$ matrices such that

- (i) $I \in \mathcal{M}$, where I is the $n \times n$ identity matrix;
- (ii) if $A \in \mathcal{M}$ and $B \in \mathcal{M}$, then either $AB \in \mathcal{M}$ or $-AB \in \mathcal{M}$, but not both;
- (iii) if $A \in \mathcal{M}$ and $B \in \mathcal{M}$, then either $AB = BA$ or $AB = -BA$;
- (iv) if $A \in \mathcal{M}$ and $A \neq I$, there is at least one $B \in \mathcal{M}$ such that $AB = -BA$.

Prove that \mathcal{M} contains at most n^2 matrices.