

Putnam $\Sigma.13$

Po-Shen Loh

16 November 2014

1 Problems

Putnam 2007/B4. Let n be a positive integer. Find the number of pairs P, Q of polynomials with real coefficients such that

$$(P(X))^2 + (Q(X))^2 = X^{2n} + 1$$

and $\deg P > \deg Q$.

Putnam 2007/B5. Let k be a positive integer. Prove that there exist polynomials $P_0(n), P_1(n), \dots, P_{k-1}(n)$ (which may depend on k) such that for any integer n ,

$$\left\lfloor \frac{n}{k} \right\rfloor^k = P_0(n) + P_1(n) \left\lfloor \frac{n}{k} \right\rfloor + \dots + P_{k-1}(n) \left\lfloor \frac{n}{k} \right\rfloor^{k-1}.$$

($\lfloor a \rfloor$ means the largest integer $\leq a$.)

Putnam 2007/B6. For each positive integer n , let $f(n)$ be the number of ways to make $n!$ cents using an unordered collection of coins, each worth $k!$ cents for some k , $1 \leq k \leq n$. Prove that for some constant C , independent of n ,

$$n^{n^2/2 - Cn} e^{-n^2/4} \leq f(n) \leq n^{n^2/2 + Cn} e^{-n^2/4}.$$