

7. Convergence

Po-Shen Loh

CMU Putnam Seminar, Fall 2018

1 Well-known statements

Infimum and supremum. The *infimum* of a set of numbers $S \subseteq \mathbb{R}$ is the largest real number x such that $x \leq s$ for all $s \in S$. The *supremum* is the smallest real number y such that $y \geq s$ for all $s \in S$.

Monotone sequence. Let $a_1 \geq a_2 \geq a_3 \geq \dots$ be a sequence of non-negative numbers. Then $\lim_{n \rightarrow \infty} a_n$ exists.

Alternating series. Let a_1, a_2, \dots be a sequence of real numbers which is decreasing in absolute value, converging to 0, and alternating in sign. Then the series $\sum_{i=1}^{\infty} a_i$ converges.

Conditionally convergent. Let a_1, a_2, \dots be a sequence of real numbers such that the series $\sum_{i=1}^{\infty} a_i$ converges to a finite number, but $\sum_{i=1}^{\infty} |a_i| = \infty$. This is called a *conditionally convergent* series. Then, for any real number r , the sequence can be rearranged so that the series converges to r . (Formally, there exists a bijection $\sigma : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ such that $\lim_{n \rightarrow \infty} \sum_{i=1}^n a_{\sigma(i)} = r$.)

2 Problems

1. Which of these series converge or diverge: $\sum_{n=1}^{\infty} \frac{1}{n}$, $\sum_{n=1}^{\infty} \frac{1}{2n}$, $\sum_{n=1}^{\infty} \frac{1}{n \log n}$, $\sum_{n=1}^{\infty} \frac{1}{n \log n \log n}$, ...?
2. Prove that the equation $x^{x^{x^{\dots}}} = 2$ is satisfied by $x = \sqrt{2}$, but that the equation $x^{x^{x^{\dots}}} = 4$ has no solution. What is the “break-point”?
3. The expression $\sqrt{-\frac{2}{9} + \sqrt{-\frac{2}{9} + \sqrt{-\frac{2}{9} + \dots}}}$ is not clearly defined. Consider the recursion $a_{n+1} = \sqrt{-\frac{2}{9} + a_n}$ with $a_1 = 0.34$. What does that sequence converge to, and why?
4. Suppose that a_1, a_2, \dots is a sequence of real numbers such that for each k , a_k is either $\frac{1}{k}$ or $-\frac{1}{k}$, and a_k has the same sign as a_{k+8} . Show that if four of a_1, a_2, \dots, a_8 are positive, then $\sum_{k=1}^{\infty} a_k$ converges. Is the converse true?
5. Prove that a sequence of positive numbers, each of which is less than the average of the previous two, is convergent.
6. Prove that every sequence of real numbers contains a monotone subsequence. Formally, show that for every sequence a_1, a_2, \dots of real numbers, there is a function $f : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ such that $f(i) < f(j)$ for all $i < j$, and either $a_{f(1)} \leq a_{f(2)} \leq a_{f(3)} \leq \dots$ or $a_{f(1)} \geq a_{f(2)} \geq a_{f(3)} \geq \dots$.
7. Given a convergent series of positive terms, $\sum_{k=1}^{\infty} a_k$, does the series $\sum_{k=1}^{\infty} \frac{a_1 + \dots + a_k}{k}$ always converge? How about the series $\sum_{k=1}^{\infty} \sqrt[k]{a_1 \times \dots \times a_k}$?

8. Let a_1, a_2, \dots be a sequence of positive real numbers satisfying $a_n < a_{n+1} + a_{n^2}$ for all n . Prove that $\sum a_n$ diverges.
9. Let a_i be real numbers such that $\sum_1^\infty a_i = 1$ and $|a_1| > |a_2| > |a_3| > \dots$. Suppose that f is a bijection from the positive integers to itself, and

$$|f(i) - i||a_i| \rightarrow 0$$

as $i \rightarrow \infty$. Prove or disprove that $\sum_1^\infty a_{f(i)} = 1$.

3 Homework

Please write up solutions to two of the problems, to turn in at next week's meeting. One of them may be a problem that we discussed in class. You are encouraged to collaborate with each other. Even if you do not solve a problem, please spend two hours thinking, and submit a list of your ideas.