

21-121 Calculus
Review Session for Exam 2
Cherry Wan
cwan@andrew.cmu.edu

$u/2$
 $7-8=30$
DH2210

Concept 1: Fundamental Theorem of Calculus

If f is continuous on $[a,b]$, then the function g defined by

PART I $g(x) = \int_a^x f(t) dt \quad a \leq x \leq b$

is continuous on $[a,b]$ and differentiable on (a,b) , and $g'(x) = f(x)$

If f is continuous on $[a,b]$, then

PART II $\int_a^b f(x) dx = F(b) - F(a)$

where F is any antiderivative of f , that is, a function such that $F' = f$.

Exercises

Use Part I of FTC to find the derivative of the following functions

1. $f(x) := \int_x^{10} \tan(t) dt$

2. $y = \int_1^{1-3x} \frac{u^3}{1+u^2} du$

Concept 2: Net Change Theorem

~ It's basically Part II of FTC

Exercises

Note: distance is the total amount travelled

displacement is just the change from the beginning to the end

3. Find the distance and displacement travelled given the following velocity function during the given time interval

$$v(t) := t^2 - 2t - 8 \quad 0 \leq t \leq 6$$

Concept 3: Integration by Substitution

Exercises

4.
$$\int x^2 (1 + 2 \cdot x^3)^3 dx$$

5.
$$\int_1^2 \frac{1}{x} \frac{e^x}{x^2} dx$$

6. $\int \sec(x) \cdot \tan(x) \cdot \sqrt{1 + \sec(x)} \, dx$

7. $\int_{e^1}^{e^4} \frac{1}{x \cdot \sqrt{\ln(x)}} \, dx$

Concept 4: Integration by Parts

$$\int u \, dv = uv - \int v \, du$$

Note: you want to pick u so that its derivative is simpler

Exercises

8. $\int (2x + 3) \cdot e^x \, dx$

9. $\int \cos(\ln(x)) \, dx$

Concept 5: Trigonometric Integrals

Rules

$$\int \sin^m(x) \cdot \cos^n(x) dx$$

If the power for cosine is odd, save one cosine factor and use $\cos^2(x) = 1 - \sin^2(x)$

Then substitute $u = \sin(x)$

If the power of sine is odd, save one sine factor and use $\sin^2(x) = 1 - \cos^2(x)$

Then substitute $u = \cos(x)$

If the powers of both sine and cosine are even, use the half-angle identities

$$\sin^2(x) = \frac{1}{2} \cdot (1 - \cos(2x))$$

$$\sin x \cdot \cos x = \frac{1}{2} \cdot \sin(2x)$$

$$\cos^2(x) = \frac{1}{2} \cdot (1 + \cos(2x))$$

$$\int \tan^m(x) \sec^n(x) dx$$

If the power of secant is even, save a factor of $\sec^2(x)$ and use $\sec^2(x) = 1 + \tan^2(x)$ to express the remaining factors in terms of $\tan(x)$

Then substitute $u = \tan(x)$

If the power of tangent is odd, save a factor of $\sec(x)\tan(x)$ and use $\tan^2(x) = \sec^2(x) - 1$ to express the remaining factors in terms of $\sec(x)$

Then substitute $u = \sec(x)$

$$\int \tan(x) dx = \ln(|\sec(x)|) + C$$

$$\int \sec(x) dx = \ln(|\sec(x) + \tan(x)|) + C$$

For the following cases on the left, considering using the properties on the right

$$\int \sin(mx) \cdot \cos(nx) \, dx$$

$$\sin(A) \cdot \cos(B) = \frac{1}{2} \cdot (\sin(A - B) + \sin(A + B))$$

$$\int \sin(mx) \cdot \sin(nx) \, dx$$

$$\sin(A) \cdot \sin(B) = \frac{1}{2} \cdot (\cos(A - B) - \cos(A + B))$$

$$\int \cos(mx) \cdot \cos(nx) \, dx$$

$$\cos(A) \cdot \cos(B) = \frac{1}{2} \cdot (\cos(A - B) + \cos(A + B))$$

Exercises

10. $\int \cos^5(x) \cdot \sin^4(x) \, dx$

11. $\int \cos^7(x) \cdot \sin^9(x) \, dx$

Concept 8: Trigonometric Substitution

Table of Trigonometric Substitution

Identity	Substitution	Expression
$1 - \sin^2(\theta) = \cos^2(\theta)$	$\frac{\pi}{2} \geq \theta \geq \frac{\pi}{2}$	$x = a \sin(\theta)$ $\sqrt{a^2 - x^2}$
$1 + \tan^2(\theta) = \sec^2(\theta)$	$\frac{\pi}{2} \geq \theta \geq \frac{\pi}{2}$	$x = a \tan(\theta)$ $\sqrt{a^2 + x^2}$
$\sec^2(\theta) - 1 = \tan^2(\theta)$	$0 \leq \theta \leq \frac{\pi}{2}$	$x = a \sec(\theta)$ $\sqrt{x^2 - a^2}$

$$12. \int \cos^4(x) \cdot \sin^2(x) dx$$

$$13. \int \tan^5(x) \cdot \sec^4(x) dx$$

$$14. \int \cos(7x) \cdot \cos(5x) dx$$

$$15. \int e^x \cdot \sin^7(e^x) dx$$

Concept 6: Trigonometric Substitution

Table of Trigonometric Substitutions

Expression	Substitution	Identity
$\sqrt{a^2 - x^2}$	$x = a \cdot \sin(\theta) \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$	$1 - \sin^2(\theta) = \cos^2(\theta)$
$\sqrt{a^2 + x^2}$	$x = a \cdot \tan(\theta) \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$	$1 + \tan^2(\theta) = \sec^2(\theta)$
$\sqrt{x^2 - a^2}$	$x = a \cdot \sec(\theta) \quad 0 \leq \theta \leq \frac{\pi}{2} \text{ or } \pi \leq \theta \leq \frac{3\pi}{2}$	$\sec^2(\theta) - 1 = \tan^2(\theta)$

Exercises

16. $\int x^3 \sqrt{9-x^2} dx$

17. $\int \frac{x^3}{\sqrt{x^2+9}} dx$

18. $\int \frac{1}{x^2 \sqrt{x^2-9}} dx$

19. $\int \frac{1}{\sqrt{t^2-6t+13}} dt$

Concept B: Area between Curves

The area between two curves $y = f(x)$ and $y = g(x)$ and between $x = a$ and $x = b$ is

$$A = \int_a^b |f(x) - g(x)| dx$$

Concept 7: Integration of Rational Functions

Note: Have to check if function is proper!

Four cases:

- (1) The denominator $Q(x)$ is a product of distinct linear factors
- (2) $Q(x)$ is a product of linear factors, some of which are repeated
- (3) $Q(x)$ contains irreducible quadratic factors, none of which is repeated
- (4) $Q(x)$ contains a repeated irreducible quadratic factor

Exercises

20.
$$\int \frac{1}{x^2 - a^2} dx$$

$a \neq 0$

21.
$$\int \frac{1}{x^2 + a^2} dx$$

14.
$$\int \cos(7x) \cos(5x) dx$$

15.
$$\int e^x \sin(e^x) dx$$

22.
$$\int \frac{x^3}{x^3 + 2x^2 + x} dx$$

23.
$$\int \frac{1}{x^3 + 1} dx$$

Concept 8: Trigonometric Substitution

Table of Trigonometric Substitutions

Expression

Substitution

Identity

$$\sqrt{a^2 - x^2}$$

$$x = a \sin(\theta) \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2}$$

$$1 - \sin^2(\theta) = \cos^2(\theta)$$

$$\sqrt{a^2 + x^2}$$

$$x = a \tan(\theta) \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2}$$

$$1 + \tan^2(\theta) = \sec^2(\theta)$$

$$\sqrt{x^2 - a^2}$$

$$x = a \sec(\theta) \quad 0 < \theta < \frac{\pi}{2} \text{ or } \frac{3\pi}{2} < \theta < 2\pi$$

$$\sec^2(\theta) - 1 = \tan^2(\theta)$$

$$24 \int \frac{(1-x+2x^2-x^3)}{x(x^2+1)^2} dx$$

$$25 \int \frac{1}{x^2-2x} dx$$

$$26 \int \frac{(2x+1)}{4x^2+12x-7} dx$$

$$27 \int \frac{(x^3-2x^2+x+1)}{x^4-3x^2-4} dx$$

Concept 8: Areas between Curves

The area between the curves $y = f(x)$ and $y = g(x)$ and between $x=a$ and $x = b$ is

$$A = \int_a^b |f(x) - g(x)| dx$$

Exercises Integration of Rational Functions

NOTE: Have to check if the denominator is prime

Set up the area between the following curves

28. $y = 2x + 1$ 29. $x + y = 1$
 $y = 9 - x^2$ $x = 2y^2$

Exercises

20. $\int \frac{1}{x^2 - 2} dx$
 $x > 0$

21. $\int \frac{1}{x^2 + 2} dx$

30. $x = 1 - y^2$
 $x = y^2 - 1$

31. $y = x^3 - x$
 $y = 3x$

22. $\int \frac{x^2}{x^2 + 2x^2 + x} dx$

23. $\int \frac{1}{x+1} dx$

Concept B: Area between Curves

The area between two curves $y = f(x)$ and $y = g(x)$ and between $x = a$ and $x = b$ is

$$A = \int_a^b |f(x) - g(x)| dx$$