

Test 0 Solutions
June 30

Name:

1. Evaluate the following integral:

$$\int_1^{e^3} \frac{(\ln x)^2}{x} dx =$$

If we let $u = \ln x$, then $du = \frac{1}{x} dx$, and

$$\begin{aligned} \int_1^{e^3} \frac{(\ln x)^2}{x} dx &= \int_0^3 u^2 du \\ &= \left[\frac{u^3}{3} \right]_0^3 \\ &= 9 - 0 = 9 \end{aligned}$$

2. Find an antiderivative:

$$\int \sin(\pi t) \cos(\pi t) dt =$$

If we let $u = \sin(\pi t)$, then $du = \pi \cos(\pi t) dt$, and

$$\begin{aligned} \int \sin(\pi t) \cos(\pi t) dt &= \int \frac{u}{\pi} du \\ &= \frac{u^2}{2\pi} + C \\ &= \frac{\sin^2(\pi t)}{2\pi} + C \end{aligned}$$

3. Put the following quantities in order, from least to greatest. Briefly explain your choices.

$$A = \int_0^{\pi/2} \cos(t) dt$$

$$B = \int_0^{\pi} \cos(t) dt$$

$$C = \int_0^3 \cos(t) dt$$

$$D = \int_0^6 \cos(t) dt$$

$$E = \int_0^{3\pi/2} \cos(t) dt$$

E, D, B, C, A

4. Let

$$K = \int_{-9}^4 3x^2 dx$$

- (a) Give an approximation for K with 39 rectangles and left endpoints.
- (b) Is your approximation an underestimate or an overestimate? Explain why.
- (c) Express K as a limit of sums.

(a)

$$\sum_{i=0}^{38} \frac{1}{3} \cdot 3\left(-9 + \frac{1}{3}i\right)^2$$

- (b) The approximation is an overestimate. We are overestimating from -9 to 0, and underestimating from 0 to 4, but overall we are overestimating.

(c)

$$K = \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{13}{n} \cdot 3\left(-9 + \frac{13}{n}i\right)^2$$

5. Do two of the following four problems:

- (a) Find all points on the graph of the function

$$f(x) = 2 \sin(x) + \sin^2(x)$$

at which the tangent line is horizontal.

- (b) Find the derivative of the function $f(x) = x^3$ at $x = 1$ directly from the definition.
 (c) Evaluate

$$\lim_{x \rightarrow \pi} \frac{e^{\sin x} - 1}{x - \pi}$$

- (d) Find an equation of the tangent line to the curve $y = e^x \cos x$ at the point $(0, 1)$.

- (a) The points where the tangent line is horizontal are the points where $f'(x) = 0$, i.e. where

$$0 = 2 \cos(x) + 2 \sin(x) \cos(x) = 2 \cos(x)[1 + \sin(x)]$$

Thus the tangent line is horizontal when $2 \cos(x) = 0$ (at $x = \frac{\pi}{2} + \pi n$) or when $\sin(x) = -1$ (at $x = \frac{3\pi}{2} + 2\pi n$).

- (b)

$$\begin{aligned} f'(1) &= \lim_{h \rightarrow 0} \frac{(1+h)^3 - 1^3}{h} \\ &= \lim_{h \rightarrow 0} \frac{1^3 + 3 \cdot 1^2 \cdot h + 3 \cdot 1 \cdot h^2 + h^3 - 1}{h} \\ &= \lim_{h \rightarrow 0} \frac{3h + 3h^2 + h^3}{h} \\ &= \lim_{h \rightarrow 0} 3 + 3h + h^2 \\ &= 3 \end{aligned}$$

- (c) Use L'Hopital's Rule:

$$\begin{aligned} \lim_{x \rightarrow \pi} \frac{e^{\sin x} - 1}{x - \pi} &= \lim_{x \rightarrow \pi} \frac{\cos(x)e^{\sin x}}{1} \\ &= \cos \pi e^{\sin \pi} \\ &= -1 \cdot 1 = -1 \end{aligned}$$

- (d) By the product rule,

$$y'(0) = -e^0 \sin(0) + \cos(0)e^0 = 0 + 1 = 1$$

Thus the tangent line passes through $(0, 1)$ and has slope 1. An equation for this line is $y = x + 1$.

6. Let f be a differentiable, non-constant, function with the property that

$$\int_0^x f(t)dt = [f(x)]^2$$

What function is f ?

If we differentiate the integral, by the Fundamental Theorem, we get $f(x)$. Thus differentiating both sides with respect to x , we get

$$f(x) = 2f(x)f'(x)$$

Thus $0 = f(x)[2f'(x) - 1]$, which implies $f(x) = 0$ or $f'(x) = 1/2$ for all x . Since f is non-constant, it can't be identically 0, and thus $f(x) = x/2 + C$. Plugging back into the original equation, we find $C = 0$, and so conclude $f(x) = x/2$.