

Math 720: Homework.

Please be aware of the late homework, and academic integrity policies in the syllabus. In particular, you may collaborate, but must write up solutions on your own. You may only turn in solutions you understand. I also recommend doing (but not turning in) the optional problems. They often involve useful concepts that will come in handy as the semester progresses.

Assignment 1 (assigned 2020-09-02, due 2020-09-09).

- Let μ be a positive measure on (X, Σ) .
 - If $A_i \in \Sigma$ are such that $A_i \subseteq A_{i+1}$, show that $\mu(\bigcup_{i=1}^{\infty} A_i) = \lim_{i \rightarrow \infty} \mu(A_i)$.
 - If $A_i \in \Sigma$ are such that $A_i \supseteq A_{i+1}$, show that $\mu(\bigcap_{i=1}^{\infty} A_i) = \lim_{i \rightarrow \infty} \mu(A_i)$, provided $\mu(A_1) < \infty$. Is it still true if $\mu(A_1) = \infty$?
- Prove any open subset of $\mathbb{R}^d$ is a countable union of cells.
- For each of the following sets, compute the Lebesgue outer measure.
 - Any countable set.
 - The Cantor set.
 - $\{x \in [0, 1] \mid x \notin \mathbb{Q}\}$.
- If $V \subset \mathbb{R}^d$ is a subspace with $\dim(V) < d$, then show that $\lambda^*(V) = 0$.
 - If $P \subset \mathbb{R}^2$ is a polygon show that $\text{area}(P) = \lambda^*(P)$.
- Does there exist a σ -algebra whose cardinality is countably infinite? Disprove, or find an example.

Optional problems and details I omitted in class.

- * Define $\mu(A)$ to be the number of elements in A . Show that μ is a measure on $(X, \mathcal{P}(X))$. (This is called the counting measure.)
- * Let $x_0 \in X$ be fixed. Define $\delta_{x_0}(A) = 1$ if $x_0 \in A$ and 0 otherwise. Show that δ_{x_0} is a measure on $(X, \mathcal{P}(X))$. (This is called the delta measure at x_0 .)
- * Show that $\lambda^*(a + E) = \lambda^*(E)$ for all $a \in \mathbb{R}^d$, $E \subset \mathbb{R}^d$.
- * Show that $\lambda^*(I) = \ell(I)$ for all cells. (I only proved it for closed cells in class.)
- * Show that $\mathcal{B}(\mathbb{R})$ has the same cardinality as $\mathbb{R}$.
- * (Challenge) Suppose $f_n : [0, 1] \rightarrow [0, 1]$ are all Riemann integrable, $0 \leq f_n \leq 1$ and $(f_n) \rightarrow 0$ pointwise. Show that $\lim_{n \rightarrow \infty} \int_0^1 f_n = 0$, using only standard tools from Riemann integration.

Assignment 2 (assigned 2020-09-09, due 2020-09-16).

- Say μ is a *translation invariant* measure on $(\mathbb{R}^d, \mathcal{L})$ (i.e. $\mu(x + A) = \mu(A)$ for all $A \in \mathcal{L}$, $x \in \mathbb{R}^d$) which is finite on bounded sets. Show that $\exists c \geq 0$ such that $\mu(A) = c\lambda(A)$.
 - Let $T : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a linear transformation, and $A \in \mathcal{L}$. Show that $T(A) \in \mathcal{L}$ and $\lambda(T(A)) = |\det(T)|\lambda(A)$.
Hint: Express T in terms of elementary transformations.
- Let $\mathcal{E} \subset \mathcal{P}(X)$, and $\rho : \mathcal{E} \rightarrow [0, \infty]$ be such that $\emptyset \in \mathcal{E}$, $\rho(\emptyset) = 0$. For any $A \subset X$ define

$$\mu^*(A) = \inf \left\{ \sum_1^{\infty} \rho(E_i) \mid E_i \in \mathcal{E}, \text{ and } A \subset \bigcup_1^{\infty} E_j \right\}.$$

Show that μ^* is an outer measure.

- Let (X, d) be any metric space, $\delta > 0$, $\alpha \geq 0$ and define

$$\mathcal{E}_{\delta} = \{A \subset X \mid \text{diam}(A) < \delta\} \quad \text{and} \quad \rho_{\alpha}(A) = \frac{\pi^{\alpha/2}}{\Gamma(1 + \frac{\alpha}{2})} \left(\frac{\text{diam}(A)}{2} \right)^{\alpha}.$$

Let $H_{\alpha, \delta}^*$ be the outer measure obtained with $\rho = \rho_{\alpha}$ and the collection of sets $\mathcal{E}_{\delta}$. Define $H_{\alpha}^* = \lim_{\delta \rightarrow 0} H_{\alpha, \delta}^*$. Show H_{α}^* is an outer measure and restricts to a measure H_{α} on a σ -algebra that contains all Borel sets. The measure H_{α} is called the *Hausdorff measure of dimension α* .

- If $X = \mathbb{R}^d$, and $\alpha = d$ show that H_d is a non-zero, finite constant multiple of the Lebesgue measure. [In fact $H_d = \lambda$ because of our choice of normalization constant, but the proof requires a little more work.]
- Let $S \in \mathcal{B}(X)$. Show that there exists (a unique) $d \in [0, \infty]$ such that $H_{\alpha}(S) = \infty$ for all $\alpha \in (0, d)$, and $H_{\alpha}(S) = 0$ for all $\alpha \in (d, \infty)$. This number is called the *Hausdorff dimension* of the set S .
- Compute the Hausdorff dimension of the Cantor set.
- Using notation from the previous question, let $\mathcal{S}_{\delta} = \{B(x, r) \mid x \in X, r \in (0, \delta)\}$. Using the collection of sets $\mathcal{S}_{\delta}$ and the function $\rho = \rho_{\alpha}$, we obtain an outer measure $S_{\alpha, \delta}^*$. As before one can show that $S_{\alpha}^* = \lim_{\delta \rightarrow 0} S_{\alpha, \delta}^*$ is an outer measure, and gives a Borel measure S_{α} .
 - Do there exist $c_1, c_2 \in (0, \infty)$ such that $c_1 S_{\alpha}(A) \leq H_{\alpha}(A) \leq c_2 S_{\alpha}(A)$ for all Borel sets A ?
 - Show by example $S_{\alpha} \neq H_{\alpha}$ in general. Prove or disprove it.
 - If $X = \mathbb{R}^d$ with the standard metric show that $S_d = \lambda$. [You may assume $\rho_d(B_r) = \lambda(B_r)$.]

Assignment 3 (assigned 2020-09-16, due 2020-09-23).

- Let μ, ν be two measures on (X, Σ) . Suppose $\mathcal{C} \subset \Sigma$ is a π -system such that $\mu = \nu$ on $\mathcal{C}$.
 - Suppose $\exists C_i \in \mathcal{C}$ such that $\bigcup_1^\infty C_i = X$ and $\mu(C_i) = \nu(C_i) < \infty$. Show that $\mu = \nu$ on $\sigma(\mathcal{C})$.
 - If we drop the finiteness condition $\mu(C_i) < \infty$ is the previous subpart still true? Prove or find a counter example.
- Let X be a metric space and μ a Borel measure on X . Suppose there exists a sequence of sets $B_n \subset X$ such that $\bar{B}_n \subset \overset{\circ}{B}_{n+1}$, $\bar{B}_n$ is compact, $X = \bigcup_1^\infty B_n$ and $\mu(B_n) < \infty$. Show that μ is regular. Further, for any $A \in \mathcal{B}(X)$ and $\varepsilon > 0$ show that there exists U open and C closed such that $C \subseteq A \subseteq U$ and $\mu(U - C) < \varepsilon$.
- Find $E \in \mathcal{B}(\mathbb{R})$ so that for all $a < b$, we have $0 < \lambda(E \cap (a, b)) < b - a$.
 - Let $\kappa \in (0, 1/2)$. Does there exist $E \in \mathcal{B}(\mathbb{R})$ such that for all $a < b \in \mathbb{R}$, we have $\kappa(b - a) \leq \lambda(I \cap (a, b)) \leq (1 - \kappa)(b - a)$? Prove it.
- Let $A \in \mathcal{L}(\mathbb{R}^d)$. Prove every subset of A is Lebesgue measurable $\iff \lambda(A) = 0$.
- Prove $\mathcal{B}(\mathbb{R}^{m+n}) = \sigma(\{A \times B \mid A \in \mathcal{B}(\mathbb{R}^m) \& B \in \mathcal{B}(\mathbb{R}^n)\})$.
 - Prove $\mathcal{L}(\mathbb{R}^{m+n}) \supsetneq \sigma(\{A \times B \mid A \in \mathcal{L}(\mathbb{R}^m) \& B \in \mathcal{L}(\mathbb{R}^n)\})$.
 - Show $\mathcal{L}(\mathbb{R}^2) \supsetneq \mathcal{B}(\mathbb{R}^2)$.

Optional problems, and details I omitted in class.

- * Is any σ -finite Borel measure on $\mathbb{R}^d$ regular?
- * Show that there exists $A \subset \mathbb{R}$ such that if $B \subset A$ and $B \in \mathcal{L}$ then $\lambda(B) = 0$, and further, if $B \subset A^c$ and $B \in \mathcal{L}$ then $\lambda(B) = 0$.
- * If $C_n \subset B(0, n+1) - \overline{B(0, n)}$ is closed, show that $\bigcup_1^\infty C_n$ is also closed.

We say $\mathcal{A} \subset \mathcal{P}(X)$ is an algebra if $\emptyset \in \mathcal{A}$, and $\mathcal{A}$ is closed under complements and finite unions. We say $\mu_0 : \mathcal{A} \rightarrow [0, \infty]$ is a (positive) *pre-measure* on $\mathcal{A}$ if $\mu_0(\emptyset) = 0$, and for any countable disjoint sequence of sets $A_i \in \mathcal{A}$ such that $\bigcup_1^\infty A_i \in \mathcal{A}$, we have $\mu_0(\bigcup_1^\infty A_i) = \sum_1^\infty \mu_0(A_i)$.

Namely, a pre-measure is a finitely additive measure on an algebra $\mathcal{A}$, which is also countably additive for disjoint unions *that belong to the algebra*.

- * (*Caratheodory extension*) If $\mathcal{A}$ is an algebra, and μ_0 is a pre-measure on $\mathcal{A}$, show that there exists a measure μ defined on $\sigma(\mathcal{A})$ that extends μ_0 .
- * (*An alternate approach to λ -systems.*) Let $\mathcal{M} \subset \mathcal{P}(X)$. We say $\mathcal{M}$ is a *Monotone Class*, if whenever $A_i, B_i \in \mathcal{M}$ with $A_i \subset A_{i+1}$ and $B_i \supset B_{i+1}$ then $\bigcup_1^\infty A_i \in \mathcal{M}$ and $\bigcap_1^\infty B_i \in \mathcal{M}$. If $\mathcal{A} \subset \mathcal{P}(X)$ is an algebra, then show that the *smallest* monotone class containing $\mathcal{A}$ is exactly $\sigma(\mathcal{A})$. [You should also address existence of a smallest monotone class containing $\mathcal{A}$.]

Assignment 4 (assigned 2020-09-23, due 2020-09-30).

- Let $C \subset \mathbb{R}^d$ be convex. Must C be Lebesgue measurable? Must C be Borel measurable? Prove or find counter examples. [The cases $d = 1$ and $d > 1$ are different.]
- Let (X, Σ, μ) be a measure space. For $A \in \mathcal{P}(X)$ define $\mu^*(A) = \inf\{\mu(E) \mid E \supset A \& E \in \Sigma\}$, and $\mu_*(A) = \sup\{\mu(E) \mid E \subset A \& E \in \Sigma\}$.
 - Show that μ^* is an outer measure.
 - Let $A_1, A_2, \dots \in \mathcal{P}(X)$ be disjoint. Show that $\mu_*(\bigcup_1^\infty A_i) \geq \sum_1^\infty \mu_*(A_i)$. [The set function μ_* is called an *inner measure*.]
 - Show that for all $A \subset X$, $\mu^*(A) + \mu_*(A^c) = \mu(X)$.
 - Let $A \subset \mathcal{P}(X)$ with $\mu^*(A) < \infty$. Show that $A \in \Sigma_\mu \iff \mu_*(A) = \mu^*(A)$.
- Let $f : [0, 1] \rightarrow [0, 1]$ be the Cantor function, and $g(x) = \inf\{f = x\}$. Show that f is Hölder continuous, and the range of g is contained in the Cantor set. What is the largest exponent α for which f is Hölder- α continuous?
- Let (X, Σ) be a measure space, and $f, g : X \rightarrow [-\infty, \infty]$ be measurable. Suppose whenever $g = 0$, $f \neq 0$, and whenever $f = \pm\infty$, $g \in (-\infty, \infty)$. Show that $\frac{f}{g} : X \rightarrow [-\infty, \infty]$ is measurable.
[Note that by the given data you will never get a ‘meaningless’ quotient of the form $\frac{0}{0}$ or $\frac{\pm\infty}{\pm\infty}$. The remainder of the quotients (e.g. $\frac{1}{\infty}$) can be defined in the natural manner.]
- Let $f_n : X \rightarrow \mathbb{R}$ be a sequence of measurable functions such that $(f_n) \rightarrow f$ almost everywhere (a.e.). Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a Borel function.
 - If for a.e. $x \in X$, g is continuous at $f(x)$, then show $(g \circ f_n) \rightarrow g \circ f$ a.e.
 - Is the previous part true without the continuity assumption on g ?

Optional problems, and details I omitted in class.

- * Prove that the completion Σ_μ we defined in class is the smallest μ -complete σ -algebra that contains Σ .
- * Show that $f : X \rightarrow [-\infty, \infty]$ is measurable if and only if any of the following conditions hold

(a) $\{f < a\} \in \Sigma$ for all $a \in \mathbb{R}$.	(c) $\{f \leq a\} \in \Sigma$ for all $a \in \mathbb{R}$.
(b) $\{f > a\} \in \Sigma$ for all $a \in \mathbb{R}$.	(d) $\{f \geq a\} \in \Sigma$ for all $a \in \mathbb{R}$.
- * Let (f_n) is a sequence of extended real valued measurable functions. Define $f(x) = \lim f_n(x)$ if the limit exists (even if the limit is $\pm\infty$), and $f(x) = 0$ otherwise. Show that f is measurable.
- * Let (X, Σ, μ) be a measure space, and $(X, \Sigma_\mu, \bar{\mu})$ it's completion. Show that $g : X \rightarrow [-\infty, \infty]$ is Σ_μ -measurable if and only if there exists two Σ -measurable functions $f, h : X \rightarrow [-\infty, \infty]$ such that $f = h$ μ -almost everywhere, and $f \leq g \leq h$ everywhere.

Assignment 5 (assigned 2020-09-30, due 2020-10-14).

- Let X be a metric space, and μ a regular Borel measure on X .
 - True or false: For any $f : X \rightarrow \mathbb{R}$ measurable and $\varepsilon > 0$ there exists $g : X \rightarrow \mathbb{R}$ continuous such that $\mu\{f \neq g\} < \varepsilon$? Prove it or find a counter example.
 - Do the previous subpart when $X = \mathbb{R}^d$.
- If $f \geq 0$ is measurable show that $\int_X f d\mu = 0 \iff f = 0$ almost everywhere.
- Let $g \geq 0$ be measurable, and define $\nu(A) = \int_A g d\mu$. Show that ν is a measure, and $\int_E f d\nu = \int_E fg d\mu$. [NOTATION: We say $d\nu = g d\mu$.]
- Suppose $I \subset \mathbb{R}^d$ is a cell, and $f : I \rightarrow \mathbb{R}$ is Riemann integrable. Show that f is measurable, Lebesgue integrable and that the Lebesgue integral of f equals the Riemann integral.
 - Is the previous subpart true if we only assume that an improper (Riemann) integral of f exists? Prove or find a counter example.
- For $p \in \mathbb{R}$ define $F(y) = \int_0^\infty \frac{\sin(xy)}{1+x^p} dx$.
 - For what $p \in \mathbb{R}$ is F defined? When defined, is F continuous? Prove it.
 - Show that F is differentiable for $p > 2$, and not differentiable when $p = 2$.
- Let for $n \in \mathbb{N}$ define $A_n = \bigcup_{k \in \mathbb{Z}} [\frac{2k}{2^n}, \frac{2k+1}{2^n})$. If $E \in \mathcal{B}(\mathbb{R})$ does $\lim_{n \rightarrow \infty} \lambda(A_n \cap E)$ exist? Prove it.

Assignment 6 (assigned 2020-09-30, due Never).

In light of your **MIDTERM** this homework is optional.

- Let μ be the counting measure on $\mathbb{N}$, and $f : \mathbb{N} \rightarrow \mathbb{R}$ a function.
 - If $\sum_1^\infty |f(n)| < \infty$, then show that $\sum_{n=1}^\infty f(n) = \int_{\mathbb{N}} f d\mu$.
 - If the series $\sum_{n=1}^\infty f(n)$ is conditionally convergent, show that $\int_{\mathbb{N}} f d\mu$ is not defined.
- Let X be a metric space $C \subset X$ be closed and $f : C \rightarrow \mathbb{R}$ be continuous.
 - If $0 \leq f \leq 1$, then show that there exists $F : X \rightarrow \mathbb{R}$ continuous such that $F(c) = f(c)$ for all $c \in C$. [HINT: Let $F(x) = f(x)$ for all $x \in C$, and $F(x) = \inf\{f(c) + \frac{d(x,c)}{d(x,C)} - 1 \mid c \in C\}$ for $x \notin C$.]
 - (Tietze extension theorem in metric spaces) Do the previous subpart without assuming $0 \leq f \leq 1$. [HINT: Put $g = \tan^{-1}(f)$, construct G by the previous subpart and set $F = \tan(G)$.]
- Find a Borel measurable function $f : [0, 1] \rightarrow \mathbb{R}$ which is not continuous almost everywhere.
- Let $s, t \geq 0$ be two simple functions. Show directly $\int_X (s+t) = \int_X s + \int_X t$.
 - Let $0 \leq s \leq t$ be two simple functions. Show $\int_X s \leq \int_X t$.
- Show directly $\int_X \alpha f = \alpha \int_X f$ for any $\alpha \in \mathbb{R}$ and integrable function f .

- Let $F : \mathbb{R}^d \rightarrow [0, \infty)$ be Lebesgue measurable. Show that $\int_{\mathbb{R}^d} F d\lambda < \infty$ if and only if for every sequence of measurable functions (f_n) such that $|f_n| \leq F$ almost everywhere, and (f_n) converges almost everywhere, we have $\lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} f_n d\lambda = \int_{\mathbb{R}^d} \lim_{n \rightarrow \infty} f_n d\lambda$.
 - Is the previous subpart true for arbitrary measure spaces?
- If f is a bounded measurable function and $\mu(X) < \infty$, then show $\int_X f d\mu = \inf\{\int_X t d\mu \mid t \geq f \text{ is simple}\}$.
 - If f, g are bounded measurable functions and $\mu(X) < \infty$ show directly that $\int_X (f+g) d\mu = \int_X f d\mu + \int_X g d\mu$.
- Let $f : [0, \infty) \rightarrow \mathbb{R}$ be a measurable function. We define the *Laplace Transform* of f to be the function $F(s) = \int_0^\infty \exp(-st)f(t) dt$ wherever defined. If f is continuous and bounded, compute $\lim_{s \rightarrow \infty} sF(s)$.
- (Pull back measures) Say ν is a measure on (Y, τ) and $f : X \rightarrow Y$ is surjective.
 - Show that $\Sigma = \{A \subset X \mid f(A) \in \tau\}$ need not be a σ -algebra. If Σ is a σ -algebra, show that $\mu(A) = \nu(f(A))$ need not be a measure on (X, Σ) .
 - Define instead $\Sigma = \{A \subset X \mid f^{-1}(f(A)) = A, \& f(A) \in \tau\}$, and $\mu(A) = \nu(f(A))$. Show that Σ is a σ -algebra and μ is a measure.
 - If $g \in L^1(Y, \nu)$, then show that $g \circ f \in L^1(X, \mu)$ and $\int_X g \circ f d\mu = \int_Y g d\nu$.
- (Linear change of variable) Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be integrable.
 - For any $y \in \mathbb{R}^d$ show that $\int_{\mathbb{R}^d} f(x+y) d\lambda(x) = \int_{\mathbb{R}^d} f(x) d\lambda(x)$.
 - If $T : \mathbb{R}^d \rightarrow \mathbb{R}^d$ an invertible linear transformation, and $E \in \mathcal{L}(\mathbb{R}^d)$. Show that

$$\int_{T^{-1}(E)} (f \circ T) |\det T| d\lambda = \int_E f d\lambda.$$

- Show that there exists $f : \mathbb{R} \rightarrow [0, \infty)$ Borel measurable such that $\int_a^b f d\lambda = \infty$ for all $a, b \in \mathbb{R}$ with $a < b \in \mathbb{R}$. [HINT: Let $g(x) = \mathbf{1}_{\{|x|<1\}} |x|^{-1/2}$, and define $h(x) = \sum_{m=-\infty}^\infty \sum_{n=1}^\infty 2^{-m-n} g(x-m/n)$. Now set $f = h^2 \mathbf{1}_{\{h < \infty\}}$.]
- Prove Hölder's inequality for $p = 1$ and $q = \infty$.
- If $p_i, q \in [1, \infty]$ with $\sum_1^N \frac{1}{p_i} = \frac{1}{q}$, show that $\|\prod_1^N f_i\|_q \leq \prod \|f_i\|_{p_i}$.
- Show that L^∞ is a Banach space.
- For $p \in [0, 1)$ show that you need not have $\|f+g\|_p \leq \|f\|_p + \|g\|_p$.
- Let $p, q \in (1, \infty)$, $\frac{1}{p} + \frac{1}{q} = 1$, $f \in L^p$ and $g \in L^q$. Show that $\int_X |fg| d\mu = \|f\|_p \|g\|_q$ if and only if there exists constants $\alpha, \beta \geq 0$ such that $\alpha f^p = \beta g^q$.
- If X is σ -finite, and $f \in L^\infty$ then show $\|f\|_\infty = \sup_{g \in L^1 - \{0\}} \frac{1}{\|g\|_1} \int_X fg d\mu$.
 - Show that the previous subpart is false if X is not σ -finite.

Assignment 7 (assigned 2020-10-14, due 2020-10-21).

1. (a) Suppose there exists $C < \infty$ such that $\int_{\mathbb{R}^d} fg \leq C \|f\|_p \|g\|_q$ for all $f \in L^p(\mathbb{R}^d)$ and $g \in L^q(\mathbb{R}^d)$. Show that $\frac{1}{p} + \frac{1}{q} = 1$.
 (b) If there exists $C < \infty$ such that $\|f\|_p \leq C \|\nabla f\|_q$ for all $f \in C_c^\infty(\mathbb{R}^d)$, find a relation between p, q and d .
2. (a) Suppose $\varphi: (a, b) \rightarrow \mathbb{R}$ is strictly convex, $f: X \rightarrow (a, b)$ is measurable, and $\mu(X) = 1$. Find a necessary and sufficient condition on f under which $\varphi(\int_X f d\mu) = \int_X \varphi \circ f d\mu$.
 (b) Use Jensen's inequality to prove Hölder's inequality for $p \in (1, \infty)$.
 (c) Hence (or otherwise) find necessary and sufficient conditions under which equality holds in Hölder's inequality.
3. (a) Suppose $p, q, r \in [1, \infty]$ with $p < q < r$. Prove that $L^p \cap L^r \subseteq L^q$. Further, find $\theta \in (0, 1)$ such that $\|f\|_q \leq \|f\|_p^\theta \|f\|_r^{1-\theta}$ for all $f \in L^p \cap L^q$.
 (b) If for some $p \in [1, \infty)$, $f \in L^p(X) \cap L^\infty(X)$ show that $\lim_{q \rightarrow \infty} \|f\|_q = \|f\|_\infty$.
 (c) If $\mu(X) = 1$ and $f \in L^1(X)$ show $\lim_{p \rightarrow 0^+} \|f\|_p = \exp(\int_X \ln|f| d\mu)$.
4. (a) For all $p \in [1, \infty)$, show that simple functions are dense in L^p .
 (b) Let X be a metric space and μ a regular Borel measure on X . Suppose there exists a sequence of sets $B_n \subset X$ such that $\bar{B}_n \subset \overset{\circ}{B}_{n+1}$, $\bar{B}_n$ is compact and $X = \bigcup_1^\infty B_n$. Show that $C_c(X)$ is dense in L^p for all $p \in [1, \infty)$.
 (c) Suppose $\Sigma = \sigma(\mathcal{C})$, where $\mathcal{C} \subset \mathcal{P}(X)$ is countable. If μ is a σ -finite measure and $1 \leq p < \infty$, show that $L^p(X)$ is separable (i.e. has a countable dense subset).
 (d) What happens to the previous three subparts when $p = \infty$?
5. (*Vitali Convergence Theorem*) If $f_n \in L^1$, $(f_n) \rightarrow f$ in measure, $\{f_n\}$ is both uniformly integrable and tight, then show that $f \in L^1$ and $(f_n) \rightarrow f$ in L^1 .

Assignment 8 (assigned 2020-10-21, due 2020-10-28).

1. (a) If $\mu(X) < \infty$, $1 \leq p < q \leq \infty$, show $L^q(X) \subset L^p(X)$ and the inclusion map from $L^q(X) \rightarrow L^p(X)$ is continuous. Find an example where $L^q(X) \subsetneq L^p(X)$. [HINT: Show $\|f\|_p \leq \mu(X)^{\frac{1}{p} - \frac{1}{q}} \|f\|_q$.]
 (b) Let $\ell^p = L^p(\mathbb{N})$ with respect to the counting measure. If $1 \leq p < q$ show that $\ell^p \subsetneq \ell^q$. Is the inclusion map $\ell^p \hookrightarrow \ell^q$ continuous? Prove your answer.
2. (a) Suppose $p \in [1, \infty)$, and $f \in L^p(\mathbb{R}^d, \lambda)$. For $y \in \mathbb{R}^d$, let $\tau_y f: \mathbb{R}^d \rightarrow \mathbb{R}$ be defined by $\tau_y f(x) = f(x - y)$. Show that $(\tau_y f) \rightarrow f$ in L^p as $|y| \rightarrow 0$.
 (b) What happens for $p = \infty$?
3. For $p \in [1, \infty)$ define $\|f\|_{L^{p,\infty}} = \sup_{\lambda > 0} \lambda \mu(\{|f| > \lambda\})^{1/p}$, and the weak L^p space (denoted by $L^{p,\infty}$) by $L^{p,\infty} = \{f \mid \|f\|_{L^{p,\infty}} < \infty\}$. [As usual, we use the convention that functions that are equal almost everywhere are identified with each other.]
 (a) If $f \in L^p$, show $f \in L^{p,\infty}$ and $\|f\|_{L^{p,\infty}} \leq \|f\|_p$. Is the converse true?
 (b) If $f, g \in L^{p,\infty}$, show that $f + g \in L^{p,\infty}$. Show further that $\|f + g\|_{L^{p,\infty}} \leq c(\|f\|_{L^{p,\infty}} + \|g\|_{L^{p,\infty}})$ for some constant c independent of f, g . [Thus $\|\cdot\|_{L^{p,\infty}}$ is called a quasi-norm, and $L^{p,\infty}$ is called a quasi-Banach space.]
 (c) If μ is σ -finite, $1 \leq p < q < r < \infty$ and $f \in L^{p,\infty} \cap L^{r,\infty}$ then show $f \in L^q$. [HINT: Show first that $\int_X |f| d\mu = \int_0^\infty \mu(|f| > t) dt$.]
4. (a) Suppose $\lim_{\lambda \rightarrow \infty} \sup_n \int_{|f_n| > \lambda} |f_n| d\mu = 0$. Show that there exists an increasing function φ with $\varphi(\lambda)/\lambda \rightarrow \infty$ as $\lambda \rightarrow \infty$, such that $\sup_n \int_X \varphi(|f_n|) < \infty$.
 (b) Suppose $\{f_n\}$ is uniformly integrable, and L^1 bounded (i.e. $\sup_n \int |f_n| < \infty$). Show that $\lim_{\lambda \rightarrow \infty} \sup_n \int_{|f_n| > \lambda} |f_n| = 0$.
 (c) Show that the previous part fails without the assumption $\sup_n \int |f_n| < \infty$.
5. Let $e_n(x) = e^{2\pi i n x}$ for $x \in \mathbb{R}$. Does (e_n) have a subsequence that converges almost everywhere? Prove it.

Optional problems, and details I omitted in class.

* Here is an alternate approach to the Hanh and Jordan decomposition.

(a) Let μ be a finite signed measure on (X, Σ) . For $A \in \Sigma$ define $|\mu|(A)$ by

$$|\mu|(A) = \sup \left\{ \sum_{B \in \pi} |\mu(B)| \mid \pi \text{ is a finite partition of } A \text{ into measurable sets.} \right\}$$

Show that $|\mu|$ is a finite positive measure on X .

(b) (*Jordan decomposition*) Show that any finite measure can be expressed uniquely as the difference of two mutually singular measures.

(c) (*Hanh decomposition*) Show that there exists a positive set $P \in \Sigma$ such that P^c is negative.

* Show that $\|\mu_n - \mu\| \rightarrow 0$ if and only if $(\mu_n(A)) \rightarrow \mu(A)$ uniformly in A , $\forall A \in \Sigma$.

Assignment 9 (assigned 2020-10-28, due 2020-11-06).

- Let $\mathcal{M}$ be the set of all finite signed measures on (X, Σ) . Show that $\mathcal{M}$ is a Banach space under the total variation norm.
- For a signed measure μ , we define $\int_X f d\mu = \int_X f d\mu^+ - \int_X f d\mu^-$. Suppose $(f_n) \rightarrow f$, $(g_n) \rightarrow g$, and $|f_n| \leq g_n$ almost everywhere with respect to $|\mu|$. If $\lim \int_X g_n d|\mu| = \int_X g d|\mu| < \infty$, show that $\lim \int_X f_n d\mu = \int_X f d\mu$.
 - Suppose $f, f_n \in L^1$, and $(f_n) \rightarrow f$ almost everywhere. Show that

$$\lim \int |f_n - f| d|\mu| = 0 \iff \lim \int |f_n| d|\mu| = \int |f| d|\mu|.$$

- Let μ, ν be two positive measures.

- If $\nu(X) < \infty$ show that

$$\nu \ll \mu \iff \forall \varepsilon > 0, \exists \delta > 0 \text{ such that } \mu(A) < \delta \implies \nu(A) < \varepsilon.$$

[NOTE: μ is not given to be σ -finite, so you can't apply the Radon-Nikodym theorem.]

- What happens without the assumption $\nu(X) < \infty$?
- Let ν_1 and ν_2 be two finite signed measures on X . Show that there exists a finite signed measure $\nu_1 \vee \nu_2$ such that $\nu_1 \vee \nu_2(A) \geq \nu_1(A) \vee \nu_2(A)$, and for any other finite signed measure ν such that $\nu(A) \geq \nu_1(A) \vee \nu_2(A)$ we must have $\nu_1 \vee \nu_2 \leq \nu$.
 - If ν_1, ν_2 above are absolutely continuous with respect to a positive σ -finite measure μ , prove $\nu_1 \vee \nu_2 \ll \mu$ and express $\frac{d(\nu_1 \vee \nu_2)}{d\mu}$ in terms of $\frac{d\nu_1}{d\mu}$ and $\frac{d\nu_2}{d\mu}$.
 - Let $(\Omega, \mathcal{F}, \mathbf{P})$ be a measure space with $\mathbf{P}(\Omega) = 1$, and $X \in L^1(\Omega, \mathcal{F}, \mathbf{P})$.
[The probabilistic interpretation is that Ω is the sample space, $A \in \mathcal{F}$ is an event, X is a random variable, and for any $B \in \mathcal{B}$, $\mathbf{P}(X \in B)$ is the chance that $X \in B$.]
 - Suppose $\mathcal{G} \subset \mathcal{F}$ is a σ -sub-algebra of $\mathcal{F}$. Show that there exists a unique $\mathcal{G}$ -measurable function, denoted by $\mathbf{E}(X|\mathcal{G})$ such that $\int_A \mathbf{E}(X|\mathcal{G}) d\mathbf{P} = \int_A X d\mathbf{P}$ for all $A \in \mathcal{G}$. [The function $\mathbf{E}(X|\mathcal{G})$ is called the *conditional expectation* of X given $\mathcal{G}$.]
 - If Y is $\mathcal{G}$ measurable, show that $\mathbf{E}(YX|\mathcal{G}) = Y\mathbf{E}(X|\mathcal{G})$.
 - (*Tower property*) If $\mathcal{H} \subset \mathcal{G}$ is a σ -sub-algebra, show that $\mathbf{E}(X|\mathcal{H}) = \mathbf{E}(\mathbf{E}(X|\mathcal{G})|\mathcal{H})$ almost everywhere.
 - (*Conditional Jensen*) If $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is convex and $\varphi(X) \in L^1$, show that $\varphi(\mathbf{E}(X|\mathcal{G})) \leq \mathbf{E}(\varphi(X)|\mathcal{G})$ almost everywhere.
 - Suppose $X \in L^2(\Omega, \mathcal{F}, \mathbf{P})$. Show that $\mathbf{E}(X|\mathcal{G})$ is the L^2 -orthogonal projection of X onto the subspace $L^2(\Omega, \mathcal{G})$. That is, show $\mathbf{E}(X|\mathcal{G}) \in L^2(\Omega, \mathcal{G})$, and $\int_\Omega (X - \mathbf{E}(X|\mathcal{G}))Y d\mathbf{P} = 0$ for all $Y \in L^2(\Omega, \mathcal{G})$.

Optional problems, and details I omitted in class.

- Show that the Radon-Nikodym theorem need not hold if μ, ν are not σ -finite.
- Let $\mathcal{M}$ be the space of all finite signed measures on (X, Σ) . Show that $\mathcal{M}$ with total variation norm (i.e. with $\|\mu\| = |\mu|(X)$) is a Banach space.

Assignment 10 (assigned 2020-11-04, due 2020-11-11).

- If X and Y are not σ -finite, show that Fubini's theorem need not hold.
 - If $\int_{[-1,1]^2} f d\lambda$ is not assumed to exist (in the extended sense), show that both iterated integrals can exist, be finite, but need not be equal.
- (*Fubini for completions.*) Suppose (X, Σ, μ) and (Y, τ, ν) are two σ -finite, complete measure spaces. Let $\varpi = (\Sigma \otimes \tau)_\pi$ denote the completion of $\Sigma \otimes \tau$ with respect to the product measure $\pi = \mu \times \nu$.
 - Show that $\Sigma \otimes \tau$ need not be π -complete (i.e. $\varpi \supsetneq \Sigma \otimes \tau$ in general).
 - Suppose $f : X \times Y \rightarrow [-\infty, \infty]$ is ϖ -measurable. Show that for μ -almost all $x \in X$, the function $y \mapsto f(x, y)$ is τ -measurable, and for ν -almost all $y \in Y$, the function $x \mapsto f(x, y)$ is Σ -measurable.
 - Suppose f is integrable on $X \times Y$ in the extended sense. Define $F(x) = \int_Y f(x, y) d\nu(y)$ and $G(y) = \int_X f(x, y) d\mu(x)$. Show F is defined μ -a.e. and Σ -measurable. Similarly show G is defined ν -a.e., and τ -measurable. Further, show and that $\int_X F d\mu = \int_Y G d\nu = \int_{X \times Y} f d\pi$.
- Compute $\int_0^\infty \frac{\sin x}{x} dx \stackrel{\text{def}}{=} \lim_{R \rightarrow \infty} \int_0^R \frac{\sin x}{x} dx$. [HINT: Substitute $\frac{1}{x} = \int_0^\infty e^{-xy} dy$.]
- If $\frac{1}{p} + \frac{1}{q} = 1$, $f \in L^p$, $g \in L^q$ show that $f * g$ is bounded and continuous. If $p, q < \infty$, show further $f * g(x) \rightarrow 0$ as $|x| \rightarrow \infty$.
- Let $\{\varphi_n\}$ be an approximate identity.
 - If $f \in C_c(\mathbb{R}^d)$, show $f * \varphi_n \rightarrow f$ uniformly.
 - If $\varphi_n \in C_c^\infty(\mathbb{R}^d)$, $p \in [1, \infty]$, and $f \in L^p(\mathbb{R}^d)$ then show that $f * \varphi_n \in C^\infty(\mathbb{R}^d)$.
- Let $A, B \in \mathcal{L}(\mathbb{R})$ be measurable, and define $A + B = \{a + b \mid a \in A, b \in B\}$. If $\lambda(A) > 0$ and $\lambda(B) > 0$ show $A + B$ contains an interval.

Optional problems, and details I omitted in class.

- To see that our characterization of the dual of L^p fails for $p = \infty$, we can (partially) construct a counter example as follows. The Hahn-Banach theorem shows that there exists $T \in (\ell^\infty)^*$ such that $Ta = \lim a_n$, for all $a = (a_n) \in \ell^\infty$ such that $\lim a_n$ exists and is finite. Show that there does not exist $b \in \ell^1$ such that $Ta = \sum a_n b_n$ for all $a \in \ell^\infty$.
- Let μ be a σ -finite measure on X , $p, q \in [1, \infty]$ with $\frac{1}{p} + \frac{1}{q} = 1$, and $g \in L^1$. If for every $f \in L^p$ we know $fg \in L^1$ and further $\sup\{\|\int fg\| \mid \|f\|_p \leq 1\} = C < \infty$, then show that $g \in L^q$ and $\|g\|_q = C$. [I used this to finish the proof of the Radon-Nikodym theorem in class.]
- Show that the Lebesgue measure on $\mathbb{R}^{m+n}$ is the product of the Lebesgue measures on $\mathbb{R}^m$ and $\mathbb{R}^n$ respectively.

Assignment 11 (assigned 2020-11-11, due 2020-11-18).

1. Let $\alpha \in (0, 1)$ and $f \in C_{per}^\alpha$. Show that $(S_N f) \rightarrow f$ uniformly, as $N \rightarrow \infty$.
2. If $\alpha \in (0, 1)$, $f \in C_{per}^\alpha([0, 1])$, show that $\sup_{n \in \mathbb{N}} |n|^\alpha |\hat{f}(n)| < \infty$.
3. Let μ be a finite signed Borel measure on $[0, 1]$. If $\forall n \in \mathbb{Z} \hat{\mu}(n) = 0$, show $\mu = 0$.
4. Let $f \in L^2([0, 1])$. Show that there exists a unique $u \in C_{per}^\infty(\mathbb{R} \times (0, \infty))$ such that $\lim_{t \rightarrow 0^+} \|u(\cdot, t) - f(\cdot)\|_{L_{per}^2} = 0$, and $\partial_t u - \partial_x^2 u = 0$. [You may assume the result of the optional problems.]
5. Let $0 \leq r < s$. Show that any bounded sequence in H_{per}^s has a subsequence that is convergent in H_{per}^r .

Optional problems, and details I omitted in class.

- * If $f \in L^p$, $g \in L^q$ with $p, q \in [1, \infty]$ and $1/p + 1/q \geq 1$, show that $f * g = g * f$.
- * If $f \in L^p$, $g \in L^q$, $h \in L^r$ with $p, q, r \in [1, \infty]$ and $1/p + 1/q + 1/r \geq 2$, show that $(f * g) * h = f * (g * h)$.
- * Define $e_n(x) = e^{2\pi i n x}$, and define the Dirichlet and Fejér kernels by $D_N = \sum_{-N}^N e_n$, $F_N = \frac{1}{N} \sum_{0}^{N-1} D_n$.
 - (a) Show that $D_N(x) = \frac{\sin((2N+1)\pi x)}{\sin(\pi x)}$. Further show $\lim_{N \rightarrow \infty} \int_\varepsilon^{1-\varepsilon} |D_N| = \infty$.
 - (b) Show that $F_N(x) = \frac{\sin^2(N\pi x)}{N \sin^2(\pi x)}$, and that $\{F_N\}$ is an approximate identity.
- * (a) If $f, g \in L_{per}^2([0, 1])$, show that $(f * g)^\wedge(n) = \hat{f}(n) \hat{g}(n)$.
 - (b) If $f, g \in L_{per}^2([0, 1])$, show that $(fg)^\wedge(n) = \hat{f} * \hat{g}(n) \stackrel{\text{def}}{=} \sum_{m \in \mathbb{Z}} \hat{f}(m) \hat{g}(n - m)$.
- * For any $s \geq 0$ show that H_{per}^s is a closed subspace of L^2 .
- * Show that $f \in H_{per}^s$ for all $s \geq 0 \iff f \in C_{per}^\infty$.
- * Let $n \in \mathbb{N} \cup \{0\}$, $\alpha \in [0, 1)$ $s > 1/2 + n + \alpha$. Show that $H_{per}^s \subseteq C_{per}^{n, \alpha}[0, 1]$ and the inclusion map is continuous. [Recall $C_{per}^{n, \alpha}[0, 1]$ is the set of all C^n periodic functions on $\mathbb{R}$ whose n^{th} derivative is Hölder continuous with exponent α .]
- * Find a function $f \in H_{per}^{1/2} - L^\infty$. [Thus, the Sobolev embedding theorem is false for $s = 1/2$.]
- * Find an example of f such that $(|n|^\alpha \hat{f}(n)) \rightarrow 0$, but $f \notin C_{per}^\alpha$.

Assignment 12 (assigned 2020-11-18, due 2020-12-02).

1. Let $s \in (0, 1]$ and $f \in L_{per}^2$. Show that $f \in H_{per}^s$ if and only if

$$\int_0^1 \left(\frac{\|f - \tau_h f\|_{L^2}}{h^s} \right)^2 \frac{dh}{h} < \infty \quad \text{for } s < 1,$$

and

$$\sup_{|h| \leq 1} \frac{\|f - \tau_h f\|_{L^2}}{h} < \infty \quad \text{for } s = 1.$$

2. (a) Let $n \in \mathbb{N}$ be even, $\frac{1}{n} + \frac{1}{n'} = 1$. If $\hat{f} \in \ell^{n'}(\mathbb{Z})$, show that $f \in L_{per}^n([0, 1])$ and $\|f\|_{L^n} \leq \|\hat{f}\|_{\ell^{n'}}$. [HINT: Let $n = 2m$. Then $\|f\|_{L^n}^n = \|(f^m)^\wedge\|_{\ell^2}^2$.]
 - (b) Let $s > \frac{1}{2} - \frac{1}{p} \geq 0$, and $\frac{1}{p} + \frac{1}{q} = 1$. If $f \in H_{per}^s$ show $\hat{f} \in \ell^q(\mathbb{Z})$. Further show that the map $f \mapsto \hat{f}$ is continuous from $H_{per}^s \rightarrow \ell^q$.
 - (c) If $n \in \mathbb{N}$ is even, $s > \frac{1}{2} - \frac{1}{n}$ then show that $H_{per}^s \subset L^n([0, 1])$ and that the inclusion map is continuous. [This is one of the Sobolev embedding theorems.]
3. Let $s > 3/2$ and $f, g \in H_{per}^s$. Show that $fg \in H^1$, and further $D(fg) = (Df)g + f(Dg)$. (Here Df is the weak derivative of f .)
4. (a) If $f \in L^1(\mathbb{R}^d)$ and f is not identically 0 (a.e.), then show that $Mf \notin L^1(\mathbb{R}^d)$. The next few subparts outline a proof that for any $p > 1$, the maximal function is an L^p bounded sublinear operator. Let $p \in (1, \infty)$, $f \in L^p(\mathbb{R}^d)$ and $f \geq 0$.
 - (b) Show that $\lambda\{Mf > \alpha\} \leq \frac{3^d}{(1-\delta)\alpha} \int_{\{f > \delta\alpha\}} f$, for any $t > 0$, $\delta \in (0, 1)$ and $f \geq 0$ measurable.
 - (c) Let $p \in (1, \infty]$, and $d \in \mathbb{N}$. Show that there exists a constant $c = c(p, d)$ such that $\|Mf\|_p \leq c\|f\|_p$ for all $f \in L^p(\mathbb{R}^d)$. [HINT: For $p < \infty$, use the previous part, the identity $\|Mf\|_p^p = \int_0^\infty p\alpha^{p-1} \lambda\{Mf > \alpha\} d\alpha$ and optimise in δ .]
5. (a) Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a right continuous increasing function. Show that there exists a finite Borel measure μ such that $\mu((x, y]) = f(y) - f(x)$ for every $x, y \in [a, b]$. Show further that $\mu = \mu_{ac} + \mu_{sc} + \sum_i \alpha_i \delta_{a_i}$, where $\mu_{ac} \ll \lambda$, $\alpha_i > 0$, $a_i \in [a, b]$, $\sum_i \alpha_i < \infty$, and $\mu_{sc} \perp \lambda$ is such that $\mu_{sc}(\{x\}) = 0$ for all $x \in \mathbb{R}$.
 - (b) Let $f : [a, b] \rightarrow \mathbb{R}$ be monotone. Show that f is differentiable almost everywhere, $f' \in L^1([a, b])$ and that $|\int_a^b f'| \leq |f(b) - f(a)|$.

Optional problems, and details I omitted in class.

- * Show that $f : [a, b] \rightarrow \mathbb{R}$ has bounded variation if and only if it is the difference of two increasing functions.
- * Let $s \geq 1$, and $f \in H_{per}^s$. Show that f has a weak derivative Df , and $Df \in H^{s-1}$. Further, show that the map $f \mapsto Df : H^s \rightarrow H^{s-1}$ is linear and continuous.
- * (*Infinite version of Vitali.*) Suppose $A \subset \cup B_\alpha$, where $\{B_\alpha\}_{\alpha \in \mathcal{A}}$ is an infinite collection of balls such that $\sup \lambda(B_\alpha) < \infty$. Show that there exists $\mathcal{A}' \subset \mathcal{A}$ such that the sub-collection $\{B_{\alpha'}\}_{\alpha' \in \mathcal{A}'}$ is disjoint and $A \subset \cup 5B_{\alpha'}$.
- * If $f \in L^1(\mathbb{R}^d)$, show that $Mf(x) \geq |f(x)|$ at all Lebesgue points of f .

Assignment 13 (assigned 2020-12-02, due 2020-12-09).

1. Let μ be a positive finite Borel measure on $\mathbb{R}^d$, and $\alpha > 0$. Show that for every $A \subset \{D\mu > \alpha\}$, we must have $\mu(A) \geq \alpha\lambda(A)$.
2. Let $f \in L^1(\mathbb{R}^d)$. Let $S^{d-1} = \{y \in \mathbb{R}^d \mid |y| = 1\}$ be the $d-1$ dimensional sphere of radius 1. Show that there exists a unique measure σ on S^{d-1} such that

$$\int_{\mathbb{R}^d} f(x) dx = \int_{r \in [0, \infty)} \int_{y \in S_1} f(ry) r^{d-1} d\sigma(y) d\lambda(r).$$

HINT: For $A \in \mathcal{B}(S^{d-1})$ define $\sigma(A) = \int \lambda(A^*)$ where $A^* = \{rx \mid x \in A, r \in [0, 1]\}$. Now for any $B \in \mathcal{B}(S^{d-1})$ prove the desired equality when $f = \mathbf{1}_A$ where $A = \{rx \mid a < r < b, x \in B\}$.

3. Let $\alpha \in [0, d]$, and $A \in \mathcal{B}(\mathbb{R}^d)$. If $H_\alpha(A) < \infty$, show $\lim_{r \rightarrow 0} \frac{H_\alpha(A \cap B(x, r))}{r^\alpha} = 0$ for H_α -almost all $x \notin A$.
4. If μ is a finite Borel measure on $\mathbb{R}^d$ define $\hat{\mu}(\xi) = \int e^{-2\pi i \langle x, \xi \rangle} d\mu(x)$. If $\hat{\mu}(\xi) = 0$ for all ξ , show that $\mu = 0$. [HINT: Show that $\int f d\mu = 0$ for all $f \in \mathcal{S}$.]
5. (*Lévy's continuity theorem*) Let μ, μ_n be a probability measures on $\mathbb{R}^d$ such that $(\hat{\mu}_n(\xi)) \rightarrow \hat{\mu}(\xi)$ for every $\xi \in \mathbb{R}^d$. Show that $\int_{\mathbb{R}^d} f d\mu_n \rightarrow \int_{\mathbb{R}^d} f d\mu$ for every bounded continuous function f .
HINT: First prove the theorem for $f \in \mathcal{S}$. Next show $\mu_n(|x| > 1/\lambda) \leq \frac{C}{\lambda} \int_{[-\lambda, \lambda]^d} (1 - \hat{\mu}_n(\xi)) d\xi$, and use this to prove the theorem for all bounded continuous f .
6. (*Central limit theorem*) Let $g \in L^1(\mathbb{R})$ be such that $g \geq 0$, $\int_{\mathbb{R}} g dx = 1$, $\int_{\mathbb{R}} xg(x) = 0$ and $\int_{\mathbb{R}} x^2 g(x) dx = \sigma^2 < \infty$. Define $g^{*n} = (g * \dots * g)$ (n -times), and $h_n(x) = \delta_{1/\sqrt{n}} g^{*n}(x) = \sqrt{n} g^{*n}(\sqrt{n}x)$. Show

$$\hat{h}_n(\xi) \xrightarrow{n \rightarrow \infty} \exp(-2\pi^2 \sigma^2 \xi^2),$$

and $\int_{\mathbb{R}} f h_n dx \rightarrow \frac{1}{\sqrt{2\pi\sigma^2}} \int_{\mathbb{R}} f(x) \exp\left(-\frac{x^2}{2\sigma^2}\right) dx$

for every bounded continuous function f .

Optional problems, and details I omitted in class.

- * Let μ be a finite signed Borel measure on $\mathbb{R}^d$ such that $\mu \perp \lambda$. Show that $D|\mu| = \infty$, μ -almost everywhere, and $D|\mu| = 0$, λ -almost everywhere.
- * Let $c_\alpha = \frac{\pi^{\alpha/2}}{\Gamma(1+\frac{\alpha}{2})}$ be the normalization constant from the definition of H_α , the Hausdorff measure of dimension α .
 - (a) If $0 < H_\alpha(A) < \infty$, show $\limsup_{r \rightarrow 0} \frac{H_\alpha(A \cap B(x, r))}{c_\alpha r^\alpha} \in [2^{-\alpha}, 1]$ for H_α -a.e. $x \in A$.
 - (b) Show that there exists $\alpha < d$ and $A \subset \mathbb{R}^d$ with $H_\alpha(A) \in (0, \infty)$ such that

$$\liminf_{r \rightarrow 0} \frac{H_\alpha(A \cap B(x, r))}{c_\alpha r^\alpha} = 0 \quad \text{and} \quad \limsup_{r \rightarrow 0} \frac{H_\alpha(A \cap B(x, r))}{c_\alpha r^\alpha} < 1,$$

for H^α -almost every $x \in \mathbb{R}^d$.

(c) If C is the Cantor set, and $\alpha = \log 2 / \log 3$, compute $\limsup_{r \rightarrow 0} \frac{H_\alpha(C \cap B(x, r))}{c_\alpha r^\alpha}$.

- * We say the family $\{E_r\}$ *shrinks nicely* to $x \in \mathbb{R}^d$ if there exists $\delta > 0$ such that for all r , $E_r \subset B(x, r)$ and $\lambda(E_r) > \delta \lambda(B(x, r))$. If $\{E_r\}$ shrinks nicely to x , show that $\lim_{r \rightarrow 0} \frac{1}{\lambda(E_r)} \int_{E_r} f = f(x)$ for all Lebesgue points of f .
- * If $f \in L^1(\mathbb{R}^d)$, show that $Mf(x) \geq |f(x)|$ at all Lebesgue points of f .
- * (a) If $p \in [1, \infty)$, $f \in L^p(\mathbb{R}^d)$, $g \in \mathcal{S}(\mathbb{R}^d)$, show that $f * g \in C^\infty(\mathbb{R}^d)$, and further $D^\alpha(f * g) = f * (D^\alpha g)$ for every multi-index α .
(b) For $p \in [1, \infty)$, show that C_c^∞ and $\mathcal{S}$ are dense subsets of L^p .
- * Show that the mapping $f \mapsto \hat{f}$ is a bijection in the Schwartz space.
- * For $f \in L^1$, the formula $\hat{f}(\xi) = \int f(x) e^{-2\pi i \langle x, \xi \rangle}$ allows us to prove many identities: E.g. $(\delta_\lambda f)^\wedge(\xi) = \hat{f}(\lambda\xi)$, etc. For $f \in L^2$, the formula $\hat{f}(\xi) = \int f(x) e^{-2\pi i \langle x, \xi \rangle}$ is no longer valid, as the integral is not defined (in the Lebesgue sense). However, most identities remain valid, and can be proved using an approximation argument. I list a couple here.
 - (a) For $f \in L^1$ we know $(\tau_x f)^\wedge(\xi) = e^{-2\pi i \langle x, \xi \rangle} \hat{f}(\xi)$. Prove it for $f \in L^2$.
 - (b) Similarly, show that $(\delta_\lambda f)^\wedge(\xi) = \hat{f}(\lambda\xi)$ for all $f \in L^2$.
 - (c) Let $\mathcal{F}$ denote the Fourier transform operator (i.e. $\mathcal{F}f = \hat{f}$), and R denote the reflection operator (i.e. $Rf(x) = f(-x)$). Note that our Fourier inversion formula (for $f \in L^1$, $\hat{f} \in L^1$) is exactly equivalent to saying $\mathcal{F}^2 f = Rf$. Prove $\mathcal{F}^2 f = Rf$ for all $f \in L^2$.

Assignment 14 (assigned 2020-12-09, due never).

1. (*Uncertainty principle*) Suppose $f \in \mathcal{S}(\mathbb{R})$. Show that

$$\left(\int_{\mathbb{R}} |xf(x)|^2 dx \right) \left(\int_{\mathbb{R}} |\xi \hat{f}(\xi)|^2 d\xi \right) \geq \frac{1}{16\pi^2} \|f\|_{L^2}^2 \|\hat{f}\|_{L^2}^2$$

[This illustrates a nice localisation principle about the Fourier transform. The first integral measures the spread of the function f . The second the spread of the Fourier transform $\hat{f}$. Here you show that this product is bounded below! The proof, once you know enough Physics, reduces to the above inequality.

Hint: Consider $\int_{\mathbb{R}} xf(x)f'(x) dx$.]

2. (*Trace theorems*) Let $p \in \mathbb{R}^m$ be fixed. Given $f : \mathbb{R}^{m+n} \rightarrow \mathbb{R}$ define $S_p f : \mathbb{R}^n \rightarrow \mathbb{R}$ by $S_p f(y) = f(p, y)$.

- (a) Let $s > m/2$, and $s' = s - m/2$. Show that there exists a constant c such that $\|S_p f\|_{H^{s'}(\mathbb{R}^n)} \leq c \|f\|_{H^s(\mathbb{R}^{m+n})}$.

- (b) Show that the section operator S_p extends to a continuous linear operator from $H^s(\mathbb{R}^{m+n})$ to $H^{s'}(\mathbb{R}^n)$.

[Given an arbitrary L^2 function on $\mathbb{R}^{m+n}$ it is of course impossible to restrict it to an m -dimensional hyper-plane. However, if your function has more than $n/2$ “Sobolev derivatives”, then you can make sense of this restriction, and the restriction still has $s - n/2$ “Sobolev derivatives”.]

3. Find $E \in \mathcal{L}(\mathbb{R}^d)$ and $x \in \mathbb{R}^d$ such that $\lim_{r \rightarrow 0} \frac{\lambda(E \cap B(x, r))}{\lambda(B(x, r))}$ does not exist.
4. Let μ be a finite Borel measure on $\mathbb{R}^d$ such that $\mu(\{x\}) = 0$ for all $x \in \mathbb{R}^d$. True or false: For any $\alpha \in [0, \mu(\mathbb{R}^d)]$ there exists $A \in \mathcal{B}(\mathbb{R}^d)$ such that $\mu(A) = \alpha$. Prove it, or find a counter example.
5. Show that the arbitrary union of closed (non-degenerate) cubes (with sides parallel to the coordinate axis) is Lebesgue measurable. [HINT: Look up and use the Vitali covering theorem (which is stronger than the covering lemma I used). More generally one can show that the arbitrary union of convex sets with nonempty interiors is Lebesgue measurable (see Balcerzak and Kharazishvili '99).]
6. Let $p \in [1, 2]$, $f \in L^p(\mathbb{R}^d)$. Given $R > 0$ define $\hat{f}_R = (\mathbf{1}_{B(0, R)} f)^\wedge$. Is there a sequence $(R_n) \rightarrow \infty$ such that $\hat{f}_{R_n}$ converges almost everywhere? Prove it, or find a counter example.
7. Let $B = B(0, 1) \subseteq \mathbb{R}^d$. Show that $|B| = \pi^{d/2} / \Gamma(1 + n/2)$, where $\Gamma(s) = \int_0^\infty x^{s-1} e^{-x} dx$. [HINT: Compute $\int_{\mathbb{R}^d} e^{-|x|^2/2} dx$.]
8. I may add to this list in the next few days.