

Putnam E.8

Po-Shen Loh

22 October 2024

1 Problems

Putnam 1987/A1. Curves A , B , C , and D are defined in the plane as follows:¹

$$\begin{aligned}A &= \left\{ (x, y) : x^2 - y^2 = \frac{x}{x^2 + y^2} \right\}, \\B &= \left\{ (x, y) : 2xy + \frac{y}{x^2 + y^2} = 3 \right\}, \\C &= \left\{ (x, y) : x^3 - 3xy^2 + 3y = 1 \right\}, \\D &= \left\{ (x, y) : 3x^2y - 3x - y^3 = 0 \right\}.\end{aligned}$$

Prove that $A \cap B = C \cap D$.

Putnam 1987/A2. The sequence of digits

123456789101112131415161718192021...

is obtained by writing the positive integers in order. If the 10^n -th digit in this sequence occurs in the part of the sequence in which the m -digit numbers are placed, define $f(n)$ to be m . For example, $f(2) = 2$ because the 100th digit enters the sequence in the placement of the two-digit integer 55. Find, with proof, $f(1987)$.

Putnam 1987/A3. For all real x , the real-valued function $y = f(x)$ satisfies

$$y'' - 2y' + y = 2e^x.$$

- (a) If $f(x) > 0$ for all real x , must $f'(x) > 0$ for all real x ? Explain.
- (b) If $f'(x) > 0$ for all real x , must $f(x) > 0$ for all real x ? Explain.

¹The equations defining A and B are indeterminate at $(0, 0)$. The point $(0, 0)$ belongs to neither.