

Fluctuations of the solutions to the KPZ equation in dimensions three and higher

Alexander Dunlap, Yu Gu, Lenya Ryzhik, Ofer Zeitouni

Abstract

We prove, using probabilistic techniques and analysis on the Wiener space, that the large scale fluctuations of the KPZ equation in $d \geq 3$ with a small coupling constant, driven by a white in time and colored in space noise, are given by the Edwards-Wilkinson model. This gives an alternative proof, that avoids perturbation expansions, to the results of Magnen and Unterberger [27].

1 Introduction

1.1 The main result

We consider the random heat equation

$$\partial_t u = \frac{1}{2} \Delta u + \beta V(t, x) u, \quad u(0, x) \equiv 1, \quad x \in \mathbb{R}^d, d \geq 3, \quad (1.1)$$

with a coupling constant $\beta > 0$, and a random process $V(t, x)$ that is white in time and colored in space, constructed from a space-time white noise $\dot{W}(t, x)$ defined on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$:

$$V(t, x) = \int_{\mathbb{R}^d} \varphi(x - y) \dot{W}(t, y) dy.$$

Here, the non-negative mollifier $\varphi \geq 0$ is in $\mathcal{C}_c^\infty(\mathbb{R}^d)$, and the product $V(t, x)u$ in (1.1) is interpreted in the Itô sense. The solution of (1.1) is then a continuous random field. We denote by $R(x)$ the spatial covariance function of $V(t, x)$:

$$R(x) = \int_{\mathbb{R}^d} \varphi(x + y) \varphi(y) dy, \quad (1.2)$$

so we can formally write

$$\mathbb{E}[V(t, x)V(s, y)] = \delta(t - s)R(x - y).$$

Note that $R \in \mathcal{C}_c^\infty(\mathbb{R}^d)$, and without additional loss of generality, we assume that $R(x) = 0$ for $|x| \geq 1$.

Consider the macroscopically rescaled solution

$$u_\varepsilon(t, x) = u\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right),$$

which solves

$$\partial_t u_\varepsilon = \frac{1}{2} \Delta u_\varepsilon + \frac{\beta}{\varepsilon^2} V\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right) u_\varepsilon.$$

The goal of this paper is to identify the asymptotic fluctuations of

$$h_\varepsilon(t, x) = \frac{1}{\beta\varepsilon^{(d-2)/2}} \log u_\varepsilon(t, x) \quad (1.3)$$

as a random distribution, that is, the asymptotic distribution, as $\varepsilon \rightarrow 0$, of

$$\int_{\mathbb{R}^d} h_\varepsilon(t, x) g(x) dx \quad (1.4)$$

for any test function $g \in \mathcal{C}_c^\infty(\mathbb{R}^d)$. Note that by the scaling property of the space-time white noise, we have

$$\frac{1}{\varepsilon^2} V\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right) \stackrel{\text{law}}{=} \varepsilon^{\frac{d-2}{2}} \dot{W}_\varepsilon(t, x), \quad \text{with} \quad \dot{W}_\varepsilon(t, x) = \frac{1}{\varepsilon^d} \int_{\mathbb{R}^d} \varphi\left(\frac{x-y}{\varepsilon}\right) \dot{W}(t, y) dy. \quad (1.5)$$

By (1.5) and the Itô formula, it is clear that $h_\varepsilon - \mathbb{E}[h_\varepsilon] \stackrel{\text{law}}{=} \tilde{h}_\varepsilon - \mathbb{E}[\tilde{h}_\varepsilon]$ with

$$\partial_t \tilde{h}_\varepsilon = \frac{1}{2} \Delta \tilde{h}_\varepsilon + \frac{1}{2} \beta \varepsilon^{\frac{d-2}{2}} |\nabla \tilde{h}_\varepsilon|^2 + \dot{W}_\varepsilon(t, x), \quad \tilde{h}_\varepsilon(0, x) \equiv 0. \quad (1.6)$$

Here is the main result of paper.

Theorem 1.1. *There exists $\beta_0 = \beta_0(d, \varphi)$ so that for $\beta < \beta_0$, $t > 0$ and test function $g \in \mathcal{C}_c^\infty(\mathbb{R}^d)$, we have*

$$\int_{\mathbb{R}^d} (h_\varepsilon(t, x) - \mathbb{E}[h_\varepsilon(t, x)]) g(x) dx \Rightarrow \int_{\mathbb{R}^d} \mathcal{U}(t, x) g(x) dx \quad (1.7)$$

in distribution as $\varepsilon \rightarrow 0$, where \mathcal{U} solves the Edwards-Wilkinson equation

$$\partial_t \mathcal{U} = \frac{1}{2} \Delta \mathcal{U} + \nu_{\text{eff}} \dot{W}(t, x), \quad \mathcal{U}(0, x) \equiv 0, \quad (1.8)$$

with the effective variance

$$\nu_{\text{eff}}^2 = \int_{\mathbb{R}^d} R(x) \mathbb{E}_B \left[\exp \left(\frac{1}{2} \beta^2 \int_0^\infty R(x + B_s) ds \right) \right] dx. \quad (1.9)$$

Theorem 1.1 is a particular case of the following more general result.

Theorem 1.2. *Assume that a function $f \in \mathcal{C}^2(\mathbb{R}_+)$ satisfies*

$$|f(y)| + |f'(y)| + |f''(y)| \leq M(y^p + y^{-p}), \quad y \in \mathbb{R}_+$$

for some $M, p > 0$. Then, there exists $\beta_0 = \beta_0(d, \varphi, p, M)$ so that, for $\beta < \beta_0$, $t > 0$ and test function $g \in \mathcal{C}_c^\infty(\mathbb{R}^d)$, we have

$$\frac{1}{\beta\varepsilon^{(d-2)/2}} \int_{\mathbb{R}^d} (f(u_\varepsilon(t, x)) - \mathbb{E}[f(u_\varepsilon(t, x))]) g(x) dx \Rightarrow \sigma_f \int_{\mathbb{R}^d} \mathcal{U}(t, x) g(x) dx \quad (1.10)$$

in distribution as $\varepsilon \rightarrow 0$, where \mathcal{U} is a solution to (1.8) and $\sigma_f = \mathbb{E}[f'(Z_\infty)Z_\infty]$. Here, Z_∞ is the positive random variable defined in (2.4) below with $\mathbb{E}[Z_\infty] = 1$.

Throughout the paper, we assume that $\beta \in (0, \beta_0)$ as in the statements of Theorems 1.1 and 1.2.

1.2 The context

The study of the KPZ equation has witnessed important progress in recent years. A lot of work was done in $d = 1$, including making sense of the equation without relying on the Hopf-Cole transform [20, 21, 22, 23, 26], proving the weak/strong universality conjecture in the one-dimensional KPZ universality class [1, 2, 3, 28], etc. We refer to the reviews [15, 35] for a more complete list of references. In $d = 2$, some relevant results can be found in [4, 7, 8, 9, 10, 17, 37].

Our result in $d \geq 3$ can be viewed as a continuation of the previous works on the stochastic heat equation (SHE) [16, 18] and as a counterpart of the recent work of Magnen and Unterberger [27], where the driving force is mollified in both temporal and spatial variables. While the proof in [27] is based on a multiscale expansion and a calculation of multi-point correlation functions, we present a probabilistic proof using the tools of Malliavin calculus.

It is well-known that in $d \geq 3$ there is a phase transition as a function of the coupling constant β , also known as the inverse temperature if we view the solution to (1.1) as the partition function of a directed polymer in a random environment, and the behaviors of the solution to the equation and the underlying polymers change drastically for different values of β . There are different notions of critical temperatures in $d \geq 3$ [13]. For our analysis, what is particularly important is that we stay deep in the weak-disorder/high-temperature regime where β is small and the $L^2(\Omega)$ norm of $u(t, x)$ is bounded uniformly in $t > 0$. For directed polymers, the diffusive behavior was proved in the early work of [6, 25], and we refer to [13] for a review of further developments and [31, 32] for the connection to the stochastic heat equation.

1.3 Connection to the stochastic heat equation

We note that the coefficient in front of the nonlinear term in (1.6) is small when $d \geq 3$, and if we naively ignore the nonlinear term, the limiting equation for \tilde{h}_ε would be

$$\partial_t \bar{h} = \frac{1}{2} \Delta \bar{h} + \nu \dot{W}(t, x),$$

with

$$\nu^2 = \int_{\mathbb{R}^d} R(x) dx < \nu_{\text{eff}}^2,$$

since $\dot{W}_\varepsilon(t, x) \rightarrow \nu \dot{W}(t, x)$ as $\varepsilon \rightarrow 0$. Thus, Theorem 1.1 shows that the “optically small” nonlinear term affects the effective variance asymptotically, even in the limit $\varepsilon \rightarrow 0$, when β is small but does not change fundamentally the nature of the Edwards-Wilkinson limit. Let us explain informally why this happens. As we have shown in [16], when β is small, the solution of (1.1), before any space-time rescaling, behaves at large times approximately as a space-time stationary random field $\Psi(t, x)$ with spatial correlations that decay, when the potential is white in time, as

$$\text{Cov}(\Psi(0, 0), \Psi(0, y)) \sim \frac{\bar{c} \beta^2 \nu_{\text{eff}}^2}{|y|^{d-2}}, \quad |y| \gg 1, \quad (1.11)$$

with a universal constant \bar{c} . Thus, $h(t, x) = \log u(t, x)$ is approximately $\log \Psi(t, x)$ with a similar decay of correlations. The law of large numbers implies then that $h(t, x)$ converges as a random distribution (after integration against a smooth test function) to a constant. The spatial decay rate of correlations in (1.11) indicates that, as a distribution, $\bar{h}(t, x) = h(t, x) - \mathbb{E}[h(t, x)]$ is of size $\varepsilon^{(d-2)/2}$ when integrated against a test function on the scale ε^{-1} . In other words, $\bar{h}(t, x)$ is close to zero as a random distribution but not point-wise. On a technical note, the slow spatial decay of correlations

of $\Psi(t, x)$ does not allow us to apply the central limit theorem directly to integrals on the macroscopic scale ε^{-1} , thinking of the integral as a sum over ε^{-d} boxes. In a sense, the hard work is to incorporate the fast temporal mixing of $V(t, x)$ into the picture, with the help of the Feynman-Kac formula. Ignoring this serious technical issue, the correlation structure of $\log \Psi(t, x)$ then dictates the rescaling by the factor $\varepsilon^{(d-2)/2}$ in (1.3) that, in turn, shows up as the factor $\varepsilon^{(d-2)/2}$ in front of the nonlinear term in (1.6). However, as $\bar{h}(t, x)$ is not close to a constant pointwise, this term is not “ ε -wise” small in the pointwise sense, only “ β -wise” small, and thus makes a non-trivial contribution to the effective variance, of order $o(1)$ as $\beta \rightarrow 0$, that survives in the limit $\varepsilon \rightarrow 0$. The precise Edwards-Wilkinson nature of the limit comes as a combination of the Gaussianity coming from the central limit theorem, modulo the technical difficulties discussed above, and the heat semi-group.

In [18, Theorem 1.2], it was proved that the fluctuations of u_ε are given by the same Edwards-Wilkinson model:

$$\frac{1}{\beta \varepsilon^{(d-2)/2}} \int_{\mathbb{R}^d} (u_\varepsilon(t, x) - \mathbb{E}[u_\varepsilon(t, x)])g(x) \Rightarrow \int_{\mathbb{R}^d} \mathcal{U}(t, x)g(x)dx, \text{ as } \varepsilon \rightarrow 0, \quad (1.12)$$

which can also be viewed as a special case of Theorem 1.2 with $f(y) = y$. In other words, when viewed as random fields, $u_\varepsilon(t, \cdot)$ and $\log u_\varepsilon(t, \cdot)$ have the same limiting distribution! While it is unclear at first glance why this should be the case, the proof in this paper helps illustrate the connection, see the discussion in Remark 3.8.

Let us make a couple of remarks on Theorem 1.2. First, if we take $f(y) = \log y - (y - 1)$ in Theorem 1.2, then

$$\sigma_f = 1 - \mathbb{E}[Z_\infty] = 0,$$

hence

$$\frac{1}{\varepsilon^{(d-2)/2}} \int_{\mathbb{R}^d} (f(u_\varepsilon(t, x)) - \mathbb{E}[f(u_\varepsilon(t, x))])g(x)dx \rightarrow 0 \quad (1.13)$$

in probability as $\varepsilon \rightarrow 0$. In other words, we have

$$\varepsilon^{-(d-2)/2} \log u_\varepsilon(t, \cdot) \approx \varepsilon^{-(d-2)/2} [u_\varepsilon(t, \cdot) - 1] \quad (1.14)$$

as random distributions (after centering). We stress that (1.14) is not a simple consequence of a linearization of $\log y$ around $y = 1$. For example, for $f(y) = y^2$, we have

$$\sigma_f = 2\mathbb{E}[Z_\infty^2] > 2\mathbb{E}[Z_\infty]^2 = 2,$$

which is different from the coefficient obtained by the linearization $f(y) \approx 2y - 1$ near $y = 1$. Such a linearization fails in general precisely because $u_\varepsilon(t, x)$ is not close to 1 but rather approaches a stationary solution in the long time limit, as discussed above.

Second, if we recall the connection between the effective variance in the Edwards-Wilkinson equation and the decay of correlations rate in (1.11) for $\Psi(t, x)$, and note that a similar connection holds for $f(\Psi(t, x))$, Theorem 1.2 says that if $\Psi(t, x)$ satisfies (1.11), then the correlations of $f(\Psi(t, x))$ decay as

$$\text{Cov}(f(\Psi(0, 0)), f(\Psi(0, y))) \sim \frac{\bar{c}\beta^2 \sigma_f^2 \nu_{\text{eff}}^2}{|y|^{d-2}}, \quad |y| \gg 1, \quad (1.15)$$

with $\sigma_f = \mathbb{E}[f'(\Psi(0, 0))\Psi(0, 0)]$. As explained in Remark 5.5 below, this is consistent with $\log \Psi(t, x)$ being a “Gaussian random field on large scales”.

For a potential that is smooth in both x and t variable, such as

$$V(t, x) = \int_{\mathbb{R}^{d+1}} \phi(t-s)\varphi(x-y)\dot{W}(s, y)dyds$$

for some $\phi \in \mathcal{C}_c^\infty(\mathbb{R})$, a result similar to (1.12) was proved in [18, Theorem 1.1], where the limiting Edwards-Wilkinson equation has also an effective diffusivity:

$$\partial_t \mathcal{U} = \frac{1}{2} \nabla \cdot a_{\text{eff}} \nabla \mathcal{U} + \nu_{\text{eff}} \dot{W}.$$

This is consistent with the result in [27], where the same limit is proved for the KPZ equation. The effective diffusion matrix a_{eff} comes from the temporal correlation of the randomness, which was previously discussed in [5, 19]. In [18, 30], as a crucial ingredient of the proof, a Markov chain was constructed to model the evolution of the path increments and the fast mixing of the Markov chain drives a central limit theorem which gives rise to the effective diffusivity. We believe that the approach developed in this paper, combined with the Markov chain techniques used in [18], will give another proof of the result in [27] for random potentials that are not white in time. We choose to work in the white in time setting since it is technically simpler to explain, but is also illustrative enough to reveal the main idea of the proof for the general case.

All the aforementioned results are on the asymptotics of the random fields after a spatial averaging. For the pointwise fluctuations, we refer to the recent work [12, 14].

1.4 Notation

We use throughout the following notations and conventions.

- (i) We use $a \lesssim b$ for $a \leq Cb$ for some constant C which is independent of ε but may depend on t .
- (ii) We use (p, q) to denote the Hölder exponents $\frac{1}{p} + \frac{1}{q} = 1$, and always choose $p \gg 1$.
- (iii) $G_t(x) = (2\pi t)^{-d/2} \exp(-|x|^2/2t)$ denotes the standard heat kernel.
- (iv) We let H denote the Hilbert space $L^2(\mathbb{R}^{d+1})$, with norm $\|\cdot\|_H$ and inner product $\langle \cdot, \cdot \rangle_H$.
- (v) $\{B_t^j, W_t^j : t \geq 0, j = 1, \dots\}$ is a family of standard independent d -dimensional Brownian motions built on another probability space $(\Sigma, \mathcal{A}, \tilde{\mathbb{P}})$. We will use $\mathbb{E}_B, \mathbb{E}_W, \mathbb{P}_B, \mathbb{P}_W$ when taking the expectation and the probability with respect to B, W separately.
- (vi) We use $d_{\text{TV}}(\cdot, \cdot)$ to denote the total variation distance between two distributions, and if X, Y are random variables of laws μ_X, μ_Y , we write $d_{\text{TV}}(X, Y)$ for $d_{\text{TV}}(\mu_X, \mu_Y)$.
- (vii) We let $\|\cdot\|_{\text{op}}$ denote the operator norm.

Acknowledgments. AD was supported by an NSF Graduate Research Fellowship, YG by NSF grant DMS-1613301/1807748 and the Center for Nonlinear Analysis of CMU, LR by NSF grant DMS-1613603 and ONR grant N00014-17-1-2145, and OZ by an Israel Science Foundation grant and the ERC advanced grant LogCorrelatedFields.

2 Sketch of the proof

We rely on the Feynman-Kac representation of the solution to (1.1):

$$u(t, x) = \mathbb{E}_B \left[\exp \left\{ \beta \int_0^t V(t-s, x + B_s) ds - \frac{1}{2} \beta^2 R(0)t \right\} \right], \quad (2.1)$$

which has the same distribution, viewed as a random field in x , with t fixed, as

$$Z(t, x) = \mathbb{E}_B [M(t, x)], \quad (2.2)$$

with

$$M(t, x) = \exp\left(\beta \int_0^t V(s, x + B_s) ds - \frac{1}{2} \beta^2 R(0) t\right). \quad (2.3)$$

We used the notation $Z(t, x)$ since it can be viewed as the partition function of a directed polymer of length t and starting at x . For fixed B and x , $M(\cdot, x)$ is a martingale. By [32, Theorem 2.1], for β small enough,

$$\lim_{t \rightarrow \infty} Z(t, 0) = Z_\infty \quad (2.4)$$

almost surely, where Z_∞ is a positive random variable satisfying $\mathbb{E}[Z_\infty] = 1$. Defining

$$Z_\varepsilon(t, x) = Z\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right), \quad M_\varepsilon(t, x) = M\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right),$$

it suffices to consider the random variable

$$X_\varepsilon(t) = \int_{\mathbb{R}^d} \log Z_\varepsilon(t, x) g(x) dx.$$

The main result (1.7) is equivalent to

$$\frac{1}{\varepsilon^{(d-2)/2}} (X_\varepsilon(t) - \mathbb{E}[X_\varepsilon(t)]) \Rightarrow \beta \int_{\mathbb{R}^d} \mathcal{U}(t, x) g(x) dx.$$

Throughout the paper, the temporal variable $t > 0$ is fixed, so sometimes we will omit the dependence. Let us define

$$\sigma_t^2 = \beta^2 \text{Var}\left[\int_{\mathbb{R}^d} \mathcal{U}(t, x) g(x) dx\right] = \beta^2 \nu_{\text{eff}}^2 \int_0^t \int_{\mathbb{R}^{2d}} g(x_1) g(x_2) G_{2s}(x_1 - x_2) dx_1 dx_2 ds, \quad (2.5)$$

where we recall $G_t(x)$ is the centered Gaussian density of variance t in \mathbb{R}^d . The proof of Theorem 1.1 consists of two steps:

Proposition 2.1. *Under the assumption of Theorem 1.1, as $\varepsilon \rightarrow 0$,*

$$\varepsilon^{-(d-2)} \text{Var}[X_\varepsilon(t)] \rightarrow \sigma_t^2.$$

Proposition 2.2. *Under the assumption of Theorem 1.1, as $\varepsilon \rightarrow 0$,*

$$\frac{X_\varepsilon(t) - \mathbb{E}[X_\varepsilon(t)]}{\sqrt{\text{Var}[X_\varepsilon(t)]}} \Rightarrow N(0, 1).$$

To prove the convergence of the variance, we use the Clark-Ocone formula to write $X_\varepsilon - \mathbb{E}[X_\varepsilon]$ as a stochastic integral with respect to \dot{W} . An appropriate decomposition enables us to carry out some explicit calculations. To prove the Gaussianity, we use the second order Poincaré inequality [11, 33] which involves estimating moments of Malliavin derivatives of X_ε and seems to be particularly handy in this context.

In the rest of this section, we describe the main steps in the proof.

2.1 Negative moments

Throughout the paper, we rely on the existence of negative moments of $Z(t, x)$ for small β , as a quantitative control on the small ball probability for $Z_\varepsilon(t, x)$.

Proposition 2.3. *There exists $\beta_0 > 0$ such that if $\beta < \beta_0$,*

$$\sup_{t>0} \mathbb{E}[Z(t, x)^{-n}] \leq C_{\beta, n},$$

for all $n \in \mathbb{N}$.

The proof is presented in Appendix B.

2.2 The Clark-Ocone representation

For each realization of the Brownian motion B , we can write

$$\int_0^{t/\varepsilon^2} V\left(s, \frac{x}{\varepsilon} + B_s\right) ds = \int_0^{t/\varepsilon^2} \left(\int_{\mathbb{R}^d} \varphi\left(\frac{x}{\varepsilon} + B_s - y\right) \dot{W}(s, y) dy \right) ds = \int_{\mathbb{R}^{d+1}} \Phi_{t,x,B}^\varepsilon(s, y) dW(s, y),$$

with

$$\Phi_{t,x,B}^\varepsilon(s, y) = \mathbb{1}_{[0, t/\varepsilon^2]}(s) \varphi\left(\frac{x}{\varepsilon} + B_s - y\right). \quad (2.6)$$

Therefore, we have

$$D_{s,y} Z_\varepsilon(t, x) = D_{s,y} \mathbb{E}_B[M_\varepsilon(t, x)] = \beta \mathbb{E}_B[M_\varepsilon(t, x) \Phi_{t,x,B}^\varepsilon(s, y)],$$

where $D_{s,y}$ denotes the Malliavin derivative operator with respect to \dot{W} .¹ By Lemma A.1, we have

$$D_{s,y} \log Z_\varepsilon(t, x) = \frac{D_{s,y} Z_\varepsilon(t, x)}{Z_\varepsilon(t, x)},$$

and the Clark-Ocone formula gives

$$\begin{aligned} X_\varepsilon - \mathbb{E}[X_\varepsilon] &= \int_{\mathbb{R}^{d+1}} \mathbb{E}[D_{s,y} X_\varepsilon | \mathcal{F}_s] dW(s, y) = \int_{\mathbb{R}^{d+1}} \mathbb{E} \left[\int_{\mathbb{R}^d} \frac{D_{s,y} Z_\varepsilon(t, x)}{Z_\varepsilon(t, x)} g(x) dx \middle| \mathcal{F}_s \right] dW(s, y) \\ &= \beta \int_0^{t/\varepsilon^2} \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} g(x) \mathbb{E} \left[\frac{\mathbb{E}_B[M_\varepsilon(t, x) \Phi_{t,x,B}^\varepsilon(s, y)]}{Z_\varepsilon(t, x)} \middle| \mathcal{F}_s \right] dx \right) dW(s, y). \end{aligned} \quad (2.7)$$

Here, \mathcal{F}_s is the filtration generated by $\dot{W}(\ell, \cdot)$ up to $\ell \leq s$.

For

$$K = \varepsilon^{-\alpha}$$

with some $\alpha > 0$ to be determined, we decompose the stochastic integral in (2.7) into three parts:

$$X_\varepsilon - \mathbb{E}[X_\varepsilon] = \beta(I_{1,\varepsilon} + I_{2,\varepsilon} + I_{3,\varepsilon})$$

with

$$I_{1,\varepsilon} = \int_0^K \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} g(x) \mathbb{E} \left[\frac{\mathbb{E}_B[M_\varepsilon(t, x) \Phi_{t,x,B}^\varepsilon(s, y)]}{Z_\varepsilon(t, x)} \middle| \mathcal{F}_s \right] dx \right) dW(s, y), \quad (2.8)$$

$$I_{2,\varepsilon} = \int_K^{t/\varepsilon^2} \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} g(x) \mathbb{E} \left[\frac{\mathbb{E}_B[M_\varepsilon(t, x) \Phi_{t,x,B}^\varepsilon(s, y)]}{Z(K, x/\varepsilon)} \left(\frac{Z(K, x/\varepsilon)}{Z(t/\varepsilon^2, x/\varepsilon)} - 1 \right) \middle| \mathcal{F}_s \right] dx \right) dW(s, y), \quad (2.9)$$

and

$$I_{3,\varepsilon} = \int_K^{t/\varepsilon^2} \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} g(x) \mathbb{E} \left[\frac{\mathbb{E}_B[M_\varepsilon(t, x) \Phi_{t,x,B}^\varepsilon(s, y)]}{Z(K, x/\varepsilon)} \middle| \mathcal{F}_s \right] dx \right) dW(s, y). \quad (2.10)$$

The goal is to show that if $1 \ll K \ll \varepsilon^{-2}$, then the contribution from $I_{1,\varepsilon}, I_{2,\varepsilon}$ is small compared to that from $I_{3,\varepsilon}$. For $I_{3,\varepsilon}$, since the integration is in $s \geq K$, the random variable $Z(K, x/\varepsilon)$ is \mathcal{F}_s -measurable, thus

$$I_{3,\varepsilon} = \int_K^{t/\varepsilon^2} \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \frac{g(x)}{Z(K, x/\varepsilon)} \mathbb{E} \left[\mathbb{E}_B[M_\varepsilon(t, x) \Phi_{t,x,B}^\varepsilon(s, y)] \middle| \mathcal{F}_s \right] dx \right) dW(s, y).$$

It turns out the inner conditional expectation can be computed explicitly, facilitating the analysis.

¹This involves an abuse of notation: we consider $D_{s,\cdot} Z_\varepsilon(t, x)$ as an element of the Hilbert space $H_1 = L^2(\mathbb{R}^d)$, which is then integrated against the cylindrical white noise $\dot{W}(s, \cdot)$. The Malliavin derivative at time s is then an element of H_1 , which we write as $D_{s,y} Z_\varepsilon(t, x)$. See e.g. [29] for background.

2.3 The second order Poincaré inequality

To simplify the notation, we define

$$Y_\varepsilon = \frac{X_\varepsilon - \mathbb{E}[X_\varepsilon]}{\sqrt{\text{Var}[X_\varepsilon]}}.$$

To show that $Y_\varepsilon \Rightarrow N(0,1)$, we apply the second order Poincaré inequality [33, Theorem 1.1]. Since $\mathbb{E}[Y_\varepsilon] = 0$ and $\text{Var}[Y_\varepsilon] = 1$, with ζ a standard centered Gaussian random variable, we have

$$d_{\text{TV}}(Y_\varepsilon, \zeta) \lesssim \mathbb{E}[\|DY_\varepsilon\|_H^4]^{1/4} \mathbb{E}[\|D^2Y_\varepsilon\|_{\text{op}}^4]^{1/4}. \quad (2.11)$$

Recall our notation convention from Section 1.4; here, $\|D^2Y_\varepsilon\|_{\text{op}}$ denotes the operator norm of the mapping $D^2Y_\varepsilon : H \rightarrow H$ given by $h \mapsto D^2Y_\varepsilon h$, that is,

$$\|D^2Y_\varepsilon\|_{\text{op}} = \sup_{h, g \in H, \|h\|_H = \|g\|_H = 1} \langle g, D^2Y_\varepsilon h \rangle_H.$$

It is clear that

$$DY_\varepsilon = \frac{DX_\varepsilon}{\sqrt{\text{Var}[X_\varepsilon]}}, \quad D^2Y_\varepsilon = \frac{D^2X_\varepsilon}{\sqrt{\text{Var}[X_\varepsilon]}}.$$

Since $\text{Var}[X_\varepsilon] \sim \varepsilon^{d-2}$ by Proposition 2.1, to show $d_{\text{TV}}(Y_\varepsilon, \zeta) \rightarrow 0$ using (2.11), we only need to prove

$$\mathbb{E}[\|DX_\varepsilon\|_H^4]^{1/4} \mathbb{E}[\|D^2X_\varepsilon\|_{\text{op}}^4]^{1/4} = o(\varepsilon^{d-2}), \quad \text{as } \varepsilon \rightarrow 0. \quad (2.12)$$

3 Convergence of the variance

Let us set

$$M_{\varepsilon,j}(t, x) = \exp\left(\beta \int_0^{t/\varepsilon^2} V(s, \frac{x}{\varepsilon} + B_s^j) ds - \frac{\beta^2 R(0)t}{2\varepsilon^2}\right),$$

where B^j are independent Brownian motions. For any set $I \subset \mathbb{R}_+$, $x \in \mathbb{R}^d$ and two Brownian motions B^i, B^j , we define

$$\mathcal{R}(I, x, B^i, B^j) = \int_I R(x + B_s^i - B_s^j) ds \quad (3.1)$$

as the weighted intersection time of B^i, B^j during the time interval I , with x being the initial distance. For $I = [0, T]$, we simply write $\mathcal{R}(T, x, B^i, B^j)$.

Before entering the proof, we present the following lemma which will be used repeatedly.

Lemma 3.1. *For any $n \in \mathbb{Z}_+$ and $q > 1$, there exists $\beta(n, q) > 0$ such that if $\beta < \beta(n, q)$, then for any random variable $F(B^1, \dots, B^n) \geq 0$, $t > 0$ and $\{x_j \in \mathbb{R}^d\}_{j=1, \dots, n}$, we have*

$$\mathbb{E}\left[\frac{\mathbb{E}_B[\prod_{j=1}^n M_{\varepsilon,j}(t, x_j) F(B^1, \dots, B^n)]}{\prod_{j=1}^n Z_\varepsilon(t, x_j)}\right] \lesssim \mathbb{E}_B[|F(B^1, \dots, B^n)|^q]^{1/q}. \quad (3.2)$$

Proof. By the Cauchy-Schwarz inequality and Proposition 2.3, the square of the l.h.s. of (3.2) is bounded by

$$\mathbb{E}\left[\left[\mathbb{E}_B\left[\prod_{j=1}^n M_{\varepsilon,j}(t, x_j) F(B^1, \dots, B^n)\right]\right]^2\right] = \mathbb{E}_B\left[\prod_{j=1}^{2n} M_{\varepsilon,j}(t, x_j) F(B^1, \dots, B^n) F(B^{n+1}, \dots, B^{2n})\right],$$

where $x_{j+n} = x_j$ for $j = 1, \dots, n$. Evaluating the expectation with respect to \tilde{W} , we obtain

$$\mathbb{E}\left[\prod_{j=1}^{2n} M_{\varepsilon,j}(t, x_j)\right] = \exp\left(\frac{\beta^2}{2} \sum_{j,k=1}^n \mathbb{1}_{j \neq k} \mathcal{R}\left(\frac{t}{\varepsilon^2}, \frac{x_j - x_k}{\varepsilon}, B^j, B^k\right)\right).$$

Taking $p = q/(q-1)$, Lemma A.2 shows that the r.h.s. of the above expression has an L^p norm that is bounded uniformly in ε, t and x_j , provided that β is chosen small. We apply Hölder's inequality to complete the proof. \square

3.1 The analysis of $I_{1,\varepsilon}$

Recall that the integral $I_{1,\varepsilon}$ is given by (2.8).

Lemma 3.2. *If $K = \varepsilon^{-\alpha}$ with $\alpha < 4/(2+d)$, we have*

$$\varepsilon^{-(d-2)} \mathbb{E}[I_{1,\varepsilon}^2] \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0.$$

Proof. Writing

$$I_{1,\varepsilon} = \int_0^K \int_{\mathbb{R}^d} \mathbb{E}[\mathbf{Y}_{s,y} | \mathcal{F}_s] dW(s, y),$$

for the appropriate $\mathbf{Y}_{s,y}$ as in (2.8), we have by Itô's isometry that

$$\begin{aligned} \mathbb{E}[I_{1,\varepsilon}^2] &= \int_0^K \int_{\mathbb{R}^d} \mathbb{E}[|\mathbb{E}[\mathbf{Y}_{s,y} | \mathcal{F}_s]|^2] dy ds \leq \int_0^K \int_{\mathbb{R}^d} \mathbb{E}[\mathbf{Y}_{s,y}^2] dy ds \\ &= \beta^2 \int_0^K \int_{\mathbb{R}^d} \int_{\mathbb{R}^{2d}} g(x_1)g(x_2) \mathbb{E}\left[\frac{\mathbb{E}_B[\prod_{j=1}^2 M_{\varepsilon,j}(t, x) \Phi_{t,x_j,B^j}^\varepsilon(s, y)]}{Z_\varepsilon(t, x_1)Z_\varepsilon(t, x_2)}\right] dx_1 dx_2 dy ds. \end{aligned}$$

Using the fact that

$$\int_0^K \int_{\mathbb{R}^d} \prod_{j=1}^2 \Phi_{t,x_j,B^j}^\varepsilon(s, y) dy ds = \int_0^K R\left(\frac{x_1 - x_2}{\varepsilon} + B_s^1 - B_s^2\right) ds = \mathcal{R}\left(K, \frac{x_1 - x_2}{\varepsilon}, B^1, B^2\right),$$

where we recall that R is the spatial covariance function defined in (1.2), we have

$$\mathbb{E}[I_{1,\varepsilon}^2] \lesssim \int_{\mathbb{R}^{2d}} g(x_1)g(x_2) \mathbb{E}\left[\frac{\mathbb{E}_B[\prod_{j=1}^2 M_{\varepsilon,j}(t, x_j) \mathcal{R}(K, \frac{x_1 - x_2}{\varepsilon}, B^1, B^2)]}{Z_\varepsilon(t, x_1)Z_\varepsilon(t, x_2)}\right] dx_1 dx_2.$$

By Lemma 3.1, we have

$$\mathbb{E}[I_{1,\varepsilon}^2] \lesssim \int_{\mathbb{R}^{2d}} g(x_1)g(x_2) \sqrt{\mathbb{E}_B[\mathcal{R}(K, \frac{x_1 - x_2}{\varepsilon}, B^1, B^2)^2]} dx_1 dx_2.$$

By the expression of \mathcal{R} in (3.1), it suffices to use the estimate

$$\mathbb{E}_B[\mathcal{R}(K, \frac{x_1 - x_2}{\varepsilon}, B^1, B^2)^2] \lesssim K^2 \mathbb{P}_B\left[\max_{s \in [0, K]} |B_s^1 - B_s^2| \geq \frac{|x_1 - x_2|}{C\varepsilon}\right] \lesssim K^2 e^{-\frac{|x_1 - x_2|^2}{C\varepsilon^2 K}}$$

for some constant $C > 0$. This implies

$$\mathbb{E}[I_{1,\varepsilon}^2] \lesssim K \int_{\mathbb{R}^{2d}} g(x_1)g(x_2) e^{-\frac{|x_1 - x_2|^2}{C\varepsilon^2 K}} dx_1 dx_2 \lesssim K^{1+\frac{d}{2}} \varepsilon^d.$$

The proof is complete. \square

3.2 The analysis of $I_{2,\varepsilon}$

Recall the definition of $I_{2,\varepsilon}$, see (2.9), and set $K = \varepsilon^{-\alpha}$. We have the following lemma.

Lemma 3.3. *For any $\alpha > 0$, there exists $\beta(\alpha)$ such that if $\beta < \beta(\alpha)$,*

$$\varepsilon^{-(d-2)} \mathbb{E}[I_{2,\varepsilon}^2] \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0.$$

Proof. By the same calculation as in the proof of Lemma 3.2, we have

$$\mathbb{E}[I_{2,\varepsilon}^2] \lesssim \int_{\mathbb{R}^{2d}} g(x_1)g(x_2)A_\varepsilon(x_1, x_2)dx_1dx_2,$$

with

$$A_\varepsilon(x_1, x_2) = \mathbb{E}\left[\frac{\mathbb{E}_B[\prod_{j=1}^2 M_{\varepsilon,j}(t, x_j)\mathcal{R}([K, t/\varepsilon^2], (x_1 - x_2)/\varepsilon, B^1, B^2)]}{Z(K, x_1/\varepsilon)Z(K, x_2/\varepsilon)} \prod_{j=1}^2 \left(\frac{Z(K, x_j/\varepsilon)}{Z(t/\varepsilon^2, x_j/\varepsilon)} - 1\right)\right].$$

Applying Proposition 2.3, Hölder's inequality, and the fact that $Z(t, x)$ is stationary in x , we have

$$\begin{aligned} A_\varepsilon(x_1, x_2) &\lesssim \left(\mathbb{E}[\mathbb{E}_B[\prod_{j=1}^2 M_{\varepsilon,j}(t, x_j)\mathcal{R}([K, t/\varepsilon^2], (x_1 - x_2)/\varepsilon, B^1, B^2)]^2]\right)^{1/2} \\ &\quad \times \left(\mathbb{E}[|Z(K, 0) - Z(t/\varepsilon^2, 0)|^{16}]\right)^{1/8}. \end{aligned} \quad (3.3)$$

For the second factor on the r.h.s. of (3.3), we have

$$\begin{aligned} \mathbb{E}[|Z(K, 0) - Z(t/\varepsilon^2, 0)|^{16}] &\lesssim \sqrt{\mathbb{E}[|Z(K, 0) - Z(t/\varepsilon^2, 0)|^2] \sup_{t>0} \mathbb{E}[|Z(t, 0)|^{30}]} \\ &\lesssim K^{-(d-2)/4} = \varepsilon^{\alpha(d-2)/4}, \end{aligned} \quad (3.4)$$

where the second “ \lesssim ” comes from e.g. [12, Proposition 2.1]. For first factor on the r.h.s. of (3.3), the same calculation as in the proof of Lemma 3.2 yields

$$\begin{aligned} &\mathbb{E}[\mathbb{E}_B[\prod_{j=1}^2 M_{\varepsilon,j}(t, x_j)\mathcal{R}([K, t/\varepsilon^2], (x_1 - x_2)/\varepsilon, B^1, B^2)]^2] \\ &\lesssim \mathbb{E}_B[|\mathcal{R}([K, t/\varepsilon^2], (x_1 - x_2)/\varepsilon, B^1, B^2)|^q]^{2/q} \lesssim \frac{1}{|x_1 - x_2|^{2(d-2)/q}} \varepsilon^{\frac{2(d-2)}{q} - \frac{4}{p}}, \end{aligned} \quad (3.5)$$

where the last step comes from Lemma 3.4 below. Combining (3.4) and (3.5), we have

$$A_\varepsilon(x_1, x_2) \lesssim \frac{\varepsilon^{\frac{(d-2)}{q} - \frac{2}{p}}}{|x_1 - x_2|^{(d-2)/q}} \times \varepsilon^{\frac{\alpha(d-2)}{32}},$$

which implies

$$\mathbb{E}[I_{2,\varepsilon}^2] \lesssim \varepsilon^{\frac{(d-2)}{q} - \frac{2}{p} + \frac{\alpha(d-2)}{32}}.$$

We choose p large enough (for fixed α) such that

$$\frac{(d-2)}{q} - \frac{2}{p} + \frac{\alpha(d-2)}{32} = \frac{d}{q} - 2 + \frac{\alpha(d-2)}{32} > d-2,$$

to complete the proof. \square

Lemma 3.4. *For any $q > 1$, we have*

$$\mathbb{E}_B[|\mathcal{R}(t/\varepsilon^2, x/\varepsilon, B^1, B^2)|^q] \lesssim \frac{\varepsilon^{d-2-\frac{2q}{p}}}{|x|^{d-2}}.$$

Proof. Since $R(x) = 0$ for $|x| \geq 1$, we have

$$|\mathcal{R}(t/\varepsilon^2, x/\varepsilon, B^1, B^2)|^q \lesssim \left(\int_0^{t/\varepsilon^2} \mathbb{1}_{|x/\varepsilon + B_s^1 - B_s^2| \leq 1} ds \right)^q \lesssim \varepsilon^{-2q/p} \int_0^{t/\varepsilon^2} \mathbb{1}_{|x/\varepsilon + B_s^1 - B_s^2| \leq 1} ds.$$

Taking the expectation, we obtain

$$\mathbb{E}[|\mathcal{R}(t/\varepsilon^2, x/\varepsilon, B^1, B^2)|^q] \lesssim \varepsilon^{-2q/p} \int_0^\infty \mathbb{E}_B[\mathbb{1}_{|x/\varepsilon + B_s^1 - B_s^2| \leq 1}] ds \lesssim \varepsilon^{-2q/p} \int_{\mathbb{R}^d} \frac{\mathbb{1}_{|x/\varepsilon + y| \leq 1}}{|y|^{d-2}} dy \lesssim \frac{\varepsilon^{d-2-\frac{2q}{p}}}{|x|^{d-2}},$$

which completes the proof. \square

3.3 The analysis of $I_{3,\varepsilon}$

Recall the definition of $I_{3,\varepsilon}$, see (2.10). Using the fact that $\mathbb{E}[M(t/\varepsilon^2, x/\varepsilon) | \mathcal{F}_s] = M(s, x/\varepsilon)$, we get

$$I_{3,\varepsilon} = \int_K^{t/\varepsilon^2} \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \frac{g(x)}{Z(K, x/\varepsilon)} \mathbb{E}_B[M(s, x/\varepsilon) \Phi_{t,x,B}^\varepsilon(s, y)] dx \right) dW(s, y).$$

For any $T > 0, x_1, x_2 \in \mathbb{R}^d$ and a standard d -dimensional Brownian motion \bar{B} , we define

$$\mathcal{H}(T, x_1, x_2) = \mathbb{E}_B \left[e^{\beta^2 \int_0^T R(x_1 + \bar{B}_{2s}) ds} | \bar{B}_{2T} = x_2 \right].$$

We introduce the following notation: for any $x, y \in \mathbb{R}^d$, the expectation $\hat{\mathbb{E}}_{x,y}$ is defined as

$$\hat{\mathbb{E}}_{x,y}[F] = \mathbb{E} \left[\frac{\mathbb{E}_B[M_1(K, x) M_2(K, y) F]}{Z(K, x) Z(K, y)} \right] \quad (3.6)$$

for any random variable F , where we recall that M_j is the M associated with B^j . In particular, we will consider functionals of

$$\mathbf{X}_K = B_K^1 - B_K^2,$$

so

$$\hat{\mathbb{E}}_{x,y}[F(\mathbf{X}_K)] = \mathbb{E} \left[\frac{\mathbb{E}_B[M_1(K, x) M_2(K, y) F(B_K^1 - B_K^2)]}{Z(K, x) Z(K, y)} \right].$$

The following three lemmas combine to show the convergence of

$$\varepsilon^{-(d-2)} \mathbb{E}[I_{3,\varepsilon}^2] \rightarrow \sigma_t^2, \quad (3.7)$$

with σ_t given in (2.5).

Lemma 3.5.

$$\varepsilon^{-(d-2)} \mathbb{E}[I_{3,\varepsilon}^2] = \int_0^{t-\varepsilon^2 K} \mathcal{G}_\varepsilon(s) ds,$$

with

$$\mathcal{G}_\varepsilon(s) = \int_{\mathbb{R}^{3d}} g(x-w) g(x) R(y) \hat{\mathbb{E}}_{-w/\varepsilon, 0} \left[G_{2s}(w + \varepsilon y - \varepsilon \mathbf{X}_K) \mathcal{H}\left(\frac{s}{\varepsilon^2}, y, \mathbf{X}_K - \frac{w}{\varepsilon} - y\right) \right] dx dy dw. \quad (3.8)$$

Lemma 3.6. *There exists $\beta_0 > 0$ so that there exists $\gamma \in (0, 1)$ such that, for all $\beta < \beta_0$, $\mathcal{G}_\varepsilon(s) \lesssim s^{-\gamma}$ for $s \in (0, t)$.*

Lemma 3.7. *For any $s \in (0, t)$,*

$$\mathcal{G}_\varepsilon(s) \rightarrow \nu_{\text{eff}}^2 \int_{\mathbb{R}^{2d}} g(x-w)g(x)G_{2s}(w)dw dx,$$

as $\varepsilon \rightarrow 0$.

Proof of Lemma 3.5. By Itô's isometry, we have

$$\mathbb{E}[I_{3,\varepsilon}^2] = \int_K^{t/\varepsilon^2} \int_{\mathbb{R}^{3d}} g(x_1)g(x_2)\mathbb{E}\left[\frac{\mathbb{E}_B[\prod_{j=1}^2 M_j(s, x_j/\varepsilon)\Phi_{t,x_j,B^j}^\varepsilon(s, y)]}{Z(K, x_1/\varepsilon)Z(K, x_2/\varepsilon)}\right]dy dx_1 dx_2 ds.$$

Conditioning on \mathcal{F}_K , we have

$$\mathbb{E}\left[\prod_{j=1}^2 M_j(s, x_j/\varepsilon) \mid \mathcal{F}_K\right] = \left(\prod_{j=1}^2 M_j(K, x_j/\varepsilon)\right) \times \exp\left(\beta^2 \int_K^s R\left(\frac{x_1-x_2}{\varepsilon} + B_r^1 - B_r^2\right)dr\right).$$

Integrating in y , we have

$$\begin{aligned} \mathbb{E}[I_{3,\varepsilon}^2] &= \int_K^{t/\varepsilon^2} \int_{\mathbb{R}^{2d}} g(x_1)g(x_2) \\ &\times \mathbb{E}\left[\frac{\mathbb{E}_B[\prod_{j=1}^2 M_j(K, x_j/\varepsilon) \exp\{\beta^2 \int_K^s R\left(\frac{x_1-x_2}{\varepsilon} + B_r^1 - B_r^2\right)dr\} R\left(\frac{x_1-x_2}{\varepsilon} + B_s^1 - B_s^2\right)]}{Z(K, x_1/\varepsilon)Z(K, x_2/\varepsilon)}\right] dx_1 dx_2 ds. \end{aligned} \quad (3.9)$$

Changing variables $x_2 \mapsto x, x_1 \mapsto x + \varepsilon y$ and $s \mapsto \frac{s}{\varepsilon^2}$, and using the stationarity of V in x , we have

$$\begin{aligned} \varepsilon^{-(d-2)}\mathbb{E}[I_{3,\varepsilon}^2] &= \int_{\varepsilon^2 K}^t \int_{\mathbb{R}^{2d}} g(x + \varepsilon y)g(x) \\ &\times \mathbb{E}\left[\frac{\mathbb{E}_B[M_1(K, y)M_2(K, 0) \exp\{\beta^2 \int_K^{s/\varepsilon^2} R(y + B_r^1 - B_r^2)dr\} R(y + B_{s/\varepsilon^2}^1 - B_{s/\varepsilon^2}^2)]}{Z(K, y)Z(K, 0)}\right] dx dy ds. \end{aligned}$$

For $r \geq K$, we can write $B_r^1 - B_r^2 = B_K^1 - B_K^2 + \bar{B}_{2(r-K)}$, where \bar{B} is another Brownian motion independent of B and the random environment V . Thus, recall the definition of $\hat{\mathbb{E}}$ in (3.6), we have

$$\begin{aligned} \varepsilon^{-(d-2)}\mathbb{E}[I_{3,\varepsilon}^2] &= \int_{\varepsilon^2 K}^t \int_{\mathbb{R}^{2d}} g(x + \varepsilon y)g(x) \\ &\times \hat{\mathbb{E}}_{y,0} \mathbb{E}_{\bar{B}}\left[\exp\{\beta^2 \int_K^{s/\varepsilon^2} R(y + \mathbf{X}_K + \bar{B}_{2(r-K)})dr\} R(y + \mathbf{X}_K + \bar{B}_{2(s/\varepsilon^2-K)})\right] dx dy ds. \end{aligned}$$

We write the expectation with respect to \bar{B} more explicitly by conditioning on the end point of \bar{B} :

$$\begin{aligned} &\mathbb{E}_{\bar{B}}\left[e^{\beta^2 \int_K^{s/\varepsilon^2} R(y + \mathbf{X}_K + \sqrt{2}\bar{B}_{r-K})dr} R(y + \mathbf{X}_K + \bar{B}_{2(s/\varepsilon^2-K)})\right] \\ &= \int_{\mathbb{R}^d} G_{2(s-\varepsilon^2 K)}(w)R(y + \mathbf{X}_K + w/\varepsilon)\mathbb{E}_{\bar{B}}\left[e^{\beta^2 \int_K^{s/\varepsilon^2} R(y + \mathbf{X}_K + \bar{B}_{2(r-K)})dr} \mid \bar{B}_{2(s/\varepsilon^2-K)} = w/\varepsilon\right] dw. \end{aligned}$$

Now we consider the integral in y and change variable $y \mapsto y - w/\varepsilon$, then the expectation in \bar{B} in the last display becomes

$$\mathbb{E}_{\bar{B}}\left[e^{\beta^2 \int_K^{s/\varepsilon^2} R(y + \mathbf{X}_K - w/\varepsilon + \bar{B}_{2(r-K)})dr} \mid \bar{B}_{2(s/\varepsilon^2-K)} = \frac{w}{\varepsilon}\right] = \mathbb{E}_{\bar{B}}\left[e^{\beta^2 \int_0^{s/\varepsilon^2-K} R(y + \mathbf{X}_K + \bar{B}_{2r})dr} \mid \bar{B}_{2(s/\varepsilon^2-K)} = -\frac{w}{\varepsilon}\right],$$

and we can write

$$\begin{aligned} \varepsilon^{-(d-2)}\mathbb{E}[I_{3,\varepsilon}^2] &= \int_{\varepsilon^2 K}^t \int_{\mathbb{R}^{3d}} g(x + \varepsilon y - w)g(x)G_{2(s-\varepsilon^2 K)}(w) \\ &\quad \times \hat{\mathbb{E}}_{y-w/\varepsilon,0}[R(y + \mathbf{X}_K)\mathcal{H}(s/\varepsilon^2 - K, y + \mathbf{X}_K, -w/\varepsilon)]dxdydw ds. \end{aligned}$$

We change back variables in the form $w \mapsto w + \varepsilon y$ to obtain

$$\begin{aligned} \varepsilon^{-(d-2)}\mathbb{E}[I_{3,\varepsilon}^2] &= \int_{\varepsilon^2 K}^t \int_{\mathbb{R}^{3d}} g(x - w)g(x)G_{2(s-\varepsilon^2 K)}(w + \varepsilon y) \\ &\quad \times \hat{\mathbb{E}}_{-w/\varepsilon,0}[R(y + \mathbf{X}_K)\mathcal{H}(s/\varepsilon^2 - K, y + \mathbf{X}_K, -w/\varepsilon - y)]dxdydw ds. \end{aligned}$$

Finally, we change variables $y \mapsto y - \mathbf{X}_K$ and $s \mapsto s + \varepsilon^2 K$ to complete the proof. \square

Proof of Lemma 3.6. By Lemma A.2, we know that \mathcal{H} is uniformly bounded for small β , so that

$$\mathcal{G}_\varepsilon(s) \lesssim \int_{\mathbb{R}^{3d}} |g(x - w)g(x)| R(y)\hat{\mathbb{E}}_{-w/\varepsilon,0}[G_{2s}(w + \varepsilon y - \varepsilon\mathbf{X}_K)]dxdydw.$$

We bound the expectation by Lemma 3.1: for any $q > 1$, if $\beta < \beta(q)$ then

$$\begin{aligned} \hat{\mathbb{E}}_{-w/\varepsilon,0}[G_{2s}(w + \varepsilon y - \varepsilon\mathbf{X}_K)] &= \mathbb{E}\left[\frac{\mathbb{E}_B[M_1(K, -w/\varepsilon)M_2(K, 0)G_{2s}(w + \varepsilon y - \varepsilon\mathbf{X}_K)]}{Z(K, -w/\varepsilon)Z(K, 0)}\right] \\ &\lesssim \mathbb{E}_B[|G_{2s}(w + \varepsilon y - \varepsilon\mathbf{X}_K)|^q]^{1/q}. \end{aligned}$$

Recalling that $\mathbf{X}_K = B_K^1 - B_K^2 \sim N(0, 2K)$, the above expectation can be computed explicitly:

$$\mathbb{E}_B[|G_{2s}(w + \varepsilon y - \varepsilon\mathbf{X}_K)|^q]^{1/q} \lesssim \left(\frac{1}{s^{d(q-1)/2}}G_{\frac{2s}{q}+2\varepsilon^2 K}(w + \varepsilon y)\right)^{1/q}.$$

Thus

$$\mathcal{G}_\varepsilon(s) \lesssim \int_{\mathbb{R}^{3d}} |g(x)| R(y) \left(\frac{1}{s^{d(q-1)/2}}G_{\frac{2s}{q}+2\varepsilon^2 K}(w)\right)^{1/q} dxdydw \lesssim s^{-\frac{d}{2p}}, \quad (3.10)$$

where (p, q) are the dual Hölder exponents and we used the fact that $s < t$ hence $2s/q + 2\varepsilon^2 K \lesssim 1$ in the last step. We choose $p > d/2$ and adjust β accordingly to complete the proof. \square

Proof of Lemma 3.7. Recall that

$$\mathcal{G}_\varepsilon(s) = \int_{\mathbb{R}^{3d}} g(x - w)g(x)R(y)\hat{\mathbb{E}}_{-w/\varepsilon,0}\left[G_{2s}(w + \varepsilon y - \varepsilon\mathbf{X}_K)\mathcal{H}\left(\frac{s}{\varepsilon^2}, y, \mathbf{X}_K - \frac{w}{\varepsilon} - y\right)\right]dxdydw.$$

Since $s > 0$ is fixed, the expectation in the above expression is bounded uniformly in x, y, w , so we only need to pass to the limit of the expectation for fixed $x, y, w \in \mathbb{R}^d$ and $w \neq 0$. The proof is divided into three steps.

(i) We show that $\hat{\mathbb{E}}_{-w/\varepsilon,0}[|G_{2s}(w + \varepsilon y - \varepsilon\mathbf{X}_K) - G_{2s}(w)|] \rightarrow 0$ as $\varepsilon \rightarrow 0$. Using the fact that

$$|G_{2s}(w + \varepsilon y - \varepsilon\mathbf{X}_K) - G_{2s}(w)| \lesssim \varepsilon|y| + \varepsilon|\mathbf{X}_K|,$$

it suffices to show $\hat{\mathbb{E}}_{-w/\varepsilon,0}[|\varepsilon\mathbf{X}_K|] \rightarrow 0$. We apply Lemma 3.1 to get

$$\hat{\mathbb{E}}_{-w/\varepsilon,0}[|\varepsilon\mathbf{X}_K|] \lesssim \sqrt{\mathbb{E}_B[|\varepsilon\mathbf{X}_K|^2]} = \sqrt{2\varepsilon^2 K} \rightarrow 0. \quad (3.11)$$

(ii) For $\alpha \in (0, 2)$, define

$$\tilde{\mathcal{H}}_\varepsilon = \mathbb{E}_{\bar{B}} \left[\exp \left\{ \beta^2 \int_0^{s/\varepsilon^\alpha} R(y + \bar{B}_{2r}) dr \right\} \middle| \bar{B}_{2s/\varepsilon^2} = \mathbf{X}_K - \frac{w}{\varepsilon} - y \right],$$

we show that

$$\hat{\mathbb{E}}_{-w/\varepsilon, 0}[\tilde{\mathcal{H}}_\varepsilon] \rightarrow \mathbb{E}_{\bar{B}} \left[\exp \left\{ \beta^2 \int_0^\infty R(y + \bar{B}_{2s}) ds \right\} \right] \quad (3.12)$$

as $\varepsilon \rightarrow 0$. Note that $\tilde{\mathcal{H}}$ can be written more explicitly by conditioning on $\bar{B}_{2s/\varepsilon^\alpha}$:

$$\begin{aligned} \tilde{\mathcal{H}}_\varepsilon &= \mathbb{E}_{\bar{B}} \left[\exp \left\{ \beta^2 \int_0^{s/\varepsilon^\alpha} R(y + \bar{B}_{2r}) dr \right\} \times \frac{G_{2s(1-\varepsilon^{2-\alpha})}(\varepsilon \mathbf{X}_K - w - \varepsilon y - \varepsilon \bar{B}_{2s/\varepsilon^\alpha})}{G_{2s}(\varepsilon \mathbf{X}_K - w - \varepsilon y)} \right] \\ &= \frac{1}{(1 - \varepsilon^{2-\alpha})^{d/2}} \mathbb{E}_{\bar{B}} \left[\exp \left\{ \beta^2 \int_0^{s/\varepsilon^\alpha} R(y + \bar{B}_{2r}) dr \right\} \right. \\ &\quad \left. \times \exp \left\{ -\frac{(\varepsilon \mathbf{X}_K - w - \varepsilon y - \varepsilon \bar{B}_{2s/\varepsilon^\alpha})^2}{4s(1 - \varepsilon^{2-\alpha})} + \frac{(\varepsilon \mathbf{X}_K - w - \varepsilon y)^2}{4s} \right\} \right]. \end{aligned}$$

There are three factors inside the above expectation. First, we note that again by an application of Lemma 3.1, we have

$$\hat{\mathbb{E}}_{-w/\varepsilon, 0}[e^{\lambda|\varepsilon \mathbf{X}_K|^2}] \lesssim 1,$$

for any $\lambda > 0$. Then by the same proof as for (i), we can replace the second factor by $e^{-w^2/4s}$ with a negligible error. Finally we use the simple inequality $|e^x - e^y| \leq (e^x + e^y)|x - y|$ to replace the third factor by $e^{w^2/4s}$ with a negligible error. This proves (3.12).

(iii) We show that

$$\hat{\mathbb{E}}_{-w/\varepsilon, 0} \left[\left| \mathcal{H}\left(\frac{s}{\varepsilon^2}, y, \mathbf{X}_K - \frac{w}{\varepsilon} - y\right) - \tilde{\mathcal{H}}_\varepsilon \right| \right] \rightarrow 0 \quad (3.13)$$

as $\varepsilon \rightarrow 0$. Since $R \geq 0$ and $R(x) = 0$ for $|x| \geq 1$, we have, using Lemma A.2

$$\begin{aligned} & \left| \mathcal{H}\left(\frac{s}{\varepsilon^2}, y, \mathbf{X}_K - \frac{w}{\varepsilon} - y\right) - \tilde{\mathcal{H}}_\varepsilon \right| \\ &= \mathbb{E}_{\bar{B}} \left[\exp \left\{ \beta^2 \int_0^{s/\varepsilon^2} R(y + \bar{B}_{2r}) dr \right\} - \exp \left\{ \beta^2 \int_0^{s/\varepsilon^\alpha} R(y + \bar{B}_{2r}) dr \right\} \middle| \bar{B}_{2s/\varepsilon^2} = \mathbf{X}_K - \frac{w}{\varepsilon} - y \right] \\ &\leq \mathbb{E}_{\bar{B}} \left[\exp \left\{ \beta^2 \int_0^{s/\varepsilon^2} R(y + \bar{B}_{2r}) dr \right\} \mathbb{1}_{\min_{r \in [s/\varepsilon^\alpha, s/\varepsilon^2]} |y + \bar{B}_{2r}| \leq 1} \middle| \bar{B}_{2s/\varepsilon^2} = \mathbf{X}_K - \frac{w}{\varepsilon} - y \right] \\ &\lesssim \sqrt{\mathbb{P}_{\bar{B}} \left[\min_{r \in [s/\varepsilon^\alpha, s/\varepsilon^2]} |y + \bar{B}_{2r}| \leq 1 \middle| \bar{B}_{2s/\varepsilon^2} = \mathbf{X}_K - \frac{w}{\varepsilon} - y \right]}. \end{aligned} \quad (3.14)$$

By Lemma A.3 and (3.11), we obtain (3.13).

To summarize, we have

$$\begin{aligned} \mathcal{G}_\varepsilon(s) &\rightarrow \int_{\mathbb{R}^{3d}} g(x-w)g(x)R(y)G_{2s}(w)\mathbb{E}_{\bar{B}} \left[\exp \left\{ \beta^2 \int_0^\infty R(y + \bar{B}_{2s}) ds \right\} \right] dx dy dw \\ &= \nu_{\text{eff}}^2 \int_{\mathbb{R}^{2d}} g(x-w)g(x)G_{2s}(w) dx dw, \end{aligned}$$

which completes the proof. \square

3.4 Proof of Proposition 2.1

Recall that $X_\varepsilon - \mathbb{E}[X_\varepsilon] = \beta(I_{1,\varepsilon} + I_{2,\varepsilon} + I_{3,\varepsilon})$. Choosing $K = \varepsilon^{-\alpha}$ with $\alpha < 4/(2+d)$ and β small, we combine Lemmas 3.2, 3.3 and (3.7) to obtain

$$\varepsilon^{-(d-2)} \text{Var}[X_\varepsilon] \rightarrow \sigma_t^2.$$

Remark 3.8. A simpler version of the proof will show the convergence of

$$\varepsilon^{-(d-2)} \text{Var} \left[\int_{\mathbb{R}^d} Z_\varepsilon(t, x) g(x) dx \right] \rightarrow \sigma_t^2,$$

that is, the convergence of the variance for the solution of the stochastic heat equation itself. A key identity in the proof of Lemma 3.5 is (3.9):

$$\begin{aligned} \mathbb{E}[I_{3,\varepsilon}^2] &= \int_K^{t/\varepsilon^2} \int_{\mathbb{R}^{2d}} g(x_1) g(x_2) \\ &\quad \times \hat{\mathbb{E}}_{\frac{x_1}{\varepsilon}, \frac{x_2}{\varepsilon}} \mathbb{E}_B \left[\exp \left\{ \beta^2 \int_K^s R \left(\frac{x_1 - x_2}{\varepsilon} + B_r^1 - B_r^2 \right) dr \right\} R \left(\frac{x_1 - x_2}{\varepsilon} + B_s^1 - B_s^2 \right) \right] dx_1 dx_2 ds. \end{aligned}$$

For the stochastic heat equation, it is straightforward to check that the above term becomes

$$\int_K^{t/\varepsilon^2} \int_{\mathbb{R}^{2d}} g(x_1) g(x_2) \mathbb{E}_B \left[\exp \left\{ \beta^2 \int_K^s R \left(\frac{x_1 - x_2}{\varepsilon} + B_r^1 - B_r^2 \right) dr \right\} R \left(\frac{x_1 - x_2}{\varepsilon} + B_s^1 - B_s^2 \right) \right] dx_1 dx_2 ds.$$

The difference between these expressions comes from the distribution of (B_K^1, B_K^2) . In the case of KPZ, B_K^j are distributed according to the polymer measure; in the case of SHE, B_K^j is not weighed by the environment hence has distribution $N(0, K)$. It is clear from (3.11) that the asymptotic behaviors of εB_K^j are the same in two cases, which leads to the same limiting variances.

4 Gaussianity

We turn to the proof of Proposition 2.2. The second order Poincaré inequality (2.11) reduces our task to showing that

$$\mathbb{E}[\|DX_\varepsilon\|_H^4]^{1/4} \mathbb{E}[\|D^2X_\varepsilon\|_{\text{op}}^4]^{1/4} = o(\varepsilon^{d-2}), \text{ as } \varepsilon \rightarrow 0.$$

Since

$$DX_\varepsilon = \int_{\mathbb{R}^d} \frac{DZ_\varepsilon(t, x)}{Z_\varepsilon(t, x)} g(x) dx,$$

we have

$$D^2X_\varepsilon = D \int_{\mathbb{R}^d} \frac{DZ_\varepsilon(t, x)}{Z_\varepsilon(t, x)} g(x) dx = \int_{\mathbb{R}^d} \frac{Z_\varepsilon(t, x) D^2Z_\varepsilon(t, x) - DZ_\varepsilon(t, x) \otimes DZ_\varepsilon(t, x)}{Z_\varepsilon^2(t, x)} g(x) dx.$$

Using the Feynman-Kac representation (2.2)-(2.3) gives

$$D^2Z_\varepsilon(t, x) = \beta^2 \mathbb{E}_B [M_\varepsilon(t, x) \Phi_{t,x,B}^\varepsilon \otimes \Phi_{t,x,B}^\varepsilon],$$

so that

$$Z_\varepsilon(t, x) D^2Z_\varepsilon(t, x) = \beta^2 \mathbb{E}_B \left[\prod_{j=1}^2 M_{\varepsilon,j}(t, x) \Phi_{t,x,B^2}^\varepsilon \otimes \Phi_{t,x,B^2}^\varepsilon \right],$$

and

$$DZ_\varepsilon(t, x) \otimes DZ_\varepsilon(t, x) = \beta^2 \mathbb{E}_B \left[\prod_{j=1}^2 M_{\varepsilon, j}(t, x) \Phi_{t, x, B^1}^\varepsilon \otimes \Phi_{t, x, B^2}^\varepsilon \right].$$

Thus, we can write

$$D^2 X_\varepsilon = \beta^2 \int_{\mathbb{R}^d} \frac{\mathbb{E}_B[\prod_{j=1}^2 M_{\varepsilon, j}(t, x) (\Phi_{t, x, B^2}^\varepsilon - \Phi_{t, x, B^1}^\varepsilon) \otimes \Phi_{t, x, B^2}^\varepsilon]}{Z_\varepsilon^2(t, x)} g(x) dx = \mathcal{P}_2 - \mathcal{P}_1,$$

where

$$\mathcal{P}_k = \beta^2 \int_{\mathbb{R}^d} \frac{\mathbb{E}_B[\prod_{j=1}^2 M_{\varepsilon, j}(t, x) \Phi_{t, x, B^k}^\varepsilon \otimes \Phi_{t, x, B^2}^\varepsilon]}{Z_\varepsilon^2(t, x)} g(x) dx \in H \otimes H.$$

Recall that $\Phi_{t, x, B}^\varepsilon$, defined in (2.6), is an element of H for each (ε, t, x, B) fixed. Thus, we have

$$\|D^2 X_\varepsilon\|_{\text{op}}^4 \lesssim \|\mathcal{P}_1\|_{\text{op}}^4 + \|\mathcal{P}_2\|_{\text{op}}^4,$$

and we only need to estimate $\mathbb{E}[\|\mathcal{P}_k\|_{\text{op}}^4]$.

4.1 The first derivative

The goal of this section is to show the following lemma.

Lemma 4.1. *For any $\delta > 0$, there exists $\beta(\delta) > 0$ such that if $\beta < \beta(\delta)$, then*

$$\mathbb{E}[\|DX_\varepsilon\|_H^4]^{1/4} \lesssim \varepsilon^{\frac{d-2}{2}-\delta}.$$

Proof. A direct calculation gives

$$\|DX_\varepsilon\|_H^4 = \beta^4 \int_{\mathbb{R}^{4d}} \prod_{j=1}^4 \frac{g(x_j)}{Z_\varepsilon(t, x_j)} \mathbb{E}_B \left[\prod_{j=1}^4 M_{\varepsilon, j}(t, x_j) \mathcal{R}\left(\frac{t}{\varepsilon^2}, \frac{x_1 - x_2}{\varepsilon}, B^1, B^2\right) \mathcal{R}\left(\frac{t}{\varepsilon^2}, \frac{x_3 - x_4}{\varepsilon}, B^3, B^4\right) \right] dx,$$

with \mathcal{R} defined in (3.1). Taking the expectation \mathbb{E} and applying Proposition 2.3 and Lemma 3.1, we have

$$\begin{aligned} \mathbb{E}[\|DX_\varepsilon\|_H^4] &\lesssim \int_{\mathbb{R}^{4d}} \prod_{j=1}^4 |g(x_j)| \mathbb{E}_B \left[\mathcal{R}^q\left(\frac{t}{\varepsilon^2}, \frac{x_1 - x_2}{\varepsilon}, B^1, B^2\right) \mathcal{R}^q\left(\frac{t}{\varepsilon^2}, \frac{x_3 - x_4}{\varepsilon}, B^3, B^4\right) \right]^{1/q} \\ &\lesssim \varepsilon^{\frac{2(d-2)}{q} - \frac{4}{p}} \int_{\mathbb{R}^{4d}} \prod_{j=1}^4 |g(x_j)| \frac{1}{|x_1 - x_2|^{\frac{d-2}{q}}} \frac{1}{|x_3 - x_4|^{\frac{d-2}{q}}} dx \lesssim \varepsilon^{\frac{2(d-2)}{q} - \frac{4}{p}}. \end{aligned}$$

We applied Lemma 3.4 in the next to last step. It remains to choose p large enough so that

$$\frac{2(d-2)}{q} - \frac{4}{p} = \frac{2d}{q} - 4 > 2(d-2) - 4\delta$$

to complete the proof. \square

4.2 The second derivative

To estimate $\|\mathcal{P}_k\|_{\text{op}}$, we use the contraction inequality [33, Proposition 4.1], which says that

$$\|\mathcal{P}_k\|_{\text{op}}^4 \leq \|\mathcal{P}_k \otimes_1 \mathcal{P}_k\|_{H \otimes H}^2.$$

4.2.1 The case $k = 1$

A direct calculation gives

$$\mathcal{P}_1 \otimes_1 \mathcal{P}_1 = \beta^4 \int_{\mathbb{R}^{2d}} \frac{\mathbb{E}_B[\prod_{j=1}^4 M_{\varepsilon,j}(t, x_j) \mathcal{R}(\frac{t}{\varepsilon^2}, \frac{x-y}{\varepsilon}, B^1, B^3) \Phi_{t,x,B^2}^\varepsilon \otimes \Phi_{t,y,B^4}^\varepsilon]}{Z_\varepsilon^2(t, x) Z_\varepsilon^2(t, y)} g(x) g(y) dx dy,$$

where we write $x_1 = x_2 = x, x_3 = x_4 = y$ to simplify the notations. Thus,

$$\begin{aligned} \|\mathcal{P}_1 \otimes_1 \mathcal{P}_1\|_{H \otimes H}^2 &= \beta^8 \int_{\mathbb{R}^{4d}} g(x) g(y) g(z) g(w) \left(\prod_{j=1}^8 Z_\varepsilon(t, x_j) \right)^{-1} \\ &\quad \times \mathbb{E}_B \left[\prod_{j=1}^8 M_{\varepsilon,j}(t, x_j) \prod_{(i,k) \in \mathcal{O}} \mathcal{R}(\frac{t}{\varepsilon^2}, \frac{x_i - x_k}{\varepsilon}, B^i, B^k) \right] dx dy dz dw, \end{aligned}$$

where $x_5 = x_6 = z, x_7 = x_8 = w$, and the set \mathcal{O} is

$$\mathcal{O} = \{(1, 3), (5, 7), (2, 6), (4, 8)\}.$$

Lemma 4.2. *For any $\delta > 0$, there exists $\beta(\delta)$ such that if $\beta < \beta(\delta)$,*

$$\mathbb{E}[\|\mathcal{P}_1 \otimes_1 \mathcal{P}_1\|_{H \otimes H}^2] \lesssim \varepsilon^{4d-8-\delta} \mathbb{1}_{d < 8} + \varepsilon^{3d-8-\delta} \mathbb{1}_{d \geq 8}.$$

Proof. Applying Lemma 3.1, we have

$$\begin{aligned} \mathbb{E}[\|\mathcal{P}_1 \otimes_1 \mathcal{P}_1\|_{H \otimes H}^2] &\lesssim \int_{\mathbb{R}^{4d}} |g(x) g(y) g(z) g(w)| \times \mathbb{E}_B \left[\prod_{(i,k) \in \mathcal{O}} \mathcal{R}^q(\frac{t}{\varepsilon^2}, \frac{x_i - x_k}{\varepsilon}, B^i, B^k) \right]^{1/q} dx dy dz dw \\ &= \int_{\mathbb{R}^{4d}} |g(x) g(y) g(z) g(w)| \times \prod_{(i,k) \in \mathcal{O}} \mathbb{E}_B \left[\mathcal{R}^q(\frac{t}{\varepsilon^2}, \frac{x_i - x_k}{\varepsilon}, B^i, B^k) \right]^{1/q} dx dy dz dw \end{aligned}$$

for some $q > 1$. We discuss two cases.

(i) $d < 8$. Applying Lemma 3.4 to all pairs $(i, k) \in \mathcal{O}$, the above integral is bounded by

$$\varepsilon^{4(\frac{d-2}{q} - \frac{2}{p})} \int_{\mathbb{R}^{4d}} \frac{|g(x) g(y) g(z) g(w)|}{(|x-y||z-w||x-z||y-w|)^{(d-2)/q}} dx dy dz dw.$$

Since $g \in \mathcal{C}_c$, by the elementary inequality

$$\int_{|y| \leq M} \frac{1}{|x-y|^{\alpha_1}} \frac{1}{|y|^{\alpha_2}} dy \lesssim \frac{1}{|x|^{\alpha_1 + \alpha_2 - d}}, \quad \text{if } \alpha_1 < d, \alpha_2 < d, \alpha_1 + \alpha_2 > d,$$

the above integral is bounded in $d < 8$. Thus, we have

$$\mathbb{E}[\|\mathcal{P}_1 \otimes_1 \mathcal{P}_1\|_{H \otimes H}^2] \lesssim \varepsilon^{4d-8-\delta}.$$

(ii) $d \geq 8$. Applying Lemma 3.4 to three pairs of $(i, k) \in \mathcal{O}$, and bounding the fourth pair simply by ε^{-2} , we have

$$\begin{aligned} \mathbb{E}[\|\mathcal{P}_1 \otimes_1 \mathcal{P}_1\|_{H \otimes H}^2] &\lesssim \varepsilon^{3(\frac{d-2}{q} - \frac{2}{p})-2} \int_{\mathbb{R}^{4d}} \frac{|g(x) g(y) g(z) g(w)|}{(|x-y||z-w||x-z|)^{(d-2)/q}} dx dy dz dw \\ &\lesssim \varepsilon^{3(d-2)-2-\delta}. \end{aligned}$$

The proof is complete. \square

4.2.2 The case $k = 2$

In this case,

$$\mathcal{P}_2 = \beta^2 \int_{\mathbb{R}^d} \frac{\mathbb{E}_B[M_\varepsilon(t, x) \Phi_{t, x, B}^\varepsilon \otimes \Phi_{t, x, B}^\varepsilon]}{Z_\varepsilon(t, x)} g(x) dx,$$

so

$$\mathcal{P}_2 \otimes_1 \mathcal{P}_2 = \beta^4 \int_{\mathbb{R}^{2d}} \frac{\mathbb{E}_B[\prod_{j=1}^2 M_{\varepsilon, j}(t, x_j) \mathcal{R}(\frac{t}{\varepsilon^2}, \frac{x_1 - x_2}{\varepsilon}, B^1, B^2) \Phi_{t, x_1, B^1}^\varepsilon \otimes \Phi_{t, x_2, B^2}^\varepsilon]}{Z_\varepsilon(t, x_1) Z_\varepsilon(t, x_2)} g(x_1) g(x_2) dx_1 dx_2,$$

and

$$\|\mathcal{P}_2 \otimes_1 \mathcal{P}_2\|_{H \otimes H}^2 = \beta^8 \int_{\mathbb{R}^{4d}} \prod_{j=1}^4 \frac{g(x_j)}{Z_\varepsilon(t, x_j)} \mathbb{E}_B \left[\prod_{j=1}^4 M_{\varepsilon, j}(t, x_j) \prod_{(i, k) \in \tilde{\mathcal{O}}} \mathcal{R}(\frac{t}{\varepsilon^2}, \frac{x_i - x_k}{\varepsilon}, B^i, B^k) \right] dx,$$

with

$$\tilde{\mathcal{O}} = \{(1, 2), (3, 4), (1, 3), (2, 4)\}.$$

Lemma 4.3. *For any $\delta > 0$, there exists $\beta(\delta)$ such that if $\beta < \beta(\delta)$,*

$$\mathbb{E}[\|\mathcal{P}_2 \otimes_1 \mathcal{P}_2\|_{H \otimes H}^2] \lesssim \varepsilon^{3(d-2)-\delta}.$$

Proof. By Lemma 3.1 and the fact that g is compactly supported, we have

$$\begin{aligned} \mathbb{E}[\|\mathcal{P}_2 \otimes_1 \mathcal{P}_2\|_{H \otimes H}^2] &\lesssim \int_{\mathbb{R}^{4d}} \prod_{j=1}^4 |g(x_j)| \times \mathbb{E}_B \left[\prod_{(i, k) \in \tilde{\mathcal{O}}} \mathcal{R}^q(\frac{t}{\varepsilon^2}, \frac{x_i - x_k}{\varepsilon}, B^i, B^k) \right]^{1/q} dx \\ &\lesssim \left(\int_{\mathbb{R}^{4d}} \prod_{j=1}^4 |g(x_j)|^q \times \mathbb{E}_B \left[\prod_{(i, k) \in \tilde{\mathcal{O}}} \mathcal{R}^q(\frac{t}{\varepsilon^2}, \frac{x_i - x_k}{\varepsilon}, B^i, B^k) \right] dx \right)^{1/q}. \end{aligned}$$

Since

$$\mathcal{R}^q(\frac{t}{\varepsilon^2}, \frac{x_i - x_k}{\varepsilon}, B^i, B^k) = \left(\int_0^{t/\varepsilon^2} R(\frac{x_i - x_k}{\varepsilon} + B_s^i - B_s^k) ds \right)^q \leq \left(\frac{t}{\varepsilon^2} \right)^{q/p} \int_0^{t/\varepsilon^2} R^q(\frac{x_i - x_k}{\varepsilon} + B_s^i - B_s^k) ds,$$

we only need to control

$$\int_{\mathbb{R}^{4d}} \prod_{j=1}^4 |g(x_j)|^q \times \mathbb{E}_B \left[\prod_{(i, k) \in \tilde{\mathcal{O}}} \int_0^{t/\varepsilon^2} R^q(\frac{x_i - x_k}{\varepsilon} + B_s^i - B_s^k) ds \right] dx.$$

Applying Lemma 4.4, we have

$$\mathbb{E}[\|\mathcal{P}_2 \otimes_1 \mathcal{P}_2\|_{H \otimes H}^2] \lesssim \left(\frac{1}{\varepsilon^2} \right)^{4/p} \varepsilon^{3(d-2)/q} = \varepsilon^{\frac{3(d-2)}{q} - \frac{8}{p}}.$$

The proof is complete. \square

Lemma 4.4. *Assume $0 \leq f, h \in \mathcal{C}_c^\infty(\mathbb{R}^d)$, then*

$$\int_{\mathbb{R}^{4d}} \mathbb{E}_B \left[\prod_{j=1}^4 f(x_j) \prod_{(i, k) \in \tilde{\mathcal{O}}} \int_0^{t/\varepsilon^2} h(\frac{x_i - x_k}{\varepsilon} + B_s^i - B_s^k) ds \right] dx \lesssim \varepsilon^{3(d-2)}.$$

Proof. Without loss of generality, assume h is even. To simplify the notation, we write

$$\begin{aligned} \prod_{(i,k) \in \bar{O}} \int_0^{t/\varepsilon^2} h\left(\frac{x_i - x_k}{\varepsilon} + B_s^i - B_s^k\right) ds &= \int_{[0, t/\varepsilon^2]^4} \prod_{j=1}^4 h\left(\frac{x_j - x_{j-1}}{\varepsilon} + B_{s_j}^j - B_{s_j}^{j-1}\right) ds \\ &= \sum_{\tau} \int_{A_{\tau}} \prod_{j=1}^4 h\left(\frac{x_j - x_{j-1}}{\varepsilon} + B_{s_j}^j - B_{s_j}^{j-1}\right) ds, \end{aligned}$$

where we let $x_0 = x_4$, $B^0 = B^4$, τ denotes the permutations of s_1, \dots, s_4 , and $A_{\tau} \subset [0, t/\varepsilon^2]^4$ corresponds to the permutation τ . Due to symmetry, there are six different permutations to consider.

Now we write the integral in the Fourier domain. Denote $\hat{f}(\xi) = \int f(x) e^{-i\xi \cdot x} dx$ as the Fourier transform of f , we have

$$\begin{aligned} \int_{\mathbb{R}^{4d}} \prod_{j=1}^4 f(x_j) h\left(\frac{x_j - x_{j-1}}{\varepsilon} + B_{s_j}^j - B_{s_j}^{j-1}\right) dx &= \frac{1}{(2\pi)^{4d}} \int_{\mathbb{R}^{8d}} \prod_{j=1}^4 f(x_j) \hat{h}(\eta_j) e^{i\eta_j \cdot (x_j - x_{j-1})/\varepsilon} e^{i\eta_j \cdot (B_{s_j}^j - B_{s_j}^{j-1})} d\eta dx \\ &= \frac{1}{(2\pi)^{4d}} \int_{\mathbb{R}^{4d}} \prod_{j=1}^4 \hat{f}\left(\frac{\eta_j - \eta_{j-1}}{\varepsilon}\right) \hat{h}(\eta_j) e^{i(\eta_j \cdot B_{s_j}^j - \eta_{j+1} \cdot B_{s_{j+1}}^j)} d\eta, \end{aligned}$$

with $\eta_5 = \eta_1$, $s_5 = s_1$. Thus, it suffices to estimate

$$\int_{A_{\tau}} \int_{\mathbb{R}^{4d}} \prod_{j=1}^4 \hat{f}\left(\frac{\eta_j - \eta_{j-1}}{\varepsilon}\right) \hat{h}(\eta_j) \mathbb{E}_B[e^{i(\eta_j \cdot B_{s_j} - \eta_{j+1} \cdot B_{s_{j+1}})}] d\eta.$$

First, we change variables

$$\eta_1 = \xi_1, \quad \eta_j = \xi_1 + \varepsilon(\xi_2 + \dots + \xi_j), \quad j = 2, 3, 4,$$

and the above integral equals to

$$\varepsilon^{3d} \int_{A_{\tau}} \int_{\mathbb{R}^{4d}} \hat{h}(\xi_1) \hat{f}(\xi_2) \hat{f}(\xi_3) \hat{f}(\xi_4) w_{\varepsilon}(\xi) \prod_{j=1}^4 \mathbb{E}_B[e^{i(\eta_j \cdot B_{s_j} - \eta_{j+1} \cdot B_{s_{j+1}})}] d\xi$$

with

$$w_{\varepsilon}(\xi) = \hat{f}(-\xi_2 - \xi_3 - \xi_4) \hat{h}(\eta_2) \hat{h}(\eta_3) \hat{h}(\eta_4) \in L^{\infty}(\mathbb{R}^{4d}).$$

Depending on the permutation τ , the factor $\prod_{j=1}^4 \mathbb{E}_B[e^{i(\eta_j \cdot B_{s_j} - \eta_{j+1} \cdot B_{s_{j+1}})}]$ can be computed explicitly. Since all six cases are treated in the same way, we only take $s_1 < s_2 < s_3 < s_4$ as an example:

$$\begin{aligned} \prod_{j=1}^4 \mathbb{E}_B[e^{i(\eta_j \cdot B_{s_j} - \eta_{j+1} \cdot B_{s_{j+1}})}] &= e^{-\frac{1}{2}|\eta_4 - \eta_1|^2 s_1} e^{-\frac{1}{2}|\eta_4|^2 (s_4 - s_1)} \prod_{j=1}^3 e^{-\frac{1}{2}|\eta_j - \eta_{j+1}|^2 s_j} e^{-\frac{1}{2}|\eta_{j+1}|^2 (s_{j+1} - s_j)} \\ &= e^{-\frac{1}{2} \sum_{j=1}^4 \lambda_j (s_j - s_{j-1})} \end{aligned}$$

with $s_0 = 0$ and

$$\begin{aligned} \lambda_1 &= \varepsilon^2 (|\xi_2|^2 + |\xi_3|^2 + |\xi_4|^2 + |\xi_2 + \xi_3 + \xi_4|^2), \\ \lambda_2 &= \varepsilon^2 |\xi_3|^2 + \varepsilon^2 |\xi_4|^2 + |\xi_1 + \varepsilon \xi_2|^2 + |\xi_1 + \varepsilon(\xi_2 + \xi_3 + \xi_4)|^2, \\ \lambda_3 &= \varepsilon^2 |\xi_4|^2 + |\xi_1 + \varepsilon \xi_2 + \varepsilon \xi_3|^2 + |\xi_1 + \varepsilon(\xi_2 + \xi_3 + \xi_4)|^2, \\ \lambda_4 &= 2|\xi_1 + \varepsilon(\xi_2 + \xi_3 + \xi_4)|^2. \end{aligned}$$

After integrating the s variables, we have

$$\int_{0 < s_1 < \dots < s_4 < t/\varepsilon^2} \prod_{j=1}^4 \mathbb{E}_B [e^{i(\eta_j \cdot B_{s_j} - \eta_{j+1} \cdot B_{s_{j+1}})}] ds \lesssim \frac{1}{\lambda_1 \lambda_2 \lambda_3 \lambda_4} \lesssim \frac{\varepsilon^{-6}}{|\xi_2|^2 |\xi_3|^2 |\xi_4|^2 |\xi_1 + \varepsilon(\xi_2 + \xi_3 + \xi_4)|^2}$$

In the end, we note that

$$\int_{\mathbb{R}^{4d}} \frac{|\hat{h}(\xi_1) \hat{f}(\xi_2) \hat{f}(\xi_3) \hat{f}(\xi_4)|}{|\xi_2|^2 |\xi_3|^2 |\xi_4|^2 |\xi_1 + \varepsilon(\xi_2 + \xi_3 + \xi_4)|^2} d\xi \lesssim 1$$

to complete the proof. \square

4.3 Proof of Proposition 2.2

Recall that

$$Y_\varepsilon = \frac{X_\varepsilon - \mathbb{E}[X_\varepsilon]}{\sqrt{\text{Var}[X_\varepsilon]}}.$$

Since

$$\begin{aligned} d_{\text{TV}}(Y_\varepsilon, \zeta) &\lesssim \mathbb{E}[\|DY_\varepsilon\|_H^4]^{1/4} \mathbb{E}[\|D^2 Y_\varepsilon\|_{\text{op}}^4]^{1/4} \\ &= \frac{1}{\text{Var}[X_\varepsilon]} \mathbb{E}[\|DX_\varepsilon\|_H^4]^{1/4} \mathbb{E}[\|D^2 X_\varepsilon\|_{\text{op}}^4]^{1/4}, \end{aligned}$$

using the fact that $\text{Var}[X_\varepsilon] \sim \varepsilon^{d-2}$ and applying Lemmas 4.1, 4.2 and 4.3, we have

$$d_{\text{TV}}(Y_\varepsilon, \zeta) \lesssim \varepsilon^{2-d} \varepsilon^{\frac{d-2}{2}-\delta} \left(\varepsilon^{4d-8-\delta} \mathbb{1}_{d < 8} + \varepsilon^{3d-8-\delta} \mathbb{1}_{d \geq 8} + \varepsilon^{3(d-2)-\delta} \right)^{\frac{1}{4}}.$$

By choosing δ small, the r.h.s. goes to zero as $\varepsilon \rightarrow 0$.

5 Proof of Theorem 1.2

The proof of Theorem 1.1 applies almost verbatim with the logarithm $\log y$ replaced by a general function $f(y)$. We need to use the assumption

$$|f(y)| + |f'(y)| + |f''(y)| \leq M(y^p + y^{-p})$$

to guarantee that

$$\mathbb{E}[|\mathbf{g}(u_\varepsilon(t, x))|^n] \lesssim 1$$

for $\mathbf{g} \in \{f, f', f''\}$, provided that β is chosen small. The only changes needed are in Section 3.3, and we sketch them here.

First, we have for general f that

$$I_{3,\varepsilon} = \int_K^{t/\varepsilon^2} \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} g(x) f'(Z(K, x/\varepsilon)) \mathbb{E} \left[\mathbb{E}_B [M_\varepsilon(t, x) \Phi_{t,x,B}^\varepsilon(s, y)] | \mathcal{F}_s \right] dx \right) dW(s, y).$$

The following two lemmas come from the same proof of Lemmas 3.5 and 3.6.

Lemma 5.1.

$$\varepsilon^{-(d-2)}\mathbb{E}[I_{3,\varepsilon}^2] = \int_0^{t-\varepsilon^2 K} \mathcal{G}_\varepsilon(s) ds,$$

with

$$\begin{aligned} \mathcal{G}_\varepsilon(s) = & \int_{\mathbb{R}^{3d}} g(x-w)g(x)R(y)\mathbb{E}\left[\mathfrak{f}'\left(Z\left(K,-\frac{w}{\varepsilon}\right)\right)\mathfrak{f}'\left(Z(K,0)\right)\right. \\ & \left. \times \mathbb{E}_B\left[M_1\left(K,-\frac{w}{\varepsilon}\right)M_2(K,0)G_{2s}(w+\varepsilon y-\varepsilon\mathbf{X}_K)\mathcal{H}\left(\frac{s}{\varepsilon^2},y,\mathbf{X}_K-\frac{w}{\varepsilon}-y\right)\right]\right] dx dy dw. \end{aligned} \quad (5.1)$$

Lemma 5.2. *There exists $\beta_0 > 0$ so that there exists $\gamma \in (0, 1)$ such that, for all $\beta < \beta_0$, $\mathcal{G}_\varepsilon(s) \lesssim s^{-\gamma}$ for $s \in (0, t)$.*

It remains to show the following lemma to complete the proof of Theorem 1.2.

Lemma 5.3. *For any $s \in (0, t)$,*

$$\mathcal{G}_\varepsilon(s) \rightarrow \nu_{\text{eff}}^2 \sigma_f^2 \int_{\mathbb{R}^{2d}} g(x-w)g(x)G_{2s}(w) dw dx,$$

as $\varepsilon \rightarrow 0$.

Proof. As in the proof of Lemma 3.7, we have

$$\mathcal{G}_\varepsilon(s) - \nu_{\text{eff}}^2 \int_{\mathbb{R}^{2d}} g(x-w)g(x)G_{2s}(w)\mathbb{E}\left[\mathfrak{f}'\left(Z\left(K,-\frac{w}{\varepsilon}\right)\right)\mathfrak{f}'\left(Z(K,0)\right)\mathbb{E}_B\left[M_1\left(K,-\frac{w}{\varepsilon}\right)M_2(K,0)\right]\right] \rightarrow 0,$$

so it remains to analyze

$$\begin{aligned} & \mathbb{E}\left[\mathfrak{f}'\left(Z\left(K,-\frac{w}{\varepsilon}\right)\right)\mathfrak{f}'\left(Z(K,0)\right)\mathbb{E}_B\left[M_1\left(K,-\frac{w}{\varepsilon}\right)M_2(K,0)\right]\right] \\ & = \mathbb{E}\left[\mathfrak{f}'\left(Z\left(K,-\frac{w}{\varepsilon}\right)\right)\mathfrak{f}'\left(Z(K,0)\right)Z\left(K,-\frac{w}{\varepsilon}\right)Z(K,0)\right] = \mathbb{E}\left[\zeta\left(K,-\frac{w}{\varepsilon}\right)\zeta(K,0)\right], \end{aligned}$$

where we defined

$$\zeta(t, x) := \mathfrak{f}'(Z(t, x))Z(t, x).$$

By stationarity in the x -variable, we write

$$\mathbb{E}\left[\zeta\left(K,-\frac{w}{\varepsilon}\right)\zeta(K,0)\right] = \text{Cov}\left[\zeta\left(K,-\frac{w}{\varepsilon}\right), \zeta(K,0)\right] + \mathbb{E}\left[\zeta(K,0)\right]^2.$$

By (2.4), we have $Z(t, 0) \rightarrow Z_\infty$ almost surely, thus, as $K \rightarrow \infty$, we have

$$\mathbb{E}\left[\zeta(K,0)\right] \rightarrow \sigma_f = \mathbb{E}\left[\mathfrak{f}'(Z_\infty)Z_\infty\right].$$

The following lemma completes the proof.

Lemma 5.4. *For any $\delta > 0$, if $\beta < \beta(\delta)$, we have for all $x \neq 0$ that*

$$\sup_{t \in [0, \varepsilon^{-2}]} \text{Cov}\left[\zeta(t, x), \zeta(t, 0)\right] \lesssim \frac{\varepsilon^{-\delta}}{|x|^{d-2}}.$$

Proof. By the covariance inequality, we have

$$|\text{Cov}[\zeta(t, x), \zeta(t, 0)]| \leq \int_{\mathbb{R}^{d+1}} \sqrt{\mathbb{E}[|D_{r,y}\zeta(t, x)|^2] \mathbb{E}[|D_{r,y}\zeta(t, 0)|^2]} dy dr.$$

Since we have

$$D_{r,y}\zeta = [\mathfrak{f}''(Z)Z + \mathfrak{f}'(Z)]D_{r,y}Z, \quad (5.2)$$

the Cauchy-Schwarz inequality yields

$$|\text{Cov}[\zeta(t, x), \zeta(t, 0)]| \leq \int_{\mathbb{R}^{d+1}} \mathbb{E}[|D_{r,y}Z(t, x)|^4]^{1/4} \mathbb{E}[|D_{r,y}Z(t, 0)|^4]^{1/4} dy dr.$$

Recalling that

$$D_{r,y}Z(t, x) = \beta \mathbb{E}_B[M(t, x) \mathbb{1}_{[0,t]}(r) \varphi(x + B_r - y)],$$

by Lemmas 3.1 and A.2 we obtain

$$\mathbb{E}[|D_{r,y}Z(t, x)|^4]^{1/4} \lesssim \mathbb{1}_{[0,t]}(r) \mathbb{E}_B[\varphi^q(x + B_r - y)]^{1/q},$$

for any $q > 1$, which implies

$$|\text{Cov}[\zeta(t, x), \zeta(t, 0)]| \lesssim \int_0^t \int_{\mathbb{R}^d} \mathbb{E}_B[\varphi^q(x + B_r - y)]^{1/q} \mathbb{E}_B[\varphi^q(B_r - y)]^{1/q} dy dr. \quad (5.3)$$

As $\varphi \in \mathcal{C}_c(\mathbb{R}^d)$, we use the simple bound

$$\mathbb{E}_B[\varphi^q(z + B_r)]^{1/q} \lesssim \mathbb{P}[|z + B_r| \leq 1]^{1/q} \lesssim \left(\mathbb{1}_{|z| \leq C} + r^{-d/2} e^{-\frac{|z|^2}{Cr}} \mathbb{1}_{|z| > C} \right)^{1/q} \lesssim \mathbb{1}_{|z| \leq C} + t^{\frac{d}{2p}} r^{-d/2} e^{-\frac{|z|^2}{qCr}} \mathbb{1}_{|z| > C},$$

for all $z \in \mathbb{R}^d$ and $r \in (0, t)$, where C is some positive constant. The r.h.s. of (5.3) is then bounded by

$$\int_0^t \int_{\mathbb{R}^d} \left(\mathbb{1}_{|x-y| \leq C} + t^{\frac{d}{2p}} r^{-d/2} e^{-\frac{|x-y|^2}{qCr}} \mathbb{1}_{|x-y| > C} \right) \left(\mathbb{1}_{|y| \leq C} + t^{\frac{d}{2p}} r^{-d/2} e^{-\frac{|y|^2}{qCr}} \mathbb{1}_{|y| > C} \right) dy dr.$$

For $|x| \gg 1$, integrating in r in the above expression to derive

$$|\text{Cov}[\zeta(t, x), \zeta(t, 0)]| \lesssim \frac{t^{\frac{d}{p}} + t^{\frac{d}{2p}}}{|x|^{d-2}}.$$

It suffices to pick $p \gg 1$ to complete the proof. \square

This also finishes the proof of Theorem 1.2. \square

Remark 5.5. By [16, Theorem 1.2], the effective variance in (1.8) is related to the asymptotic decorrelation rate of the stationary solution $\Psi(t, x)$ of the stochastic heat equation, as in (1.11):

$$\text{Cov}[\Psi(t, x), \Psi(t, y)] \approx \frac{c(\beta) \nu_{\text{eff}}^2}{|x - y|^{d-2}}, \quad \text{for } |x - y| \gg 1, \quad (5.4)$$

with $c(\beta) = \bar{c}\beta^2$. Theorem 1.2 further indicates that for any smooth function \mathfrak{f} we also have

$$\text{Cov}[\mathfrak{f}(\Psi(t, x)), \mathfrak{f}(\Psi(t, y))] \approx \frac{\sigma_{\mathfrak{f}}^2 c(\beta) \nu_{\text{eff}}^2}{|x - y|^{d-2}}, \quad \text{for } |x - y| \gg 1, \quad (5.5)$$

with $\sigma_{\mathfrak{f}} = \mathbb{E}[\mathfrak{f}'(\Psi(t, x))\Psi(t, x)]$, so that

$$\text{Cov}[\mathfrak{f}(\Psi(t, x)), \mathfrak{f}(\Psi(t, y))] \approx \sigma_{\mathfrak{f}}^2 \text{Cov}[\Psi(t, x), \Psi(t, y)] \text{ for } |x - y| \gg 1. \quad (5.6)$$

Recall also that

$$\mathbb{E}[\Psi(t, x)] = \mathbb{E}[Z_{\infty}] = 1. \quad (5.7)$$

We restate these properties in terms of the stationary solution $H(t, x) = \log \Psi(t, x)$ of the KPZ equation:

$$\text{Cov}[e^{H(t, x)}, e^{H(t, y)}] \approx \frac{c(\beta)\nu_{\text{eff}}^2}{|x - y|^{d-2}}, \text{ for } |x - y| \gg 1, \quad (5.8)$$

and

$$\text{Cov}[\mathfrak{f}(e^{H(t, x)}), \mathfrak{f}(e^{H(t, y)})] \approx \frac{\sigma_{\mathfrak{f}}^2 c(\beta)\nu_{\text{eff}}^2}{|x - y|^{d-2}} \approx \sigma_{\mathfrak{f}}^2 \text{Cov}[e^{H(t, x)}, e^{H(t, y)}], \text{ for } |x - y| \gg 1, \quad (5.9)$$

with $\sigma_{\mathfrak{f}} = \mathbb{E}[\mathfrak{f}'(e^{H(t, x)})e^{H(t, x)}]$, and

$$\mathbb{E}[e^{H(t, x)}] = 1. \quad (5.10)$$

We can now illustrate the origin of $\sigma_{\mathfrak{f}}$ through a toy calculation. Let X and Y be two jointly Gaussian $N(0, 1)$ variables, with $\text{Cov}[X, Y] = \delta \ll 1$, and define $\mathcal{X} = e^{X - \frac{1}{2}}$, $\mathcal{Y} = e^{Y - \frac{1}{2}}$, so that

$$\mathbb{E}[\mathcal{X}] = \mathbb{E}[\mathcal{Y}] = 1. \quad (5.11)$$

We think of \mathcal{X} as representing $e^{H(t, x)}$ and \mathcal{Y} as representing $e^{H(t, y)}$, so as to fit (5.10), although we emphasize that there is no real claim of Gaussianity of $H(t, \cdot)$ in the pointwise sense. With this approximation, we may write

$$Y = \delta X + \sqrt{1 - \delta^2}W,$$

with W another $N(0, 1)$ variable independent of X . Then, first, we have

$$\text{Cov}[\mathcal{X}, \mathcal{Y}] = \mathbb{E}[\mathcal{X}\mathcal{Y}] - \mathbb{E}[\mathcal{X}]\mathbb{E}[\mathcal{Y}] = e^{\delta} - 1 = \delta + o(\delta),$$

and second, we have

$$\begin{aligned} \text{Cov}[\mathfrak{f}(\mathcal{X}), \mathfrak{f}(\mathcal{Y})] &= \mathbb{E}[\mathfrak{f}(\mathcal{X})\mathfrak{f}(\mathcal{Y})] - \mathbb{E}[\mathfrak{f}(\mathcal{X})]\mathbb{E}[\mathfrak{f}(\mathcal{Y})] = \delta \mathbb{E}[\mathfrak{f}'(e^{X - \frac{1}{2}})X] \mathbb{E}[\mathfrak{f}'(e^{W - \frac{1}{2}})e^{W - \frac{1}{2}}] + o(\delta) \\ &= \delta \sigma_{\mathfrak{f}}^2 + o(\delta). \end{aligned} \quad (5.12)$$

We have denoted $\sigma_{\mathfrak{f}} = \mathbb{E}[\mathfrak{f}'(\mathcal{X})\mathcal{X}]$ for smooth functions \mathfrak{f} , as before. Here, we used the identity

$$\mathbb{E}[\mathfrak{f}(e^{X - \frac{1}{2}})X] = \mathbb{E}[\mathfrak{f}'(e^{X - \frac{1}{2}})e^{X - \frac{1}{2}}], \quad (5.13)$$

obtained via integration by parts. Thus, we have

$$\text{Cov}[\mathfrak{f}(\mathcal{X}), \mathfrak{f}(\mathcal{Y})] \approx \sigma_{\mathfrak{f}}^2 \text{Cov}[\mathcal{X}, \mathcal{Y}]. \quad (5.14)$$

Unravelling the definitions, this parallels (5.9).

As emphasized above, there was nothing Gaussian in the pointwise sense in the field $H(t, x)$. However, the above computation would be essentially unchanged if we replace $H(t, x)$ by the field

$$\tilde{H}(t, x) = \mathcal{G}(t, x) + E(t, x), \quad (5.15)$$

where $\mathcal{G}(t, \cdot)$ is a mean-zero spatially stationary Gaussian random field with $\mathbb{E}[\mathcal{G}(t, x)^2] = \sigma^2$, and correlation function

$$\mathbb{E}[\mathcal{G}(t, x)\mathcal{G}(t, y)] \approx \frac{\sigma^2}{|x - y|^{d-2}}, \text{ for } |x - y| \gg 1, \quad (5.16)$$

while $E(t, \cdot)$ is a spatial stationary random field of negative mean, independent of $\mathcal{G}(t, \cdot)$, with a rapidly (in space) decaying correlation function, such that

$$\mathbb{E}[e^{\mathcal{G}(t, x)}]\mathbb{E}[e^{E(t, x)}] = 1. \quad (5.17)$$

(This would be consistent with (5.8)-(5.9).) Repeating the above computation with $x, y \in \mathbb{R}^d$ such that $|x - y| \gg 1$ and $\delta = \text{Cov}[\mathcal{G}(t, x), \mathcal{G}(t, y)] \ll 1$, we then obtain again that

$$\text{Cov}[\mathfrak{f}(e^{\tilde{H}(t, x)}), \mathfrak{f}(e^{\tilde{H}(t, y)})] \approx \sigma_{\mathfrak{f}}^2 \text{Cov}[e^{\tilde{H}(t, x)}, e^{\tilde{H}(t, y)}]. \quad (5.18)$$

Thus, any field of the form (5.15) satisfies conditions (5.8)-(5.10), with the variance $\sigma_{\mathfrak{f}}$ as in Theorem 1.2. It is tempting to speculate that the stationary solution $H(t, x)$ of the KPZ equation has a decomposition of the form (5.15); we unfortunately do not have any supporting evidence for such a speculation, and at this point cannot even agree whether such a decomposition is behind the results of this paper or, alternatively, that temporal mixing plays an additional important role.

A Auxiliary lemmas

Lemma A.1. *For any $t > 0, x \in \mathbb{R}^d$, we have*

$$D \log Z(t, x) = \frac{DZ(t, x)}{Z(t, x)} \in L^2(\Omega; H).$$

Proof. Recall that

$$\begin{aligned} Z(t, x) &= \mathbb{E}_B \left[\exp \left\{ \beta \int_0^t V(s, x + B_s) ds - \frac{1}{2} \beta^2 R(0)t \right\} \right], \\ \int_0^t V(s, x + B_s) ds &= \int_{\mathbb{R}^{d+1}} \mathbb{1}_{[0, t]}(s) \varphi(x + B_s - y) dW(s, y), \end{aligned}$$

so, for each t and x fixed, we have

$$D_{s, y} Z(t, x) = \beta \mathbb{E}_B \left[\exp \left\{ \beta \int_0^t V(s, x + B_s) ds - \frac{1}{2} \beta^2 R(0)t \right\} \mathbb{1}_{[0, t]}(s) \varphi(x + B_s - y) \right] \in L^n(\Omega; H), \quad (\text{A.1})$$

for any $n \in \mathbb{Z}_+$, where we recall that H is the $L^2(\mathbb{R}^{d+1})$ -space with respect to the s, y -variables. To deal with the logarithm function, which is singular at the origin and grows at infinity, we use an approximation $f_n \in C_c^\infty(\mathbb{R})$ such that $f_n(x) = \log x$ for $x \in [1/n, n]$ and $|f'_n(x)| \leq |x|^{-1}$. It is clear that

$$Df_n(Z(t, x)) = f'_n(Z(t, x)) DZ(t, x) \in L^2(\Omega; H),$$

and the error

$$\begin{aligned} \mathbb{E} \left[\left\| Df_n(Z(t, x)) - \frac{DZ(t, x)}{Z(t, x)} \right\|_H^2 \right] &\lesssim \mathbb{E} \left[\frac{\|DZ(t, x)\|_H^2}{|Z(t, x)|^2} (\mathbb{1}_{Z(t, x) < \frac{1}{n}} + \mathbb{1}_{Z(t, x) > n}) \right] \\ &\lesssim \sqrt{\mathbb{E}[|Z(t, x)|^{-4} (\mathbb{1}_{Z(t, x) < \frac{1}{n}} + \mathbb{1}_{Z(t, x) > n})]} \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$, where we used Proposition 2.3 in the last step, together with (A.1). By [34, Proposition 1.2.1], the proof is complete. \square

Lemma A.2. *There exists $\beta_0 > 0$ such that if $\beta < \beta_0$, we have in $d \geq 3$ that*

$$\sup_{x \in \mathbb{R}^d} \mathbb{E}_B \left[\exp \left\{ \beta \int_0^\infty R(x + B_s) ds \right\} \right] < \infty, \quad (\text{A.2})$$

and

$$\sup_{t > 0, x, y \in \mathbb{R}^d} \mathbb{E}_B \left[\exp \left\{ \beta \int_0^t R(x + B_s) ds \right\} \middle| B_t = y \right] < \infty. \quad (\text{A.3})$$

Proof. The statement in (A.2) follows from Portenko's lemma, see [32, (3.1), (3.2)].

We turn to proving (A.3). Conditioned on $B_t = y$, the process $\{B_s\}_{s \leq t}$ is a Brownian bridge, and solves the SDE

$$dX_s = -\frac{X_s - y}{t - s} ds + d\bar{B}_s, \quad X_0 = 0, \quad (\text{A.4})$$

where \bar{B} is a standard Brownian motion. (See [36, Exercise IV.3.18], or simply compute the covariance.) In particular, it has a Markovian representation. Thus, again by Portenko's lemma, it is enough to show that

$$\beta \sup_{t > 0, x, y \in \mathbb{R}^d} \mathbb{E}_B \left[\int_0^t R(x + B_s) ds \middle| B_t = y \right] < 1, \quad (\text{A.5})$$

for all β small enough. By symmetry, as X_s is a Brownian bridge, it suffices to show that

$$\sup_{t > 0, x, y \in \mathbb{R}^d} \mathbb{E}_B \left[\int_0^{t/2} R(x + B_s) ds \middle| B_t = y \right] < \infty. \quad (\text{A.6})$$

Note that X_s has mean sy/t and variance $s(t-s)/t$, which in our range is larger than Cs . In particular, for $s \leq t/2$, we have

$$\sup_{x, y \in \mathbb{R}^d} \mathbb{P}_B \left[|B_s - x| \leq 1 \middle| B_t = y \right] \leq C(1+s)^{-d/2}.$$

Since $s^{-d/2}$ is integrable as $s \rightarrow +\infty$, this yields (A.6) and completes the proof of the lemma. \square

Lemma A.3. *Let B be a standard Brownian motion in $d \geq 3$. For any $\alpha \in (0, 2)$ and $t > 0, w \neq 0$, we have*

$$\mathbb{P}_B \left[\min_{\ell \in [t/\varepsilon^\alpha, t/\varepsilon^2]} |B_\ell| \leq 1 \middle| \varepsilon B_{t/\varepsilon^2} = w \right] \rightarrow 0$$

as $\varepsilon \rightarrow 0$.

Proof. It is convenient to handle separately the intervals $\ell \in [t/\varepsilon^\alpha, t/2\varepsilon^2]$, $\ell \in [t/2\varepsilon^2, t/\varepsilon^2 - t/\varepsilon^\gamma]$, and $\ell \in [t/\varepsilon^2 - t/\varepsilon^\gamma, t/\varepsilon^2]$, with $\gamma \in (4/d, 2)$.

In the regime $\ell \in [t/\varepsilon^\alpha, t/2\varepsilon^2]$, note that it follows from the representation (A.4) that the law of the process $\{X_s\}_{s \in [0, r]}$ is absolutely continuous with respect to the law of the process $\{B_s\}_{s \in [0, r]}$, with Radon-Nykodim derivative given by

$$\Lambda_t(r) = \exp \left\{ - \int_0^r \frac{X_s - w/\varepsilon}{t/\varepsilon^2 - s} \cdot dX_s - \frac{1}{2} \int_0^r \frac{|X_s - w/\varepsilon|^2}{(t/\varepsilon^2 - s)^2} ds \right\} = \exp \left\{ - \frac{|X_r - w/\varepsilon|^2}{2(t/\varepsilon^2 - r)} + \frac{w^2}{2t} + \frac{d}{2} \int_0^r \frac{ds}{t/\varepsilon^2 - s} \right\}, \quad (\text{A.7})$$

where the second equality follows by integration by parts. Thus, for $r \in [t/\varepsilon^\alpha, t/2\varepsilon^2]$, we have the bound $\Lambda_t(r) \leq \Lambda_t(t/2\varepsilon^2) \leq C$ where $C = C(w)$ is a universal constant. Therefore, we have

$$\mathbb{P}_B \left[\min_{\ell \in [t/\varepsilon^\alpha, t/2\varepsilon^2]} |B_\ell| \leq 1 \mid \varepsilon B_{t/\varepsilon^2} = w \right] \leq C \mathbb{P}_B \left[\min_{\ell \in [t/\varepsilon^\alpha, \infty]} |B_\ell| \leq 1 \right] \xrightarrow{\varepsilon \rightarrow 0} 0, \quad (\text{A.8})$$

where the limit is obtained from integrating the heat kernel with respect to time over $[t/\varepsilon^\alpha, \infty)$, using $d \geq 3$.

We argue similarly in the regime $\ell \in [t/2\varepsilon^2, t/\varepsilon^2 - t/\varepsilon^\gamma]$. Indeed, in that regime we have from (A.7) that $\Lambda_t(r) \leq \Lambda_t(t/\varepsilon^2 - t/\varepsilon^\gamma) \leq \varepsilon^{-(2-\gamma)d/2}$, and thus

$$\mathbb{P}_B \left[\min_{\ell \in [t/2\varepsilon^2, t/\varepsilon^2 - t/\varepsilon^\gamma]} |B_\ell| \leq 1 \mid \varepsilon B_{t/\varepsilon^2} = w \right] \leq C \varepsilon^{-(2-\gamma)d/2} \mathbb{P}_B \left[\min_{\ell \in [t/2\varepsilon^2, \infty)} |B_\ell| \leq 1 \right]. \quad (\text{A.9})$$

A Green function computation gives immediately that

$$\mathbb{P}_B \left[\min_{\ell \in [t/2\varepsilon^2, \infty)} |B_\ell| \leq 1 \right] \leq C \varepsilon^{d-2}.$$

Using this in (A.9), we conclude that the left side there goes to zero as long as $\gamma > 4/d$.

For the regime $\ell \in [t/\varepsilon^2 - t/\varepsilon^\gamma, t/\varepsilon^2]$ we use a different representation of the Brownian bridge: if \mathcal{B}_s is a Brownian bridge on $[0, t/\varepsilon^2]$ with $\mathcal{B}_{t/\varepsilon^2} = w/\varepsilon$ then

$$\mathcal{B}_s = \frac{s\varepsilon^2}{t} (w/\varepsilon - \mathcal{B}_{t/\varepsilon^2}) + B_s,$$

for some Brownian motion B . In particular, this gives a modulus of continuity for the Brownian bridge, of the following form. Let $t_0 = t/\varepsilon^2 - t/\varepsilon^\gamma$ and let $t_i = t_0 + \frac{i}{A \log(1/\varepsilon)}$, where A is a large constant to be determined below, and set

$$M_\varepsilon = \max_{i: t_{i-1} \leq t/\varepsilon^2} \max_{s \in [0, 1/A \log(1/\varepsilon)]} |\mathcal{B}_{s+t_i} - \mathcal{B}_{t_i}|. \quad (\text{A.10})$$

Note that

$$\mathcal{B}_{s+t_i} - \mathcal{B}_{t_i} = \frac{s\varepsilon w}{t} - \frac{s\varepsilon^2}{t} B_{t/\varepsilon^2} + (B_{s+t_i} - B_{t_i}).$$

Since t, w are fixed, we have $s\varepsilon w/t \rightarrow 0$ as $\varepsilon \rightarrow 0$ for $s \leq \frac{1}{A \log(1/\varepsilon)}$, and if we further assume that $B_{t/\varepsilon^2} \leq t/4\varepsilon^2$, then

$$\max_{s \in [0, 1/A \log(1/\varepsilon)]} \left| \frac{s\varepsilon w}{t} - \frac{s\varepsilon^2}{t} B_{t/\varepsilon^2} \right| < \frac{1}{2}.$$

Therefore, we have

$$\begin{aligned} \mathbb{P}_B[M_\varepsilon \geq 1] &\leq \mathbb{P}_B[B_{t/\varepsilon^2} > t/4\varepsilon^2] + \frac{At \log(1/\varepsilon)}{\varepsilon^2} \mathbb{P}_B \left[\max_{s \in [0, 1/A \log(1/\varepsilon)]} |B_s| > 1/2 \right] \\ &\leq C e^{-c/\varepsilon} + CA \varepsilon^{-2} \log(1/\varepsilon) e^{-A |\log \varepsilon|} \xrightarrow{\varepsilon \rightarrow 0} 0, \end{aligned} \quad (\text{A.11})$$

for A large enough (independent of ε). We now have

$$\mathbb{P}_B \left[\min_{\ell \in [t/2\varepsilon^2, t/\varepsilon^2]} |B_\ell| \leq 1 \mid \varepsilon B_{t/\varepsilon^2} = w \right] = \mathbb{P}_B \left[\min_{\ell \in [t/2\varepsilon^2, t/\varepsilon^2]} |B_\ell| \leq 1 \right] \leq \mathbb{P}_B[M_\varepsilon \geq 1] + \sum_{i: t_{i-1} < t/\varepsilon^2} \mathbb{P}_B[|\mathcal{B}_{t_i}| \leq 2]. \quad (\text{A.12})$$

The mean of \mathcal{B}_{t_i} is $\varepsilon t_i w/t$ which is greater in modulus than $|w|/2\varepsilon$ (for $t_i > \frac{t}{2\varepsilon^2}$), and its variance is $t_i(t - \varepsilon^2 t_i)/t \leq t/\varepsilon^\gamma$. It follows that

$$\mathbb{P}_B [|\mathcal{B}_{t_i}| \leq 2] \leq C e^{-C|w|^2 \varepsilon^{\gamma-2}}.$$

Thus, the summation term in (A.12) goes to zero as $\varepsilon \rightarrow 0$ since $\gamma < 2$. \square

B Negative moments of $Z(t, x)$

We now prove Proposition 2.3. The goal is to show there exists $\beta_0 > 0$ such that if $\beta < \beta_0$ and $n \in \mathbb{Z}_+$, we have

$$\sup_{t>0} \mathbb{E}[Z(t, x)^{-n}] \leq C_{\beta, n}, \quad (\text{B.1})$$

with some constant $C_{\beta, n} > 0$. We adapt to our setting the proof of [24, Corollary 4.8], which deals with the case when the noise is also singular in space. The same proof applies to our situation, and we only present the details for the convenience of the readers.

Since $Z(t, x)$ has the same distribution as $u(t, x)$, it suffices to estimate the small ball probability $\mathbb{P}[u(t, x) \leq r]$ for $r \ll 1$. We define an approximation of the spacetime white noise

$$W_\varepsilon(t, x) = e^{-\varepsilon(t^2 + |x|^2)} \int_{\mathbb{R}^{d+1}} \phi_\varepsilon(t-s, x-y) dW(s, y),$$

where $\phi_\varepsilon(t, x) = \varepsilon^{-d-2} \phi(t/\varepsilon^2, x/\varepsilon)$ with $\phi \in \mathcal{C}_c^\infty(\mathbb{R}^{d+1})$ such that $\phi \geq 0$ is even and $\|\phi\|_{L^1} = 1$. It is clear that for fixed $\varepsilon > 0$, $W_\varepsilon \in L^2(\mathbb{R}^{d+1}) \cap \mathcal{C}^\infty(\mathbb{R}^{d+1})$ almost surely. We will use $\|\cdot\|_2$ to denote the $L^2(\mathbb{R}^{d+1})$ norm. Define

$$V_\varepsilon(t, x) = \int_{\mathbb{R}^d} \varphi(x-y) W_\varepsilon(t, y) dy, \quad \mathcal{R}_\varepsilon(t, s, x, y) = \mathbb{E}[V_\varepsilon(t, x) V_\varepsilon(s, y)],$$

and

$$\mathcal{U}_\varepsilon(t, x) = \mathbb{E}_B [e^{\mathcal{V}_t^\varepsilon(B)}],$$

with

$$\mathcal{V}_t^\varepsilon(B) = \beta \int_0^t V_\varepsilon(t-s, x+B_s) ds - \frac{1}{2} \beta^2 \mathcal{Q}_\varepsilon(t, x, x, B, B),$$

where

$$\mathcal{Q}_\varepsilon(t, x, y, B^1, B^2) = \int_0^t \int_0^t \mathcal{R}_\varepsilon(t-s, t-\ell, x+B_s^1, y+B_\ell^2) ds d\ell.$$

By [24, Proposition 4.2], $\mathcal{U}_\varepsilon(t, x) \rightarrow u(t, x)$ in probability so we only need to estimate $\mathbb{P}[\mathcal{U}_\varepsilon(t, x) \leq r]$ for $r \ll 1$.

With any given W_ε , define the expectation

$$\mathbb{E}_B^{W_\varepsilon} [F(B^1, B^2)] = \frac{\mathbb{E}_B [F(B^1, B^2) e^{\mathcal{V}_t^\varepsilon(B^1) + \mathcal{V}_t^\varepsilon(B^2)}]}{\mathbb{E}_B [e^{\mathcal{V}_t^\varepsilon(B^1) + \mathcal{V}_t^\varepsilon(B^2)}]}.$$

To emphasize the dependence of \mathcal{U}_ε on W_ε , we write $\mathcal{U}_\varepsilon(t, x) = \mathcal{U}_\varepsilon(t, x, W_\varepsilon)$. For any $\lambda > 0$, define the set

$$A_\lambda(t, x) = \left\{ W_\varepsilon : \mathcal{U}_\varepsilon(t, x, W_\varepsilon) > \frac{1}{2}, \int_0^t \mathbb{E}_B^{W_\varepsilon} [R(B_s^1 - B_s^2)] ds \leq \lambda \right\}.$$

Lemma B.1. For any $\tilde{W}_\varepsilon \in A_\lambda(t, x)$, we have

$$\mathcal{U}_\varepsilon(t, x, W_\varepsilon) \geq \frac{1}{2} e^{-\sqrt{\lambda} \|W_\varepsilon - \tilde{W}_\varepsilon\|_2},$$

with $\|\cdot\|_2$ denoting the $L^2(\mathbb{R}^{d+1})$ norm.

Proof. We write

$$\begin{aligned} \mathcal{U}_\varepsilon(t, x, W_\varepsilon) &= \mathbb{E}_B[e^{\mathcal{V}_t^\varepsilon(B)}] = \mathbb{E}_B[e^{\tilde{\mathcal{V}}_t^\varepsilon(B)}] \frac{\mathbb{E}_B[e^{\mathcal{V}_t^\varepsilon(B) - \tilde{\mathcal{V}}_t^\varepsilon(B)} e^{\tilde{\mathcal{V}}_t^\varepsilon(B)}]}{\mathbb{E}_B[e^{\tilde{\mathcal{V}}_t^\varepsilon(B)}]} \\ &= \mathcal{U}_\varepsilon(t, x, \tilde{W}_\varepsilon) \mathbb{E}_B^{\tilde{W}_\varepsilon}[e^{\mathcal{V}_t^\varepsilon(B) - \tilde{\mathcal{V}}_t^\varepsilon(B)}], \end{aligned}$$

where $\tilde{\mathcal{V}}_t^\varepsilon$ is obtained by replacing $W_\varepsilon \mapsto \tilde{W}_\varepsilon$ in the expression of $\mathcal{V}_t^\varepsilon$. Since $\tilde{W}_\varepsilon \in A_\lambda$, by Jensen's inequality we have

$$\mathcal{U}_\varepsilon(t, x, W_\varepsilon) \geq \frac{1}{2} \exp\left(\mathbb{E}_B^{\tilde{W}_\varepsilon}[\mathcal{V}_t^\varepsilon(B) - \tilde{\mathcal{V}}_t^\varepsilon(B)]\right).$$

It remains to show that

$$|\mathbb{E}_B^{\tilde{W}_\varepsilon}[\mathcal{V}_t^\varepsilon(B) - \tilde{\mathcal{V}}_t^\varepsilon(B)]| \leq \sqrt{\lambda} \|W_\varepsilon - \tilde{W}_\varepsilon\|_2. \quad (\text{B.2})$$

We write

$$\begin{aligned} \mathcal{V}_t^\varepsilon(B) - \tilde{\mathcal{V}}_t^\varepsilon(B) &= \beta \int_0^t [V_\varepsilon(t-s, x+B_s) - \tilde{V}_\varepsilon(t-s, x+B_s)] ds \\ &= \beta \int_0^t \int_{\mathbb{R}^d} \varphi(x+B_s-y) [W_\varepsilon(t-s, y) - \tilde{W}_\varepsilon(t-s, y)] dy ds, \end{aligned}$$

and apply the Cauchy-Schwarz inequality to get

$$\begin{aligned} |\mathbb{E}_B^{\tilde{W}_\varepsilon}[\mathcal{V}_t^\varepsilon(B) - \tilde{\mathcal{V}}_t^\varepsilon(B)]| &\leq \beta \|W_\varepsilon - \tilde{W}_\varepsilon\|_2 \sqrt{\int_0^t \int_{\mathbb{R}^d} |\mathbb{E}_B^{\tilde{W}_\varepsilon}[\varphi(x+B_s-y)]|^2 dy ds} \\ &\leq \beta \|W_\varepsilon - \tilde{W}_\varepsilon\|_2 \sqrt{\int_0^t \mathbb{E}_B^{\tilde{W}_\varepsilon}[R(B_s^1 - B_s^2)] ds} \leq \sqrt{\lambda} \|W_\varepsilon - \tilde{W}_\varepsilon\|_2, \end{aligned}$$

which completes the proof. \square

Lemma B.2. There exists universal constants $\lambda, c > 0$ such that $\mathbb{P}[A_\lambda(t, x)] \geq c$.

Proof. We have

$$\mathbb{P}[A_\lambda(t, x)] \geq \mathbb{P}[\mathcal{U}_\varepsilon(t, x, W_\varepsilon) > \frac{1}{2}] - \mathbb{P}[B_\lambda(t, x)],$$

with

$$B_\lambda(t, x) = \left\{ W_\varepsilon : \mathcal{U}_\varepsilon(t, x, W_\varepsilon) > \frac{1}{2}, \int_0^t \mathbb{E}_B^{W_\varepsilon}[R(B_s^1 - B_s^2)] ds > \lambda \right\}.$$

Using the fact that $\mathbb{E}[\mathcal{U}_\varepsilon(t, x, W_\varepsilon)] = 1$ and the Paley-Zygmund's inequality, we have

$$\mathbb{P}\left[\mathcal{U}_\varepsilon(t, x, W_\varepsilon) > \frac{1}{2}\right] \geq \frac{1}{4\mathbb{E}[\mathcal{U}_\varepsilon(t, x, W_\varepsilon)^2]} = \frac{1}{4\mathbb{E}_B[e^{\beta^2 \mathcal{Q}_\varepsilon(t, x, x, B^1, B^2)}]}.$$

For $B_\lambda(t, x)$, we have, as $\mathcal{U}_\varepsilon(t, x, W_\varepsilon) > 1/2$,

$$\begin{aligned} \mathbb{P}[B_\lambda(t, x)] &\leq \mathbb{P}\left[\int_0^t \mathbb{E}_B[R(B_s^1 - B_s^2)] e^{\mathcal{V}_t^\varepsilon(B^1) + \mathcal{V}_t^\varepsilon(B^2)} ds > \frac{\lambda}{4}\right] \\ &\leq \frac{4}{\lambda} \mathbb{E}_B\left[e^{\beta^2 \mathcal{Q}_\varepsilon(t, x, x, B^1, B^2)} \int_0^t R(B_s^1 - B_s^2) ds\right] \leq \frac{4C}{\lambda} \mathbb{E}_B\left[e^{2\beta^2 \mathcal{Q}_\varepsilon(t, x, x, B^1, B^2)}\right]^{1/2}, \end{aligned} \quad (\text{B.3})$$

with some constant $C > 0$. By Lemma B.3 below and choosing λ large, there exists some constants $c, \lambda > 0$ independent of ε, t, x such that $\mathbb{P}[A_\lambda(t, x)] \geq c$. \square

Lemma B.3. *There exists $\beta_0 > 0$ such that if $\beta < \beta_0$, we have*

$$1 \leq \mathbb{E}_B \left[e^{\beta \mathcal{Q}_\varepsilon(t, x, x, B^1, B^2)} \right] \leq C_\beta.$$

Proof. Recall that

$$\mathcal{Q}_\varepsilon(t, x, x, B^1, B^2) = \int_0^t \int_0^t \mathcal{R}_\varepsilon(t-s, t-\ell, x+B_s^1, x+B_\ell^2) ds d\ell.$$

We write \mathcal{R}_ε explicitly:

$$\begin{aligned} \mathcal{R}_\varepsilon(t_1, t_2, x_1, x_2) &= \int_{\mathbb{R}^{2d}} \varphi(x_1 - y_1) \varphi(x_2 - y_2) \mathbb{E}[W_\varepsilon(t_1, y_1) W_\varepsilon(t_2, y_2)] dy_1 dy_2 \\ &= \int_{\mathbb{R}^{2d}} \varphi(x_1 - y_1) \varphi(x_2 - y_2) e^{-\varepsilon(t_1^2 + t_2^2 + |y_1|^2 + |y_2|^2)} \phi_\varepsilon \star \phi_\varepsilon(t_1 - t_2, y_1 - y_2) dy_1 dy_2 \\ &\leq \int_{\mathbb{R}^{2d}} \varphi(x_1 - y_1) \varphi(x_2 - y_2) \phi_\varepsilon \star \phi_\varepsilon(t_1 - t_2, y_1 - y_2) dy_1 dy_2, \end{aligned}$$

with \star denoting the convolution. By the fact that φ, ϕ have compact supports, we have

$$\mathcal{R}_\varepsilon(t_1, t_2, x_1, x_2) \lesssim \varepsilon^{-2} \mathbb{1}_{|x_1 - x_2| \leq C, |t_1 - t_2| \leq C\varepsilon^2}.$$

for some $C > 0$. Thus, \mathcal{Q}_ε is essentially measuring the mutual “intersection” time of B^1, B^2 . By [18, Corollary 4.4] and the fact that $d \geq 3$, the proof is complete. \square

Now we can write

$$\mathbb{P}[\mathcal{U}_\varepsilon(t, x, W_\varepsilon) \leq r] \leq \mathbb{P}\left[\frac{1}{2} e^{-\sqrt{\lambda} \text{dist}(W_\varepsilon, A_\lambda(t, x))} \leq r\right] \leq \mathbb{P}\left[\text{dist}(W_\varepsilon, A_\lambda(t, x)) \geq \frac{\log(2r)^{-1}}{\sqrt{\lambda}}\right], \quad (\text{B.4})$$

where $\text{dist}(W_\varepsilon, A_\lambda(t, x)) = \inf\{\|W_\varepsilon - \tilde{W}_\varepsilon\|_2 : \tilde{W}_\varepsilon \in A_\lambda(t, x)\}$. Now we can apply [24, Lemma 4.5] to obtain

$$\mathbb{P}\left[\text{dist}(W_\varepsilon, A_\lambda(t, x)) \geq \tau + 2\sqrt{\log \frac{2}{c}}\right] \leq 2e^{-\tau^2/4} \quad (\text{B.5})$$

for all $\tau > 0$, where $\lambda, c > 0$ are chosen as in Lemma B.2. Combining (B.4) and (B.5), we have

$$\mathbb{P}[\mathcal{U}_\varepsilon(t, x, W_\varepsilon) \leq r] \leq 2 \exp\left(-\frac{1}{4} \left(\frac{\log(2r)}{\sqrt{\lambda}} + 2\sqrt{\log \frac{2}{c}}\right)^2\right),$$

which implies $\mathbb{E}[\mathcal{U}_\varepsilon(t, x, W_\varepsilon)^{-n}] \lesssim 1$ and completes the proof of (B.1).

References

- [1] T. ALBERTS, K. KHANIN, AND J. QUASTEL, The intermediate disorder regime for directed polymers in dimension $1+1$, Ann. Probab., **42** (2014), pp. 1212–1256. [3](#)

- [2] G. AMIR, I. CORWIN, AND J. QUASTEL, Probability distribution of the free energy of the continuum directed random polymer in $1+1$ dimensions, *Comm. Pure Appl. Math.*, **64** (2011), pp. 466–537. [3](#)
- [3] L. BERTINI AND G. GIACOMIN, Stochastic Burgers and KPZ equations from particle systems, *Comm. Math. Phys.*, **183** (1997), pp. 571–607. [3](#)
- [4] L. BERTINI AND N. CANCRINI, The two-dimensional stochastic heat equation: renormalizing a multiplicative noise, *Journal of Physics A: Mathematical and General*, **31** (1998), p. 615. [3](#)
- [5] V. BETZ AND H. SPOHN, A central limit theorem for Gibbs measures relative to Brownian motion, *Prob. Theory Rel. Fields*, **131** (2005), pp. 459–478. [5](#)
- [6] E. BOLTHAUSEN, A note on the diffusion of directed polymers in a random environment, *Comm. Math. Phys.*, **123** (1989), pp. 529–534. [3](#)
- [7] F. CARAVENNA, R. SUN, AND N. ZYGOURAS, Universality in marginally relevant disordered systems, *Ann. Appl. Prob.*, **27** (2017), pp. 3050–3112. [3](#)
- [8] F. CARAVENNA, R. SUN AND N. ZYGOURAS, On the moments of the $(2+1)$ -dimensional directed polymer and stochastic heat equation in the critical window, arXiv:1808.03586 (2018). [3](#)
- [9] F. CARAVENNA, R. SUN AND N. ZYGOURAS, The two-dimensional KPZ equation in the entire subcritical regime, arXiv:1812.03911 (2018). [3](#)
- [10] S. CHATTERJEE AND A. DUNLAP, Constructing a solution of the $(2+1)$ -dimensional KPZ equation, arXiv preprint arXiv:1809.00803, (2018). [3](#)
- [11] S. CHATTERJEE, Fluctuations of eigenvalues and second order Poincaré inequalities, *Probability Theory and Related Fields*, **143** (2009), pp. 1–40. [6](#)
- [12] F. COMETS, C. COSCO AND C. MUKHERJEE, Fluctuation and rate of convergence for the stochastic heat equation in weak disorder, arXiv preprint arXiv:1807.03902, (2018). [5](#), [10](#)
- [13] F. COMETS, Directed polymers in random environments, *Lecture Notes in Mathematics*, vol. 2175, Springer, Cham, 2017, Lecture notes from the 46th Probability Summer School held in Saint-Flour, 2016. [3](#)
- [14] F. COMETS AND Q. LIU, Rate of convergence for polymers in a weak disorder, *J. Math. Anal. Appl.*, **455** (2017), pp. 312–335. [5](#)
- [15] I. CORWIN, The Kardar–Parisi–Zhang equation and universality class, *Random matrices: Theory and applications*, **1** (2012), p. 1130001. [3](#)
- [16] A. DUNLAP, Y. GU, L. RYZHIK, AND O. ZEITOUNI, The random heat equation in dimensions three and higher: the homogenization viewpoint, arXiv preprint arXiv:1808.07557, (2018). [3](#), [22](#)
- [17] Z. S. FENG, Rescaled Directed Random Polymer in Random Environment in Dimension $1+2$, PhD thesis, University of Toronto (Canada), 2016. [3](#)
- [18] Y. GU, L. RYZHIK, AND O. ZEITOUNI, The Edwards–Wilkinson limit of the random heat equation in dimensions three and higher, *Comm. Math. Phys.*, **363** (2018), pp. 351–388. [3](#), [4](#), [5](#), [29](#)

- [19] M. GUBINELLI, Gibbs measures for self-interacting Wiener paths, *Markov Proc. Rel. Fields*, **12** (2006), pp. 747–766. [5](#)
- [20] M. GUBINELLI, P. IMKELLER, AND N. PERKOWSKI, Paracontrolled distributions and singular PDEs, in *Forum of Mathematics, Pi*, vol. 3, Cambridge Univ Press, 2015, p. e6. [3](#)
- [21] M. GUBINELLI AND N. PERKOWSKI, KPZ reloaded, *Communications in Mathematical Physics*, **349** (2017), pp. 165–269. [3](#)
- [22] M. HAIRER, Solving the KPZ equation, *Ann. Math.*, **178** (2013), pp. 559–664. [3](#)
- [23] ———, A theory of regularity structures, *Inv. Math.*, **198** (2014), pp. 269–504. [3](#)
- [24] Y. HU AND K. LÊ, Asymptotics of the density of parabolic Anderson random fields, arXiv preprint arXiv:1801.03386, (2018). [27](#), [29](#)
- [25] J. Z. IMBRIE AND T. SPENCER, Diffusion of directed polymers in a random environment, *Jour. Stat. Phys.*, **52** (1988), pp. 609–626. [3](#)
- [26] A. KUPIAINEN, Renormalization group and stochastic PDEs, in *Annales Henri Poincaré*, vol. 17, Springer, 2016, pp. 497–535. [3](#)
- [27] J. MAGNEN AND J. UNTERBERGER, The Scaling Limit of the KPZ Equation in Space Dimension 3 and Higher, *Jour. Stat. Phys.*, **171** (2018), pp. 543–598. [1](#), [3](#), [5](#)
- [28] K. MATETSKI, J. QUASTEL, AND D. REMENIK, The KPZ fixed point, arXiv preprint arXiv:1701.00018, (2016). [3](#)
- [29] E. MAYER-WOLF AND M. ZAKAI, The Clark-Ocone formula for vector valued Wiener functionals, *J. Funct. Anal.*, **229** (2005), pp. 143–154. [7](#)
- [30] C. MUKHERJEE, A central limit theorem for the annealed path measures for the stochastic heat equation and the continuous directed polymer in $d \geq 3$, arXiv preprint arXiv:1706.09345, (2017). [5](#)
- [31] C. MUKHERJEE AND Y. BROEKER, Quenched central limit theorem for the stochastic heat equation in weak disorder,, arXiv preprint arXiv:1710.00631, (2017). [3](#)
- [32] C. MUKHERJEE, A. SHAMOV, AND O. ZEITOUNI, Weak and strong disorder for the stochastic heat equation and continuous directed polymers in $d \geq 3$, *Electr. Comm. Prob.*, **21** (2016). [3](#), [6](#), [25](#)
- [33] I. NOURDIN, G. PECCATI, AND G. REINERT, Second order Poincaré inequalities and CLTs on Wiener space, *Journal of Functional Analysis*, **257** (2009), pp. 593–609. [6](#), [8](#), [16](#)
- [34] D. NUALART, *The Malliavin calculus and related topics*, vol. 1995, Springer, 2006. [24](#)
- [35] J. QUASTEL AND H. SPOHN, The one-dimensional KPZ equation and its universality class, *Journal of Statistical Physics*, **160** (2015), pp. 965–984. [3](#)
- [36] D. REVUZ AND M. YOR, *Continuous martingales and Brownian motion*, third edition, Springer (1999). [25](#)
- [37] F. TONINELLI, $(2+1)$ -dimensional interface dynamics: mixing time, hydrodynamic limit and anisotropic KPZ growth, arXiv preprint arXiv:1711.05571, (2017). [3](#)