## CA LECTURE 3

## SCRIBE: CLAIRE TOMESCH

Recall that R/I is an ID iff I is prime and R/I is a field iff I is maximal.

**Definition 1.** A poset is  $(\mathbb{P}, \leq)$  where  $\leq$  is transitive  $(a \leq b \text{ and } b \leq c \text{ implies } a \leq c)$ , reflexive  $(a \leq a)$  and antisymmetric  $(a \leq b \text{ and } b \leq a \text{ implies } a = b)$ .

It is easy to see that any family of sets ordered by inclusion is a poset. Conversely if  $\mathbb{P}$  is a poset and we define  $C_a = \{b : b \leq a\}$  then the set of  $C_a$  ordered by inclusion forms a poset isomorphic to  $\mathbb{P}$ .

A maximal element in  $\mathbb{P}$  is an element q such that  $\forall r \ r \geq q \implies r = q$ .

We say that  $C \subseteq \mathbb{P}$  is a *chain* iff C is linearly ordered that is for all  $a, b \in C$  we have  $a \leq b$  or  $b \leq a$ . b is an *upper bound* for C iff  $\forall a \in C \ a \leq b$ .

Zorn's Lemma: If  $\mathbb{P}$  is a poset such that every chain has an upper bound then for every p there is  $q \ge p$  such that q is maximal.

It is important to note that not all chains are necessarily of the form

 $a_0 \leq a_1 \leq a_2 \leq \dots$ 

For example let X be uncountable and let  $\mathbb{P}$  be the poset of countable subsets of X ordered by inclusion. Then  $\mathbb{P}$  has no maximal element even though every *countable* chain has an upper bound.

The following is easy:

**Lemma 1.** Let R be a ring,  $\mathcal{F}$  a nonempty family of ideals linearly ordered by inclusion. Then the union  $\bigcup \mathcal{F}$  is an ideal.

**Theorem 1.** Let R be a ring,  $I \neq R$ . Then there is  $J \supseteq I$  with J maximal.

*Proof.* Let  $\mathbb{P}$  be the set of all ideals J with  $J \neq R$ , ordered by inclusion. Clearly the maximal elements of  $\mathbb{P}$  are the maximal ideals, so it suffices to show that chains in  $\mathbb{P}$  have bounds. Let C be a chain. Then  $1 \notin \bigcup C$  so  $\bigcup C \neq R$ , and as we just saw  $\bigcup C$  is an ideal.  $\Box$ 

In particular if  $R \neq 0$  then (0) is an ideal not equal to R so that R has at least one maximal ideal.

We now discuss ring elements which are in some sense "pathological".

We say that  $a \in R$  is *nilpotent* iff there is n > 0 with  $a^n = 0$ . a is a zero-divisor iff ab = 0 for some nonzero b.

Let  $a \neq 0$  be nilpotent and let n > 1 be least such that  $a^n = 0$ . Then  $a^{n-1} \neq 0$  and  $aa^{n-1} = 0$  so a is a zero-divisor. In particular in an ID 0 is the only nilpotent element.

If a is nilpotent then 1 + a is a unit with inverse  $1 - a + \ldots a^{n-1}$ . In fact u + a is a unit for all units u.

If J is an ideal then  $\sqrt{J}$  is the set of a so that  $a^n \in J$  for some n > 0, equivalently a + J is nilpotent in R/J. We claim that  $\sqrt{J}$  is an ideal. If  $b^n \in J$  then  $(ab)^n =$ 

 $a^n b^n \in J$ . Also if  $a^m, b^n \in J$  then  $(a+b)^{m+n-1} \in J$  (expand it out as a sum of terms  $a^i b^j$  with i+j=m+n-1, so that  $i \ge m$  or  $j \ge n$ ).

J is radical iff  $J = \sqrt{J}$  or equivalently iff 0 is the only nilpotent in R/J. If P is prime then R/P is an ID, so by the remarks above P must be radical. Note that  $(0) \subseteq P$  and so  $\sqrt{(0)} \subseteq \sqrt{P} = P$ , that is any nilpotent element is in every prime ideal.

Remark: the intersection of any nonempty family of ideals is an ideal. So the ideal  $\sqrt{(0)}$  of nilpotent elements is contained in the ideal formed by intersecting all prime ideals of R. We claim that these ideals are equal, which amounts to showing that for every non-nilpotent a there is a prime P with  $a \notin P$ .

**Definition 2.** Let R be a ring.  $S \subseteq R$  is multiplicatively closed iff  $1 \in S$  and  $\forall a, b \in S \ ab \in S$ .

Remark: it is allowed that  $0 \in S$  but this will be a pathological case.

We can use the next result to manufacture prime ideals. If  $S = \{1\}$  it is just the trick for producing maximal ideals we already discussed.

**Theorem 2.** Let S be multiplicatively closed with  $0 \notin S$  and let  $\mathbb{P}$  be the set of ideals disjoint from S. Then  $\mathbb{P}$  is nonempty,  $\mathbb{P}$  is closed under unions of chains, and any maximal element of  $\mathbb{P}$  is prime.

*Proof.* Since  $0 \notin S$ ,  $(0) \in \mathbb{P}$ . Clearly it is closed under union of chains. Let I be maximal. Then  $1 \notin I$  as  $1 \in S$ , so  $I \neq R$ . Also if  $a \notin I$  then  $I \subsetneq I + (a)$ , so by maximality I + (a) meets S.

So if  $a, b \notin I$  we may choose  $s, t \in S$  with  $s \in I + (a)$  and  $t \in I + (b)$ . Then  $st \in I + (ab)$ , and since  $st \in S$  while I is disjoint from S we see that  $ab \notin I$ .  $\Box$