


Graph Theory


Simple Graph $G = (V, E)$.
 $V = \{\text{vertices}\}$, $E = \{\text{edges}\}$.



$V = \{a, b, c, d, e, f, g, h, k\}$
 $E = \{(a, b), (a, g), (a, h), (a, k), (b, c), (b, k), \dots, (h, k)\}$ $|E| = 16$.

Eulerian Graphs


Can you draw the diagram below without taking your pen off the paper or going over the same line twice?




Bipartite Graphs

G is bipartite if $V = X \cup Y$ where X and Y are disjoint and every edge is of the form (x, y) where $x \in X$ and $y \in Y$.

In the diagram below, A, B, C, D are women and a, b, c, d are men. There is an edge joining x and y iff x and y like each other. The thick edges form a "perfect matching" enabling everybody to be paired with someone they like. Not all graphs will have perfect matching!



Vertex Colouring



Let $C = \{\text{colours}\}$. A vertex colouring of G is a map $f : V \rightarrow C$. We say that $v \in V$ gets coloured with $f(v)$.

The colouring is *proper* iff $(a, b) \in E \Rightarrow f(a) \neq f(b)$.



The *Chromatic Number* $\chi(G)$ is the minimum number of colours in a proper colouring.

Application: $V = \{\text{exams}\}$. (a, b) is an edge iff there is some student who needs to take both exams. $\chi(G)$ is the minimum number of periods required in order that no student is scheduled to take two exams at once.


Subgraphs

$G' = (V', E')$ is a *subgraph* of $G = (V, E)$ if $V' \subseteq V$ and $E' \subseteq E$.

G' is a *spanning* subgraph if $V' = V$.



NOT SPANNING




SPANNING

If $V' \subseteq V$ then

$$G[V'] = (V', \{(u, v) \in E : u, v \in V'\})$$

is the subgraph of G *induced* by V' .



$G[\{a, b, c, d, e\}]$


Isomorphism

Similarly, if $E_1 \subseteq E$ then $G[E_1] = (V_1, E_1)$ where

$$V_1 = \{v \in V : \exists e \in E_1 \text{ such that } v \in e\}$$

is also *induced* (by E_1).



$$E_1 = \{(a, b), (a, d)\}$$



$G[E_1]$

$G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are *isomorphic* if there exists a bijection $f : V_1 \rightarrow V_2$ such that

$$(v, w) \in E_1 \leftrightarrow (f(v), f(w)) \in E_2.$$



$f(a)=A$ etc.

Complete Graphs


$$K_n = ([n], \{(i, j) : 1 \leq i < j \leq n\})$$

is the complete graph on n vertices.

$$K_{m,n} = ([m] \cup [n], \{(i, j) : i \in [m], j \in [n]\})$$

is the complete bipartite graph on $m + n$ vertices.

(The notation is a little imprecise but hopefully clear.)



K_5



$K_{2,3}$

Vertex Degrees


$d_G(v)$ = degree of vertex v in G

= number of edges incident with v

$$\delta(G) = \min_v d_G(v)$$

$$\Delta(G) = \max_v d_G(v)$$

G



$d_G(a)=2, d_G(g)=4$ etc.


$\delta(G)=2, \Delta(G)=4$.

Matrices and Graphs

Incidence matrix M : $V \times E$ matrix.

$$M(v, e) = \begin{cases} 1 & v \in e \\ 0 & v \notin e \end{cases}$$


	e_1	e_2	e_3	e_4	e_5	e_6	e_7	e_8
a		1	1				1	
b	1	1			1			
c	1			1		1		
d			1	1				1
e					1	1	1	1



Adjacency matrix A : $V \times V$ matrix.

$$A(v, w) = \begin{cases} 1 & v, w \text{ adjacent} \\ 0 & \text{otherwise} \end{cases}$$

	a	b	c	d	e
a		1		1	1
b	1		1		1
c		1		1	1
d	1		1		1
e	1	1	1	1	



+

+

+

+

Theorem 1

$$\sum_{v \in V} d_G(v) = 2|E|$$

Proof Consider the incidence matrix M .
Row v has $d_G(v)$ 1's. So

$$\# \text{ 1's in matrix } M \text{ is } \sum_{v \in V} d_G(v).$$

Column e has 2 1's. So

$$\# \text{ 1's in matrix } M \text{ is } 2|E|.$$

□

Corollary 1 *In any graph, the number of vertices of odd degree, is even.*

Proof Let $ODD = \{\text{odd degree vertices}\}$
and $EVEN = V \setminus ODD$.

$$\sum_{v \in ODD} d(v) = 2|E| - \sum_{v \in EVEN} d(v)$$

is even.

So $|ODD|$ is even.

□

+

13

+

14