

## Trees



A *tree* is a graph which is

- (a) Connected and
- (b) has no cycles (*acyclic*).


1

**Lemma 1** Let the components of  $G$  be


$C_1, C_2, \dots, C_r$ , Suppose  $e = (u, v) \notin E, u \in C_i, v \in C_j$ .

- (a)  $i = j \Rightarrow \omega(G + e) = \omega(G)$ .
- (b)  $i \neq j \Rightarrow \omega(G + e) = \omega(G) - 1$ .

(a)



(b)



**Proof** Every path  $P$  in  $G + e$  which is not in  $G$  must contain  $e$ . Also,

$$\omega(G + e) \leq \omega(G).$$

Suppose

$$(x = u_0, u_1, \dots, u_k = u, u_{k+1} = v, \dots, u_\ell = y)$$

is a path in  $G + e$  that uses  $e$ . Then clearly  $x \in C_i$  and  $y \in C_j$ .

(a) follows as now no new relations  $x \sim y$  are added.


(b) Only possible new relations  $x \sim y$  are for  $x \in C_i$  and  $y \in C_j$ . But  $u \sim v$  in  $G + e$  and so  $C_i \cup C_j$  becomes (only) new component.  $\square$

**Lemma 2**  $G = (V, E)$  is *acyclic* (forest) with (tree) components  $C_1, C_2, \dots, C_k$ .  $|V| = n$ .  $e = (u, v) \notin E, u \in C_i, v \in C_j$ .

- (a)  $i = j \Rightarrow G + e$  contains a cycle.
- (b)  $i \neq j \Rightarrow G + e$  is *acyclic* and has one less component.
- (c)  $G$  has  $n - k$  edges.

3


4



Suppose  $G + e$  contains the cycle  $C$ .  $e \in C$  else  $C$  is a cycle of  $G$ .

$$C = (u = u_0, u_1, \dots, u_\ell = v, u_0).$$

But then  $G$  contains the path  $(u_0, u_1, \dots, u_\ell)$  from  $u$  to  $v$  – contradiction.



The drop in the number of components follows from Lemma 1.

The rest follows from

**Corollary 1** *If a tree  $T$  has  $n$  vertices then*

**(a)** *It has  $n - 1$  edges.*

**(b)** *It has at least 2 vertices of degree 1, ( $n \geq 2$ ).*

**Proof** (a) is part (c) of previous lemma.  $k = 1$  since  $T$  is connected.


(b) Let  $s$  be the number of vertices of degree 1 in  $T$ . There are no vertices of degree 0 – these would form separate components. Thus

$$2n - 2 = \sum_{v \in V} d_T(v) \geq 2(n - s) + s.$$

So  $s \geq 2$ . □

(a)  $u, v \in C_i$  implies there exists a path  $(u = u_0, u_1, \dots, u_\ell = v)$  in  $G$ .

So  $G + e$  contains the cycle  $u_0, u_1, \dots, u_\ell, u_0$ .



(c) Suppose  $E = \{e_1, e_2, \dots, e_r\}$  and  $G_i = (V, \{e_1, e_2, \dots, e_i\})$  for  $0 \leq i \leq r$ .

**Claim:**  $G_i$  has  $n - i$  components.

Induction on  $i$ .

$i = 0$ :  $G_0$  has no edges.

$i > 0$ :  $G_{i-1}$  is acyclic and so is  $G_i$ . It follows from part (a) that  $e_i$  joins vertices in distinct components of  $G_{i-1}$ . It follows from (b) that  $G_i$  has one less component than  $G_{i-1}$ .

**End of proof of claim**

Thus  $r = n - k$  (we assumed  $G$  had  $k$  components). □

**Theorem 1** Suppose  $|V| = n$  and  $|E| = n - 1$ . The following three statements become equivalent.

(a)  $G$  is connected.

(b)  $G$  is acyclic.

(c)  $G$  is a tree.

Let  $E = \{e_1, e_2, \dots, e_{n-1}\}$  and  $G_i = (V, \{e_1, e_2, \dots, e_i\})$  for  $0 \leq i \leq n - 1$ .

(a)  $\Rightarrow$  (b):  $G_0$  has  $n$  components and  $G_{n-1}$  has 1 component. Addition of each edge  $e_i$  must reduce the number of components by 1 – Lemma 2(b). Thus  $G_{i-1}$  acyclic implies  $G_i$  is acyclic. (b) follows as  $G_0$  is acyclic.


(b)  $\Rightarrow$  (c): We need to show that  $G$  is connected. Since  $G_{n-1}$  is acyclic,  $\omega(G_i) = \omega(G_{i-1}) - 1$  for each  $i$  – Lemma 2(b). Thus  $\omega(G_{n-1}) = 1$ .

(c)  $\Rightarrow$  (a): trivial.

9

10


**Corollary 2** If  $v$  is a vertex of degree 1 in a tree  $T$  then  $T - v$  is also a tree.



**Proof** Suppose  $T$  has  $n$  vertices and  $n$  edges. Then  $T - v$  has  $n - 1$  vertices and  $n - 2$  edges. It is acyclic and so must be a tree.  $\square$

11

### Cut edges



$e$  is a *cut edge* of  $G$  if  $\omega(G - e) > \omega(G)$ .


**Theorem 2**  $e = (u, v)$  is a cut edge iff  $e$  is not on any cycle of  $G$ .

**Proof**  $\omega$  increases iff there exist  $x \sim y \in V$  such that all walks from  $x$  to  $y$  use  $e$ .

Suppose there is a cycle  $(u, P, v, u)$  containing  $e$ . Then if  $W = x, W_1, u, v, W_2, y$  is a walk from  $x$  to  $y$  using  $e$ ,  $x, W_1, P, W_2, y$  is a walk from  $x$  to  $y$  that doesn't use  $e$ . Thus  $e$  is not a cut edge.

12

x



If  $e$  is not a cut edge then  $G - e$  contains a path  $P$  from  $u$  to  $v$  ( $u \sim v$  in  $G$  and relations are maintained after deletion of  $e$ ). So  $(v, u, P, v)$  is a cycle containing  $e$ .  $\square$

**Corollary 3** A connected graph is a tree iff every edge is a cut edge.

13

**Corollary 4** Every finite connected graph  $G$  contains a spanning tree.

**Proof** Consider the following process: starting with  $G$ ,

1. If there are no cycles – **stop**.
2. If there is a cycle, delete an edge of a cycle.


Observe that (i) the graph remains connected – we delete edges of cycles. (ii) the process must terminate as the number of edges is assumed finite.

On termination there are no cycles and so we have a connected acyclic spanning subgraph i.e. we have a spanning tree.  $\square$

14

**Theorem 3** Let  $T$  be a spanning tree of  $G = (V, E)$ ,  $|V| = n$ . Suppose  $e = (u, v) \in E \setminus T$ .

- (a)  $T + e$  contains a unique cycle  $C(e)$ .
- (b)  $f \in C(e)$  implies that  $T + e - f$  is a spanning tree of  $G$ .



15

**Proof** (a) Lemma 2(a) implies that  $T + e$  has a cycle  $C$ . Suppose that  $T + e$  contains another cycle  $C' \neq C$ . Let edge  $g \in C' \setminus C$ .  $T' = T + e - g$  is connected, has  $n - 1$  edges. But  $T'$  contains a cycle  $C$ , contradicting Lemma 2(b).

(b)  $T + e - f$  is connected and has  $n - 1$  edges. Therefore it is a tree.  $\square$

16


## Maximum weight trees

$G = (V, E)$  is a connected graph.

$w : E \rightarrow \mathbf{R}$ .  $w(e)$  is the *weight* of edge  $e$ .

For spanning tree  $T$ ,  $w(T) = \sum_{e \in T} w(e)$ .

**Problem:** find a spanning tree of maximum weight.



17

## Greedy Algorithm

Sort edges so that  $E = \{e_1, e_2, \dots, e_m\}$  where

$$w(e_1) \geq w(e_2) \geq \dots \geq w(e_m).$$

**begin**

$T := \emptyset$

**for**  $i = 1, 2, \dots, m$  **do**

**begin**

**if**  $T + e_i$  does not contain a cycle

**then**  $T \leftarrow T + e_i$

**end**

Output  $T$

**end**

Greedy always adds the maximum weight edge which does not make a cycle with previously chosen edges.

18

**Theorem 4** *The tree constructed by GREEDY has maximum weight.*

**Proof** Let the edges of the *greedy tree* be  $e_1^*, e_2^*, \dots, e_{n-1}^*$  in order of choice. Note that  $w(e_i^*) \geq w(e_{i+1}^*)$  since neither makes a cycle with  $e_1^*, e_2^*, \dots, e_{i-1}^*$ .

Let  $f_1, f_2, \dots, f_{n-1}$  be the edges of any other tree where  $w(f_1) \geq w(f_2) \geq \dots \geq w(f_{n-1})$ .

We show that


$$w(e_i^*) \geq w(f_i) \quad 1 \leq i \leq n-1. \quad (1)$$

Suppose (1) is false. There exists  $k > 0$  such that

$$w(e_i^*) \geq w(f_i), \quad 1 \leq i < k \text{ and } w(e_k^*) < w(f_k).$$

Each of  $f_i$ ,  $1 \leq i \leq k$  makes a cycle with  $e_1^*, e_2^*, \dots, e_{k-1}^*$ . Otherwise one of them would have been chosen in preference to  $e_k^*$ .

Let the components of the forest  $(V, \{e_1^*, e_2^*, \dots, e_{k-1}^*\})$  be  $C_1, C_2, \dots, C_{n-k+1}$ . Each  $f_i$ ,  $1 \leq i \leq k$  has both of its endpoints in the same component.



19

20

Let  $\mu_i$  be the number of  $f_j$  which have both endpoints in  $C_i$  and let  $\nu_i$  be the number of vertices of  $C_i$ . Then

$$\mu_1 + \mu_2 + \cdots + \mu_{n-k+1} = k \quad (2)$$

$$\nu_1 + \nu_2 + \cdots + \nu_{n-k+1} = n \quad (3)$$

It follows from (2) and (3) that there exists  $t$  such that

$$\mu_t \geq \nu_t. \quad (4)$$

[Otherwise


$$\begin{aligned} \sum_{i=1}^{n-k+1} \mu_i &\leq \sum_{i=1}^{n-k+1} (\nu_i - 1) \\ &= \sum_{i=1}^{n-k+1} \nu_i - (n - k + 1) \\ &= k - 1. \end{aligned} \quad ]$$

But (4) implies that the edges  $f_j$  such that  $f_j \subseteq C_t$  contain a cycle.  $\square$



21

## How many trees? – Cayley's Formula




n=4



n=5



n=6



22

## Prüfer's Correspondence

There is a 1-1 correspondence  $\phi_V$  between spanning trees of  $K_V$  (the complete graph with vertex set  $V$ ) and sequences  $V^{n-2}$ . Thus for  $n \geq 2$

$$\tau(K_n) = n^{n-2} \quad \text{Cayley's Formula.}$$

Assume some arbitrary ordering  $V = \{v_1 < v_2 < \cdots < v_n\}$ .

$\phi_V(T)$ :

**begin**

$T_1 := T$ ;

**for**  $i = 1$  **to**  $n - 2$  **do**


**begin**

$s_i :=$  neighbour of least leaf  $\ell_i$  of  $T_i$ .

$T_{i+1} = T_i - \ell_i$ .

**end**  $\phi_V(T) = s_1 s_2 \dots s_{n-2}$

**end**



6,4,5,14,2,6,11,14,8,5,11,4,2

23


24

**Lemma 3**  $v \in V(T)$  appears exactly  $d_T(v) - 1$  times in  $\phi_V(T)$ .

**Proof** Assume  $n = |V(T)| \geq 2$ . By induction on  $n$ .

$n = 2$ :  $\phi_V(T) = \Lambda$  = empty string.

Assume  $n \geq 3$ :



$\phi_V(T) = s_1 \phi_{V_1}(T_1)$  where  $V_1 = V - \{s_1\}$ .

$s_1$  appears  $d_{T_1}(s_1) - 1 + 1 = d_T(s_1) - 1$  times – induction.

$v \neq s_1$  appears  $d_{T_1}(v) - 1 = d_T(v) - 1$  times – induction.  $\square$

25

### Construction of $\phi_V^{-1}$

Inductively assume that for all  $|X| < n$  there is an inverse function  $\phi_X^{-1}$ . (True for  $n = 2$ ).

Now define  $\phi_V^{-1}$  by

$\phi_V^{-1}(s_1 s_2 \dots s_{n-2}) = \phi_{V_1}^{-1}(s_2 \dots s_{n-2})$  plus edge  $s_1 \ell_1$ ,

where  $\ell_1 = \min\{s : s \notin s_1, s_2, \dots, s_{n-2}\}$  and  $V_1 = V - \{\ell_1\}$ . Then

$$\begin{aligned} \phi_V(\phi_V^{-1}(s_1 s_2 \dots s_{n-2})) &= s_1 \phi_{V_1}(\phi_{V_1}^{-1}(s_2 \dots s_{n-2})) \\ &= s_1 s_2 \dots s_{n-2}. \end{aligned}$$

Thus  $\phi_V$  has an inverse and the correspondence is established.

26

### Number of trees with a given degree sequence

**Corollary 5** If  $d_1 + d_2 + \dots + d_n = 2n - 2$  then the number of spanning trees of  $K_n$  with degree sequence  $d_1, d_2, \dots, d_n$  is

$$\binom{n-2}{d_1-1, d_2-1, \dots, d_n-1} = \frac{(n-2)!}{(d_1-1)!(d_2-1)! \dots (d_n-1)!}.$$

**Proof** From Prüfer's correspondence and Lemma 3 this is the number of sequences of length  $n - 2$  in which 1 appears  $d_1 - 1$  times, 2 appears  $d_2 - 1$  times and so on.  $\square$