

Day 21

June 21, 2012

1 Counting Proofs

We have already seen some interesting properties of numbers witnessed by counting. For example:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

The left is, by definition, the number of ways that we can choose k objects from a set of size n . The right, however, is just some numbers, which incidentally count the same thing. The proof of this equality was showing that both expressions counted the same set. That is, they both count the number of ways we can choose k objects from n . An interesting corollary of this fact is that $\frac{n!}{k!(n-k)!}$ is an integer. It's not immediately clear that it is, but because it counts a set of objects, it must be an integer.

As you would probably guess, this is not the only such proof which is interesting. There are many combinatorial identities. To prove them, we can usually use several methods. Sometimes we can use arithmetic and identities already proved; the above is a particularly important one because it allows us to actually do arithmetic with binomial coefficients. Another way we can prove combinatorial identities, often enough is induction. We have already seen, for instance, a proof that 2^n counts the subsets of a set of size n by induction.

What we are interested in, however, is combinatorial proofs, as in the one above.

1.1 Some Examples of Combinatorial Proofs

Example 1. We have already seen many examples of combinatorial proofs, in a way. Recall we counted the number of ways to permute the English alphabet such that between x and y there are exactly 5 letters. We did this several different ways

- Step 1** Decide if x or y comes first 2 ways
- Step 2** Arrange the remaining letters; the first five go between x and y 24! ways
- Step 3** Pick a spot in the alphabet for the beginning of the x, y block. The other letters go around this position in the order chosen 20 ways

This makes the total $2 \cdot 20 \cdot 24!$.

But, we could have equally well counted this a different way:

- Step 1** Decide if x or y comes first 2 ways
- Step 2** Arrange the remaining letters; the first 5 go between x and y 24! ways
- Step 3** Decide how many of the remaining 19 letters go before and after this x, y block by breaking them into two groups $\binom{19+2-1}{19}$

Note, it is fairly easy to see these are equal from looking at it, but they do represent different processes of counting. This proves therefore that

$$2 \cdot 20 \cdot 24! = 2 \cdot 24! \cdot \binom{20}{19}$$

There is no arithmetic or algebraic argument needed. They are equal because they both count the number of ways to permute the alphabet so there is exactly five letters between x and y .

Theorem 3.

$$\sum_{i=1}^n i \cdot \binom{n}{i} = n \cdot 2^{n-1}$$

Proof. The right hand side is overwhelmingly simpler, so we first try to count using that. Note, that n chooses an object from a set of size n , and 2^{n-1} seems to make a subset. Therefore, the set we choose to count is the number of committees that we can form from n people with a chairperson.

- Step 1** Choose chairperson n ways
- Step 2** Pick the remaining committee 2^{n-1} ways

So, we wish to count this in a different ways. Note, the sum on the left hand side suggests that we should so cases. From the form, it looks as though the cases should be done on the size of the committee being formed.

Case 1	The committee has one person	
Step 1	Choose the committee	$\binom{n}{1}$ ways
Step 2	Choose the chairperson among those people	1 way
Case 2	The committee has two people	
Step 1	Choose the committee	$\binom{n}{2}$ ways
Step 2	Choose the chairperson among those people	2 ways
\vdots		
Case n	The committee has n people	
Step 1	Choose the committee	$\binom{n}{n}$ ways
Step 2	Choose the chairperson among those people	n ways

We sum across the cases, giving us the right hand side.

Since we counted the same set using both the left hand side of the above and the right hand side they are equal. □

Theorem 4.

$$\sum_{i=0}^n \binom{n-i}{k} = \binom{n+1}{k+1}$$

Proof. It is plain one thing that the right hand side counts: the number of ways to choose subset of of size $k + 1$ from a universe of size $n + 1$. But instead, here we will look at the left hand side as binary sequences (sequences of 0's and 1's) and the right hand side is the the number of binary sequences of length n with $k + 1$ many 1's.

For the left hand side, we do cases of where the first 1 appears (I am thinking of i as the number of 0's that appear before the first 1).

In the case when there are no 0's before the first 1, we know our sequence begins with a 1, and then there are k other positions we must put 1 in the remaining n spots. So the total is $\binom{n}{k}$. If there is one 0 before 1, we know the sequence begins 01, and then there are k other positions where we must put 1 in the remaining $n - 1$ spots. Continuing in this fashion, we have that the total number of ways for each i is $\binom{n-i}{k}$. Summing across all the possible number of 0's before the first 1, we get the left hand side. □