

OPERATIONS RESEARCH II 21-393

Homework 3: Due Monday October 1.

Q1 Solve the following problem by a cutting plane algorithm:

$$\begin{aligned} &\text{minimise} && 4x_1 + 5x_2 + 3x_3 \\ &\text{subject to} && \\ &&& 2x_1 + x_2 - x_3 \geq 2 \\ &&& x_1 + 4x_2 + x_3 \geq 13 \end{aligned}$$

$$x_1, x_2, x_3 \geq 0 \text{ and integer.}$$

Solution

Initial tableau

x_1	x_2	x_3	x_4	x_5	R.H.S	
-4	-5	-3	0	0	0	z
-2	-1	1	1	0	-2	x_4
-1	-4	-1	0	1	-13	$x_5 \leftarrow$
	\uparrow					

x_1	x_2	x_3	x_4	x_5	R.H.S	
$\frac{-11}{4}$	0	$\frac{-7}{4}$	0	$\frac{-5}{4}$	$\frac{65}{4}$	z
$\frac{-7}{4}$	0	$\frac{5}{4}$	1	$\frac{-1}{4}$	$\frac{5}{4}$	x_4
$\frac{1}{4}$	1	$\frac{1}{4}$	0	$\frac{-1}{4}$	$\frac{13}{4}$	x_2

Primal feasible, but the solution is not integral.

We add a cut which eliminates the current optimal solution.

$$\frac{1}{4}x_1 + \frac{1}{4}x_3 + \frac{3}{4}x_5 - y_1 = \frac{1}{4}$$

x_1	x_2	x_3	x_4	x_5	y_1	R.H.S	
$\frac{-11}{4}$	0	$\frac{-7}{4}$	0	$\frac{-5}{4}$	0	$\frac{65}{4}$	Z
$\frac{-7}{4}$	0	$\frac{5}{4}$	1	$\frac{-1}{4}$	0	$\frac{5}{4}$	x_4
$\frac{4}{4}$	1	$\frac{4}{4}$	0	$\frac{-1}{4}$	0	$\frac{13}{4}$	x_2
$\frac{-11}{4}$	0	$\frac{-1}{4}$	0	$\frac{-3}{4}$	+1	$\frac{-1}{4}$	$y_1 \leftarrow$
				\uparrow			

We do a dual simplex pivot to obtain

x_1	x_2	x_3	x_4	x_5	y_1	R.H.S	
$\frac{-7}{3}$	0	$\frac{-4}{3}$	0	$\frac{-5}{3}$	0	$\frac{50}{3}$	Z
$\frac{-5}{3}$	0	$\frac{4}{3}$	1	0	$\frac{-1}{3}$	$\frac{4}{3}$	x_4
$\frac{1}{3}$	1	$\frac{1}{3}$	0	0	$\frac{-1}{3}$	$\frac{10}{3}$	x_2
$\frac{1}{3}$	0	$\frac{1}{3}$	0	1	$\frac{-4}{3}$	$\frac{1}{3}$	x_5

The solution is primal feasible and so optimal but still not integer.

We add a cut which eliminates the current optimal solution.

$$-\frac{1}{3}x_1 - \frac{1}{3}x_3 + y_2 = \frac{1}{3}$$

We obtain tableau

x_1	x_2	x_3	x_4	x_5	y_2	R.H.S	
$\frac{-7}{3}$	0	$\frac{-4}{3}$	0	0	0	$\frac{50}{3}$	Z
$\frac{-5}{3}$	0	$\frac{4}{3}$	1	0	0	$\frac{4}{3}$	x_4
$\frac{1}{3}$	1	$\frac{1}{3}$	0	0	0	$\frac{10}{3}$	x_2
$\frac{1}{3}$	0	$\frac{1}{3}$	0	1	0	$\frac{1}{3}$	x_5
$\frac{-1}{3}$	0	$\frac{-1}{3}$	0	0	1	$\frac{1}{3}$	$y_2 \leftarrow$
		\uparrow					

We do a dual simplex pivot to obtain

x_1	x_2	x_3	x_4	x_5	y_2	R.H.S	
-1	0	0	0	0	-4	18	Z
-3	0	0	1	0	4	0	x_4
0	1	0	0	0	1	3	x_2
0	0	0	0	1	1	0	x_5
1	0	1	0	0	-3	1	x_3

Which is optimal integral.

- Q2** Formulate the following as an integer program: There are n students and exams $E_1, E_2, \dots, E_m \subseteq [n]$ need to be scheduled. There are s rooms available and each room can hold r students. The rules are
- (i) A student must not be asked to take more than one exam per day;
 - (ii) Several different exams can be held in the same room provided there is capacity in the room to hold the students.
 - (iii) No student has to take 3 exams in 3 consecutive days.

The problem is to minimise the number of days needed to carry out all of the exams.

Solution Define

$$x_{i,j,k} = \begin{cases} 1 & \text{Exam } i \text{ takes place in room } j \text{ on day } k \\ 0 & \text{Otherwise} \end{cases}$$

and

$$y_i = \begin{cases} 1 & \text{At least one exam takes place on day } j \\ 0 & \text{Otherwise} \end{cases}$$

Then we have to solve the problem

$$\text{Minimise } \sum_{i=1}^m y_i$$

Subject to

$$\sum_{j=1}^s \sum_{k=1}^n x_{i,j,k} = 1 \quad \text{for all } i.$$

$$x_{i,j,k} \leq y_k \quad \text{for all } i, j, k.$$

$$\sum_{i=1}^m |E_i| x_{i,j,k} \leq r \quad \text{for all } j, k.$$

$$x_{i,j,k} + x_{i',j,k} \leq 1, \quad \text{for all } E_i \cap E_{i'} \neq \emptyset, j, k.$$

$$x_{i,j,k} + x_{i',j',k+1} + x_{i'',j'',k+2} \leq 2, \quad \text{for all } E_i \cap E_{i'} \cap E_{i''} \neq \emptyset.$$

- Q3** An assembly line consists of a sequence of locations called work stations. The manufacture of a certain object requires m separate jobs to be undertaken with job i requiring t_i minutes. The jobs are to be allocated to work stations so that each station completes a set of jobs and then passes

the object onto the next station on the line and waits to receive the next object from the previous station on the line. The combined time of all jobs assigned to any station must not exceed T the cycle time. Also there are a number of precedence relations between jobs indicated by the digraph $D = (V, A)$ where $(i, j) \in A$ if job i must precede job j . The problem is to open as few work stations as possible consistent with the cycle time. Formulate this as an integer programming problem.

Solution

$$\begin{aligned}
 & \text{Minimize } \sum_{i=1}^m y_i \\
 & \text{s.t.} \\
 & \quad x_{ij} \leq y_i \quad \forall i, j \in \{1, \dots, m\} \\
 & \quad \sum_{j=1}^m x_{ij}t_j \leq T \quad \forall i \in \{1, 2, \dots, m\} \\
 & \quad \sum_{i=1}^m x_{ij} = 1 \quad \forall j \in \{1, 2, \dots, m\} \\
 & \quad \sum_{i=1}^m ix_{ij_1} \leq \sum_{i=1}^m ix_{ij_2} \quad \forall (j_1, j_2) \in A \\
 & \quad x_{ij} \in \{0, 1\} \quad \forall i, j \in \{1, \dots, m\} \\
 & \quad y_i \in \{0, 1\} \quad \forall i \in \{1, \dots, m\}
 \end{aligned}$$

x_{ij} is 1 when job j is done at station i and 0 otherwise.

y_i is 1 if at least one job is done at station i .

The first constraint ensures that if any job is done at station i , the variable y_i is 1.

The second constraint ensures that each station satisfies the cycle time T .

The third constraint ensures that each job is scheduled on some machine.

The last constraint ensures the j_1 is done before job j_2 if there is a precedence constraint between them.