Math 268: Homework.

The late homework policy on the class website will be strictly followed. I will often assign harder optional problems. I recommend doing (but not turning in) the optional problems. They often involve useful concepts that will come in handy as the semester progresses.

Assignment 1 (assigned 2020-01-15, due 2020-01-22).

- 1. Determine (without proving) whether the following sets are open. (No proof required; just guess.)
 - $\begin{array}{ll} \text{(a)} & \{x \in \mathbb{R}^3 \mid 1 < |x| < 10\}. \\ \text{(b)} & \{x \in \mathbb{R}^3 \mid 1 \leqslant |x| < 10\}. \\ \end{array} \\ \begin{array}{ll} \text{(c)} & \{x \in \mathbb{R}^3 \mid x_1^2 + x_2^2 < |x_3|\}. \\ \text{(d)} & \{x \in \mathbb{R}^{268} \mid \sum_{i < 268} x_i^2 < |x_{268}|\}. \end{array} \\ \end{array}$
- 2. Decide whether $\{x \in \mathbb{R}^2 \mid x_1 + x_2 < x_2^2\}$ is open. Prove your answer.
- 3. (a) If U_1, U_2, \ldots are infinitely many open sets, must the union $U = U_1 \cup U_2 \cdots$ also be open? Prove it.
 - (b) If U_1, U_2, \ldots, U_N are finitely many open sets, must the intersection U = $U_1 \cap U_2 \cdots \cap U_N$ also be open? Prove it.
 - (c) Find infinitely many open sets U_1, U_2, \ldots , such that the intersection U = $U_1 \cap U_2 \cdots$ is not open.

$$\begin{split} \bar{A} &\stackrel{\text{def}}{=} \left\{ x \in \mathbb{R}^d \mid \forall \varepsilon > 0, \ B(x,\varepsilon) \cap A \neq \emptyset \right\}, \\ & \mathring{A} \stackrel{\text{def}}{=} \left\{ x \in \mathbb{R}^d \mid \exists \varepsilon > 0 \text{ such that } B(x,\varepsilon) \subseteq A \right\}, \\ & \partial A \stackrel{\text{def}}{=} \left\{ x \in \mathbb{R}^d \mid \forall \varepsilon > 0, \ B(x,\varepsilon) \cap A \neq \emptyset, \text{ and } B(x,\varepsilon) \cap A^c \neq \emptyset \right\}. \end{split}$$

Show that $\partial A = \overline{A} - A$.

- 5. For each set in the first question, find the closure, interior and boundary. (No proof required: just make an educated guess.)
- 6. Let $U \subset \mathbb{R}^d$ be a domain, $a \in U$, and $f: U \to \mathbb{R}^n$ be a function.
 - (a) If $\lim_{x\to a} f(x) = \ell$, then for every $i \in \{1, \ldots, n\}$ must $\lim_{x\to a} f_i(x) = \ell_i$? Prove it, or find a counter example. [Here f is the vector function (f_1, f_2, \ldots, f_n) .]
 - (b) Conversely, if for every $i \in \{1, \ldots, n\}$ we have $\lim_{x \to a} f_i(x) = \ell_i$, then must $\lim_{x\to a} f(x) = \ell$? Prove it, or find a counter example.

Optional fun questions to think about

* (a) Does
$$\lim_{|x|\to 0} \frac{x_1 x_2}{|x|^2}$$
 exist? Justify. [Here $x \in \mathbb{R}^2$.]
(b) Does $\lim_{|x|\to 0} \frac{x_1 x_2}{|x|}$ exist? Justify. [Here $x \in \mathbb{R}^2$.]

* Prove that any open set in \mathbb{R} can be expressed as a countable union of open sets.

Assignment 2 (assigned 2020-01-22, due 2020-01-29).

1. For each of the following functions defined on $\mathbb{R}^2 - \{(0,0)\}$ determine if the function has a limit as $(x, y) \rightarrow (0, 0)$. Prove your answer.

(a)
$$f(x,y) = \frac{3x^2y^2}{x^2 + y^2}$$

(b) $f(x,y) = \frac{xy(x^2 - y^2)}{x^4 + y^4}$
(c) $f(x,y) = \frac{x^3y^4}{|x|^5 + y^6}$
(d) $f(x,y) = \frac{x^2y^3}{(x^4 + y^6)^{1/3}}$

- 2. (a) Suppose $f:[a,b] \to \mathbb{R}$ is differentiable, and f' is increasing. Show that for all $x, y \in [a, b], \theta \in [0, 1]$ we have $f(\theta x + (1 - \theta)y) \leq \theta f(x) + (1 - \theta)f(y)$.
 - (b) Conversely, suppose for all $x, y \in [a, b], \theta \in [0, 1]$ we have $f(\theta x + (1 \theta)y) \leq \theta$ $\theta f(x) + (1 - \theta) f(y)$. If f is differentiable, show that f' is increasing.

A function that satisfies either of the above properties is called convex. (The second form is preferable since it doesn't assume differentiability.)

- (c) If p, q > 1 with 1/p + 1/q = 1 and $x, y \in \mathbb{R}$ show that $xy \leq |x|^p/p + |y|^q/q$. [Hint: This is part (c) of a question. Google "Young's inequality" if you need more help.]
- (d) Does $\lim_{x \to 0} \frac{x_1 x_2}{(|x_1|^{4/3} + x_2^4)^{.99}}$ exist? How about $\lim_{x \to 0} \frac{x_1 x_2}{(|x_1|^{4/3} + x_2^4)^{1.01}}$ exist? Prove it. [HINT: This is the last part of a question. (However, there is also an elegant short solution to this part without using any of the previous parts.)]
- 4. Given a set $A \subseteq \mathbb{R}^d$ define the closure \overline{A} , interior \mathring{A} and boundary ∂A as follows: 3. Let $U \subseteq \mathbb{R}^d$ be open and $f: \mathbb{R}^d \to \mathbb{R}^n$ be continuous. True or false: For every open set $V \subseteq \mathbb{R}^n$, the set $f^{-1}(V)$ is also open. Prove it, or find a counter example. [Once you do this problem, you might want to revisit some examples of open sets from the first homework.]
 - 4. Compute the following partial derivatives. (You should determine from context whether $x \in \mathbb{R}$, or $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$.)
 - (a) $\partial_x \tan^{-1}\left(\frac{y}{r}\right)$ (e) $\partial_t \int_0^t \exp(-|x|^2 - s^2) ds$ (b) $\partial_1 \ln |x|$ (f) $\partial_{x_1} \int_0^t \exp(-|x|^2 - s^2) ds$ (c) $\frac{\partial}{\partial x} x^y$

(d)
$$\frac{\partial}{\partial y} x^y$$

1

5. Let $f(x, y) = \frac{x^2y}{x^2 + y^2}$ for $(x, y) \neq 0$, and f(0, 0) = 0.

- (a) At what points in \mathbb{R}^2 do $\partial_x f$ and $\partial_y f$ exist? Justify.
- (b) At what points in \mathbb{R}^2 are $\partial_x f$ and $\partial_y f$ continuous? Justify.
- (c) Find all $v \in \mathbb{R}^2$ so that $D_v f(0,0)$ exists. Justify.
- (d) Is f differentiable at (0,0)? Justify.

Assignment 3 (assigned 2020-01-29, due 2020-02-05).

- 1. For each of the functions below, decide at what points the function is differentiable (no proof required; just guess). Further compute the derivative.
 - (d) $z = (x 2y)^5 e^{xy}$. (a) f(x, y) = x/y(b) $f(x) = \ln|x|, x \in \mathbb{R}^2$. (e) $f(x, y) = \tan^{-1}(y/x)$. (c) $f(x,y) = \frac{xy}{1-x-y}$ (f) $f(x) = 1/|x|, x \in \mathbb{R}^3$.
- 2. Let $f: \mathbb{R}^2 \to \mathbb{R}$ be defined by $f(x) = |x|^2 \sin(1/|x|)$ when $x \neq 0$, and f(0) = 0.
 - (a) Show that $\partial_1 f$ and $\partial_2 f$ are continuous in \mathbb{R}^2 except at 0.
 - (b) Show that f is differentiable at all points in \mathbb{R}^2 , including at a = 0.
- 3. (a) Let $U = \{(x,y) \in \mathbb{R}^2 \mid x > 0\}$. Define $r(x,y) = \sqrt{x^2 + y^2}, \ \theta(x,y) = \sqrt{x^2 + y^2}$ $\tan^{-1}(y/x)$, and $\varphi: U \to \mathbb{R}^2$ by $\varphi(x,y) = (r(x,y), \theta(x,y))$. Compute $D\varphi$ and $\det(D\varphi)$.
 - (b) Let $V = \{(r, \theta) \in \mathbb{R}^2 \mid r > 0, |\theta| < \pi/2\}$. Let $x(r, \theta) = r \cos \theta$, and $y(r, \theta) = r \cos \theta$. $r\sin\theta$ and $\psi(r,\theta) = (x(r,\theta), y(r,\theta))$. Compute $D\psi$ and $\det(D\psi)$.
- 4. Determine whether each of these statements are true or false. If true, prove it. If not, find a counter example. (Assume d > 1 below.)
 - necessarily continuous.
 - (b) If $f : \mathbb{R}^d \to \mathbb{R}$ is such that $\partial_i f$ exists for all $i \in \{1, \ldots, d\}$, then f is necessarily continuous.
 - (c) If $f : \mathbb{R}^d \to \mathbb{R}$ is such that $D_v f(a)$ exists for all $v \in \mathbb{R}^d \{0\}$, then f is necessarily continuous at a.
 - (d) Any differentiable function $f : \mathbb{R}^m \to \mathbb{R}^n$ is continuous.
 - (e) Any continuous function $f : \mathbb{R}^m \to \mathbb{R}^n$ is differentiable.
- 5. Suppose $f, q: \mathbb{R}^d \to \mathbb{R}$ are both differentiable at a. Show that fq is also differentiable at a, and find a formula for $D(fg)_a$ in terms of f(a), g(a), Df_a and Dg_a .
 - HINT: To a formula for $D(fg)_a$, compute $\partial_i(fg)$ using the standard one variable product rule. Now show that fq is differentiable directly using the definition of the derivative.

Assignment 4 (assigned 2020-02-05, due Never).

- In light of your Midterm on 2/12, this homework is not due. However, many of the problems are good practice, and I recommend doing (or at least thinking) about them before your exam. Some of these problems will find their way onto your next homework.
- 1. For each of the functions y below express the derivative (with respect to x) as a product of two matrices, and evaluate the matrices at the given values of x below.

(a)
$$y = \begin{pmatrix} u_1 u_2 - 3u_1 \\ u_2^2 + 2u_1 u_2 + 2u_1 - u_2 \end{pmatrix}$$
, $u = \begin{pmatrix} x_1 \cos(3x_2) \\ x_1 \sin(3x_2) \end{pmatrix}$ at $x = 0$.
(b) $y = \begin{pmatrix} u_1^2 + u_2^2 - 3u_1 + u_3 \\ u_1^2 - u_2^2 + 2u_1 - 3u_3 \end{pmatrix}$, $u = \begin{pmatrix} x_1 x_2 x_3^2 \\ x_1 x_2^2 x_3^2 \\ x_1^2 x_2 x_3 \end{pmatrix}$, at $x = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$.

- 2. (a) Let F(x,y) = xy. Given two differentiable functions $f, g: \mathbb{R}^d \to \mathbb{R}$, let $G: \mathbb{R}^d \to \mathbb{R}^2$ be defined by G(x) = (f(x), g(x)). Observe $D(fg) = D(F \circ G)$. Compute this using the chain rule, and derive the product rule.
 - (b) Derive the quotient rule from the chain rule using a method similar to the previous question.
- (a) If $f: \mathbb{R}^d \to \mathbb{R}$ is such that $\partial_i f$ exists for some $i \in \{1, \ldots, d\}$, then f is 3. Let $U = \mathbb{R}^2 \{(x, 0) \mid x \leq 0\}$, and $V = \{(r, \theta) \mid r > 0, \theta \in (-\pi, \pi)\}$. Given a differentiable function f defined on U, we treat it as a function of the coordinates x and y. Using the relation $x = r \cos \theta$ and $y = r \sin \theta$ for $(r, \theta) \in V$, we can now treat f as a function of r and θ .
 - (a) Express $\partial_r f$ and $\partial_\theta f$ in terms of $\partial_x f$, $\partial_u f$, r and θ .
 - (b) Let $u = x^2 + y^2$ and v = y/x. Explicitly express u, v in terms of r and θ and compute $\partial_r u$, $\partial_{\theta} u$, $\partial_r v$ and $\partial_{\theta} v$ directly. Verify that this agrees with the formulae in the previous part.
 - (c) If further r, θ are functions functions of variables s and t, compute $\partial_s f$ in terms of $\partial_x f$, $\partial_\mu f$, r, θ , $\partial_s r$ and $\partial_s \theta$.
 - (d) Express r, θ in terms of x and y.
 - (e) Suppose now q is a differentiable function defined on V, which we treat as a function as a function of r and θ . Using the previous part, we can treat q as a function of x and y. Compute $\partial_x g$ and $\partial_y g$ in terms of $\partial_r g$, $\partial_\theta g$, x and y. Verify your formula is correct for the function $q(r, \theta) = r\theta$.
 - 4. Let $f: \mathbb{R}^d \to \mathbb{R}^n$ be differentiable, and Γ be the graph of f, and $(a, f(a)) \in \Gamma$. Let T be the tangent space at this point, and N be the normal space (i.e. $N = \{ v \in \mathbb{R}^{d+n} \mid u \cdot v = 0 \ \forall u \in T \} \}.$
 - (a) Show that $\dim(T) = d$.
 - (b) Find a basis of N explicitly in terms of $\nabla f_1(a), \ldots, \nabla f_n(a)$. [Note, for n = 1, we showed $(\nabla f(a), -1)$ was a basis of N. Figure out the right generalization of this.]
 - 5. Use the proof outline from class to give a full ε - δ proof of the chain rule.

Assignment 5 (assigned 2020-02-12, due 2020-02-19).

- 1. Do questions 3, 4 and 5 from homework 4.
- 2. If $f : \mathbb{R}^d \to \mathbb{R}$ is C^2 define $\Delta f = \sum_{i=1}^d \partial_i^2 f$ (this is called the Laplacian of f).
 - (a) Suppose $u, v : \mathbb{R}^2 \to \mathbb{R}$ are C^2 functions such that $\partial_x u = \partial_y v$ and $\partial_y u = -\partial_x v$. Show that $|\nabla u|^2 = |\nabla v|^2$, $\nabla u \cdot \nabla v = 0$, and $\Delta u = \Delta v = 0$.
 - (b) If $f : \mathbb{R}^2 \to \mathbb{R}$ is C^2 , and let $g = \Delta f$. Compute $\Delta(f(u, v))$ in terms of g, u and v. [If you do it correctly, your answer will only involve g and $|\nabla u|^2$.]

In class we stated (but didn't prove) the fact that if mixed partials are equal they must be continuous. Here's a proof

- 3. Let $f : \mathbb{R}^2 \to \mathbb{R}$ be a C^2 function. Suppose h, k > 0 and let $\delta = \delta(h, k)$ be defined by $\delta(h, k) = f(x + h, y + k) f(x, y + k) f(x + h, y) + f(x, y)$.
 - (a) Use the mean value theorem to show that there exists $\theta \in (0, h)$ so that

$$\delta(h,k) = h(\partial_x f(x+\theta, y+k) - \partial_x f(x+\theta, y))$$

- (b) Conclude $\delta(h,k) = hk\partial_y\partial_x f(x+\theta,y+\eta)$ for some $\theta \in (0,h)$ and $\eta \in (0,k)$.
- (c) Show that $\lim_{(h,k)\to(0,0)} \frac{\delta(h,k)}{hk} = \partial_y \partial_x f(x,y)$, provided $\partial_y \partial_x f$ is continuous.
- (d) Similarly, show $\lim_{(h,k)\to(0,0)} \frac{\delta(h,k)}{hk} = \partial_x \partial_y f(x,y)$, if $\partial_x \partial_y f$ is continuous.
- (e) Conclude $\partial_x \partial_y f = \partial_y \partial_x f$ if $f \in C^2$.

Optional problems, and details from class I left for you to check.

- * Suppose $u, v : \mathbb{R}^3 \to \mathbb{R}^3$ are two C^1 functions. Let $\alpha, \beta \in \mathbb{R}$ be two constants and A be a 3×3 (constant) matrix. Prove the following identities. [You don't need the limit definition, or ε - δ 's for even one of these. You can do them all by using partial derivatives and your one variable differentiation rules.]
 - (a) D(au + bv) = aDu + bDv(b) D(Au) = A(Du).(c) $\nabla (u \cdot v) = (Du)^T v + (Dv)^T u$ (d) $\partial_i (u \times v) = \partial_i u \times v + u \times \partial_i v$
- * (Mean value theorem) Let $f : \mathbb{R}^d \to \mathbb{R}$ be a differentiable function. For any $a, b \in \mathbb{R}^d$ show that there exists ξ on the line segment joining a and b such that $f(b) f(a) = (b a) \cdot \nabla f(\xi)$.

Assignment 6 (assigned 2020-02-19, due 2020-02-26).

- 1. Let $f : \mathbb{R}^d \to \mathbb{R}$ be C^2 . Show that f is convex if and only if its Hessian is always positive semi-definite. [Recall, a function $f : \mathbb{R}^d \to \mathbb{R}$ is *convex* if for every $x, y \in \mathbb{R}^d$ and $\theta \in [0, 1]$ we have $f(\theta x + (1 \theta)y) \leq \theta f(x) + (1 \theta)f(y)$.]
- 2. Let $a, b, c \in \mathbb{R}$ be such that $ac b^2 \neq 0$. Find all critical points of $ax^2 + 2bxy + cy^2$. Find conditions on a, b, c that would classify this as a local minimum, maximum or saddle.
- 3. Find the critical points of each of these functions. For each critical point, determine whether it is a local maximum, local minimum, saddle or neither.

(a)
$$\frac{x}{x^2+y^2}$$
 (c) $\sin x \cosh y$

(b)
$$[x^2 + (y+1)^2][x^2 + (y-1)^2]$$
 (d) $x^2 - 2xy + y^2$

4. (a) Suppose $f : \mathbb{R}^2 \to \mathbb{R}$ is a C^2 function. Compute

$$\lim_{h \to 0} \frac{1}{4h^2} (f(x+h,y+h) - f(x-h,y+h) + f(x-h,y-h) - f(x+h,y-h))$$

and express your answer in terms of derivatives of f.

(b) Do the same for

$$\lim_{h \to 0} \frac{1}{h^2} (f(x+h,y) + f(x,y+h) + f(x-h,y) + f(x,y-h) - 4f(x,y)).$$

- 5. Let $f : \mathbb{R}^d \to \mathbb{R}$ be a C^3 function, $a, h \in \mathbb{R}^d$ and define g(t) = f(a + th).
 - (a) Using the third order (one dimensional) Taylor expansion for g(1) g(0), show that there exists ξ on the line segment joining a and a + h such that

$$f(a+h) = f(a) + \nabla f(a) \cdot h + \frac{1}{2}(Hf_ah) \cdot h + \frac{1}{6}\sum_{i,j,k=1}^d \partial_i \partial_j \partial_k f(\xi) h_i h_j h_k$$

(b) Conclude there exists a function $R_3(h)$ such that

$$f(a+h) = f(a) + \sum \partial_i f(a)h_i + \frac{1}{2} \sum_{i,j=1}^d \partial_i \partial_j f(a)h_i h_j + \frac{1}{6} \sum_{i,j,k=1}^d \partial_i \partial_j \partial_k f(a)h_i h_j h_k + R_3(h)$$

and $\lim_{h\to 0} R_3(h)/|h|^3 = 0$. [This can of course be continued inductively to obtain higher order Taylor expansions of f.]

Optional problems, and details from class I left for you to check.

- * (a) Let $A = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$. Compute the eigenvalues of A in terms of a, b, c.
 - (b) Show that A is positive semi-definite if and only if $a \ge 0$, $c \ge 0$ and $ac-b^2 \ge 0$.

Assignment 7 (assigned 2020-02-26, due 2020-03-04).

- 1. (Spherical coordinates) Let $V = \{(r, \theta, \phi) \mid r > 0, \theta \in (-\pi, \pi), \phi \in (0, \pi)\}$ and define $\varphi(r, \theta, \phi) = (x, y, z)$ where $x = r \sin \phi \cos \theta$, $y = r \sin \phi \sin \theta$ and $z = r \cos \phi$. (Geometrically, ϕ is the angle between (x, y, z) and the positive z-axis, and θ is the angle between the projection (x, y) and the positive x-axis.)
 - (a) Let $U \subseteq \mathbb{R}^3$ be the set of points where either x > 0 or $y \neq 0$. Show $\det(D\varphi) \neq 0$ in V, and conclude (by the inverse function theorem) that r, θ, φ can be (locally) expressed as differentiable functions of x, y, z.
 - (b) Explicitly express r, θ, ϕ as functions of x, y, z and show that $\varphi : V \to U$ is bijective (and hence a coordinate change function). Let $(r, \theta, \phi) = \psi(x, y, z)$ denote the inverse function. Compute det $D\psi$ explicitly and verify det $D\psi \neq 0$ in U.
 - (c) If f is a differentiable function compute $\partial_x f$, $\partial_y f$ and $\partial_z f$ in terms of $\partial_r f$, $\partial_\theta f$, $\partial_\phi f$, r, θ and ϕ .
 - (d) If g is a differentiable function compute $\partial_r g$, $\partial_{\theta} g$ and $\partial_{\phi} g$ in terms of $\partial_x g$, $\partial_y g$ and $\partial_z g$, x, y and z.
- 2. For each of the equations below near the given point, which variables can be solved for and expressed as differentiable functions of the remaining variables (according to the implicit function theorem). For each of these variables, compute all the partials at the given point. (That is, if you say w, x can be expressed as differentiable functions of y, z, compute $\partial_y w$, $\partial_z w$, $\partial_y x$ and $\partial_z x$ at the given point.)

(a)
$$x^2 + y^2 - \cos(xy) = 0$$
 near (1,0).
(b) $e^{xz} + y\sin(yz) + z = 0$ near (0,0,-1).
(c) $\sin(xy) + \sin(yz) + \sin(xz) = 0$ and $e^{xyz} + x + y + z = 2$, near (0,0,1).

- 3. Decide whether each of the following implicitly defined sets are curves or surfaces. At the given point, find a basis of the tangent space, and as many many linearly independent normal vectors as possible. Also find the tangent line (or tangent plane).
 - (a) $x\sin(x) = y + xe^y$ at (0,0).

(b)
$$\ln(xy) = y - x$$
 at (1,1)

- (c) $x \sin y + y \sin z + z \sin x = 0$ at $(0, \pi, 2\pi)$.
- (d) $z^2 = x^2 + y^2 1$ and 2(x 1) + y z = 0 at (1, 0, 0).
- 4. Let $f, g: \mathbb{R}^3 \to \mathbb{R}$ be two differentiable functions, $c, d \in \mathbb{R}$. Consider the two surfaces $\Gamma = \{x \in \mathbb{R}^3 \mid f(x) = c\}$ and $\Delta = \{x \in \mathbb{R}^3 \mid g(x) = d\}$. Suppose $C = \Gamma \cap \Delta$ is a curve, $a \in C$ and that the vectors $\nabla f(a)$ and $\nabla g(a)$ are linearly independent. Find the tangent space of C at a in terms of ∇f and ∇g . Verify your formula by explicitly computing it when $f(x, y, z) = z^2 x^2 y^2 + 1$ and g(x, y, z) = 2(x 1) + y z at the point (1, 0, 0).

Assignment 8 (assigned 2020-03-04, due 2020-03-18).

- 1. Parametrize the following curves. [For closed curves C, a parametrization is a C^1 function $\gamma: [0,1] \to C$ such that $\gamma' \neq 0$, $\gamma(0) = \gamma(1)$ and $\gamma: [0,1) \to C$ is bijective.]
 - (a) $\frac{(x-x_0)^2}{a^2} + \frac{(y-y_0)^2}{b^2} = 1$, where a, b > 0 are constants.
 - (b) $\cos x \cos y = 1/2$ for $x, y \in (-\frac{\pi}{2}, \frac{\pi}{2})$.
- 2. Let $\gamma : \mathbb{R} \to \mathbb{R}^d$ be a differentiable function so that $\gamma(t)$ represents the position of a particle at time t. Let $v(t) = \gamma'(t)$ denotes the velocity, and a(t) = v'(t)denote the acceleration. Note $\gamma(t) = (\gamma_1(t), \ldots, \gamma_d(t))$ and $\gamma'(t)$ is defined to be the vector function $(\gamma'_1(t), \ldots, \gamma'_d(t))$. We know that if γ parametrises a curve Γ , then v(t) is tangent to Γ at the point $\gamma(t)$. The acceleration a, however, need not be tangent to Γ , as we will see below.
 - (a) If |v(t)| = 1 for all t, show that a(t) and v(t) are perpendicular.

If when moving while keeping the *magnitude* of the velocity constant, a particle experiences acceleration it is because the path taken by the particle is curved. The sharper the curve, the more the acceleration experienced (think about the force you feel when driving around a sharp curve).

Definition: If a curve Γ is parametrized by the function γ , then we define the curvature at the point $\gamma(t)$ by $\kappa = \frac{1}{|\gamma'(t)|} |(\frac{\gamma'(t)}{|\gamma'(t)|})'| = \frac{1}{|v|} |(\frac{v}{|v|})'|$.

If $|\gamma'(t)| = 1$ for all t, then the curvature is exactly the magnitude of the acceleration. (Note, one can show that the curvature κ only depends on the curve, and not the parametrization.)

- (b) Compute the curvature at any point on a circle of radius R.
- (c) Compute the curvature of the curve $y^2 = x^2 1$ at the point (x, 1).
- (d) If d = 3, show $\kappa^2 = |v|^{-6} (|a|^2 |v|^2 (a \cdot v)^2)$, and hence conclude $\kappa = \frac{|a \times v|}{|v|^3}$.
- 3. (a) Let $f : \mathbb{R} \to \mathbb{R}$ is differentiable, and $\Gamma = \{(x, f(x)) \mid x \in \mathbb{R}\}$ be the graph of f. Show that the curvature of Γ is $|f''|/(1 + (f')^2)^{3/2}$.
 - (b) Let $g : \mathbb{R}^2 \to \mathbb{R}$ be C^2 , $c \in \mathbb{R}$ and Γ be the curve $\{(x, y) \mid g(x, y) = c\}$. Suppose further $\nabla g \neq 0$ on the curve Γ . Show that the curvature of Γ is

$$\frac{\partial_x^2 g(\partial_y g)^2 + \partial_y^2 g(\partial_x g)^2 - 2\partial_x \partial_y g \partial_x g \partial_y g|}{|\nabla g|^3}$$

HINT: Let $(a, b) \in \Gamma$, and suppose $\partial_y g(a, b) \neq 0$. Use the implicit function theorem to write Γ as the curve y = f(x) near the point (a, b). Now express f'' in terms of derivatives of g by implicit differentiation, and use the previous part.

- 4. Let $S \subseteq \mathbb{R}^3$ be the surface $z^2 = x^2 + y^2 1$, and $(a, b, c) \in S$. There are exactly two lines that pass through (a, b, c) that are *completely contained* in the surface S. Find them. Express them as $\ell_i = \{(a, b, c) + t(u_i, v_i, w_i) \mid t \in \mathbb{R}\}$ for $i \in \{1, 2\}$ and two linearly independent vectors (u_1, v_1, w_1) and (u_2, v_2, w_2) . [HINT: What is the intersection of S with it's tangent plane? Start with (a, b, c) = (1, 0, 0).]
- 5. (Optional challenge) If Γ is any curve contained on the sphere $x^2 + y^2 + z^2 = 1$, then show that the curvature at any point on Γ is at least 1.

Assignment 9 (assigned 2020-03-18, due Never).

In light of your Midterm on 3/25, this homework is not due. However, many of the problems are good practice, and I recommend doing (or at least thinking) about them before your exam. Some of these problems will find their way onto your next homework.

1. The plane x + y + 2z = 2 intersects the paraboloid $z = x^2 + y^2$ in an ellipse. Find the points on this ellipse that are nearest to and farthest from the origin.

Sometimes when maximising a function in the region $\{g \leq c\}$, the maximum (or minimum) could be attained on the boundary $\{g = c\}$. In this case you can of course find all *interior* local maxima and minima by solving $\nabla f = 0$ and looking at Hf. For maxima and minima on the boundary, it is often convenient to use Lagrange multipliers.

- 2. Find the absolute maxima and minima of e^{-xy} on the set $x^2 + 4y^2 \leq 1$.
- 3. For each of the curves below, find the local maxima and minima of the curvature.
 - (a) A circle of radius R.
 - (b) A straight line.
 - (c) The parabola $y = x^2$.
 - (d) The ellipse $x^2/a^2 + y^2/b^2 = 1$, for a, b > 0.
 - (e) The curve $\cos x \cos y = 1/2$ in the region $x, y \in (-\pi/2, \pi/2)$.
- 4. Maximise the volume of an open box given the surface area is $3a^2$. (That is, maximise xyz under the constraint $xy + 2(yz + zx) = 3a^2$.)
- 5. Maximise the volume of a cylinder given that the total surface area is $6\pi a^2$. [For fun, check if the proportions of your optimal cylinder agrees with your standard coke can; if not, write to Coco-cola with a proposal to save money and the environment...]
- 6. Let p, q > 1 be such that 1/p + 1/q = 1. In the region x, y > 0 maximise xy subject to the constraint $x^p/p + y^q/q = C$. Use this to give a different proof of Young's inequality from homework 3.
- 7. (Cauchy-Schwartz inequality) If $x, y \in \mathbb{R}^n$ show $|x \cdot y| \leq |x||y|$. [HINT: Maximise $x \cdot y$ subject to the constraint |x| = a and |y| = b.]

Assignment 10 (assigned 2020-03-25, due 2020-04-01).

- 1. Do question 2 from assignment 9.
- 2. Suppose $f: \mathbb{R}^{n+d} \to \mathbb{R}$ is C^2 , and $g: \mathbb{R}^{n+d} \to \mathbb{R}^n$ is C^1 . Let $M = \{g = 0\}$, and assume for all $x \in M$ we have rank $(Dg_x) = n$. Suppose further f attains a constrained local minimum at a, subject to the constraint g = 0.
 - (a) True or false: Hf_a must necessarily be positive semi-definite (here Hf_a is the Hessian of f at the point a.) Prove it, or find a counter example.
 - (b) True or false: For every u in the tangent space of M at the point a, we must have $u \cdot (Hf_a u) \ge 0$. Prove it, or find a counter example.
 - (c) Redo the previous part with the additional assumption that g is linear.

3. Let $f: \mathbb{R}^d \to \mathbb{R}^m$ and $g: \mathbb{R}^d \to \mathbb{R}^n$ be C^1 . Let $c \in \mathbb{R}^m$, $d \in \mathbb{R}^n$, $M = \{f = c\}$, and $N = \{g = d\}$. Suppose for all $x \in M$, $y \in N$ we have $\operatorname{rank}(Df_x) = m$ and $\operatorname{rank}(Dg_y) = n$. Let $a \in M$ and $b \in N$ be two points that minimize the distance between M and N (i.e. $|x - y| \ge |a - b|$ whenever $x \in M$ and $y \in N$). True or false: a - b is orthogonal to the tangent space of M at a, and also orthogonal to the tangent space of N at b. Prove it, or find a counter example.

- 4. In each of the cases below, find $\int_B f(x, y) dA$.
 - (a) $f(x,y) = 16 x^2 y^2$, and R is the triangle with vertices (0,0), (1,1) and (2,1).
 - (b) f(x,y) = y, where R is bounded by the curves $x = y^2$ and y = x 2.
 - (c) $f(x, y) = x^2 + 2y$, where R is bounded by the curves y = x, $y = x^3$ in the region $x \ge 0$.
- 5. Compute the integrals integrals:

(a)
$$\int_{x=-1}^{1} \int_{y=1}^{2} \sin(xy^2) \, dy \, dx.$$
 (b) $\int_{y=0}^{1} \int_{x=-\cos^{-1} y}^{\cos^{-1} y} e^{\sin x} \, dx \, dy.$

[Just for fun, try plugging these into a computer. Most computer algebra systems won't be able to evaluate these integrals symbolically.]

6. (Optional) Compute both iterated integrals of the function

$$f(x,y) = \frac{x^2 - y^2}{(x^2 + y^2)^2}$$

on the region $[0,1] \times [0,1]$. Are they equal? [HINT: First compute $\partial_x \partial_y \tan^{-1}(y/x)$.]

Assignment 11 (assigned 2020-04-01, due 2020-04-08).

- 1. Evaluate $\int_{B} f(x, y, z) dV$ in the following cases:
 - (a) $f(x, y, z) = x^2 y^2 z$ where R is the cylinder $x^2 + y^2 < 1, 0 < z < 1$.
 - (b) $f(x, y, z) = x^2 + z^2$, where R is the pyramid with vertices $(\pm 1, \pm 1, 0)$ and (0, 0, 1).
- 2. (a) Let $D = B(0,1) \subseteq \mathbb{R}^2$ be the two dimensional disk of radius 1 and center (0,0). For what $p \in \mathbb{R}$ is $\int_D \frac{1}{|x|^p} dA < \infty$. [Here $x = (x_1, x_2) \in \mathbb{R}^2$.]
 - (b) Let $S = B(0,1) \subseteq \mathbb{R}^3$ be the three dimensional sphere with radius 1 and center (0,0,0). For what $p \in \mathbb{R}$ is $\int_S \frac{1}{|x|^p} dV < \infty$. [Here $x = (x_1, x_2, x_3) \in \mathbb{R}^3$.]
- 3. Let $D \subseteq \mathbb{R}^2$ represent an irregular plate whose density is given by $\rho(x, y)$. Let $\ell \subseteq \mathbb{R}^2$ be a straight line, representing a knife edge upon which D is balanced. The magnitude of the *torque* experienced when D is balanced on ℓ is given by $T_{\ell} = \int_D \rho d \, dx \, dy$. Here $d = p \cdot \hat{n}$, where \hat{n} is a unit vector perpendicular to ℓ , and p = p(x, y) is the vector from (x, y) to the closest point on ℓ . The plate D will balance on a knife edge along ℓ if (and only if) $T_{\ell} = 0$.
 - (a) Given $a \in \mathbb{R}$, let $\ell = \{(a,t) \mid t \in \mathbb{R}\}$. Show that $T_{\ell} = 0$ if and only if $a = (\int_D x \rho \, dA) / \int_D \rho dA$.
 - (b) Given $b \in \mathbb{R}$ let $\ell = \{(t, b) \mid t \in \mathbb{R}\}$. Show that $T_{\ell} = 0$ if and only if $b = (\int_D y\rho \, dA) / \int_D \rho dA$.

The point (a, b) above for which $T_{\ell} = 0$ is called the *center of mass* of D.

- (c) Find the center of mass of the triangle with vertices (0,0), (a,b), (0,c) that has a uniform density. [Assume 0 < b < c and a > 0.]
- (d) If ℓ is any line (not necessarily parallel to the coordinate axis) passing through the center of mass of D must $T_{\ell} = 0$? Prove or disprove.
- 4. Let $U \subseteq \mathbb{R}^2$ be a (bounded) oddly shaped region containing the origin. Define $C \subseteq \mathbb{R}^3$ to be the cone with vertex (0, 0, h) and base U given by

$$C = \left\{ (x, y, z) \mid 0 < z < h, \left(\frac{hx}{h-z}, \frac{hy}{h-z}\right) \in U \right\}$$

Find vol(C) in terms of h and area(U). [HINT: Let u = hx/(h-z), v = hy/(h-z) and w = z, and transform $\int_C 1 dx dy dz$ into u, v, w coordinates.]

- 5. Compute $\int_{\mathbb{R}^2} \frac{1}{y^2 + 4\sqrt{x^2 + y^2} + 4} \frac{dx \, dy}{\sqrt{x^2 + y^2}}$. [HINT: Let $x = u^2 v^2$ and y = 2uv and change coordinates.]
- 6. Compute $\int_0^\infty \frac{\sin x}{x} dx$.

HINT: Substitute $1/x = \int_0^\infty e^{-xy} dy$ above and switch the order of integration. This is one situation where the hypothesis of Fubini's theorem *won't* be satisfied; however, flipping the order of the integrals can still be justified using other methods. Extra kudos for justifying it!

Assignment 12 (assigned 2020-04-08, due 2020-04-15).

- 1. A non-uniform wire is bent along the semi-circle $x^2 + y^2 = 1$ with $y \ge 0$. The density of the wire is given by $\rho(x, y) = 2 y$. Find the total mass and the *center of mass* of the wire. Is the center of mass on the wire? [You've derived a formula for the center of mass of a plate previously. See if you can use your intuition from there to guess a formula for the center of mass in this context.]
- 2. Let $S \subseteq \mathbb{R}^3$ be a surface whose boundary is the closed curve Γ . Amperes law says that the *total current* passing through the surface S is given by $\frac{1}{\mu_0} \oint_{\Gamma} B \cdot d\ell$, where μ_0 is the magnetic constant and $B : \mathbb{R}^3 \to \mathbb{R}^3$ is the magnetic field. Compute the current through the rectangle with (-1, -1, -1), (-1, 1, -1), (1, 1, 1) and (1, -1, 1) (traversed in the stated order) when

$$B(x, y, z) = \frac{1}{\sqrt{x^2 + y^2}} \begin{pmatrix} -y \\ x \\ 0 \end{pmatrix}.$$

3. Let $\Gamma \subseteq \mathbb{R}^2$ be a closed, piecewise C^1 curve that does not pass through the origin. Define the *winding number* of Γ about the origin to be

$$W(\Gamma) = \frac{1}{2\pi} \oint_{\Gamma} \frac{-y \, dx + x \, dy}{x^2 + y^2}$$

- (a) Compute the winding number of a circle of radius R and center (0,0), traversed counter clockwise. What happens if you traverse it clockwise instead?
- (b) Compute the winding number of a circle of radius 1 and center (0, 2) traversed in either direction.

The remainder of this problem is devoted to showing that the winding number is an integer. Let $\gamma : [0,1] \to \Gamma$ be a parametrization of Γ and write $\gamma(t) = (x(t), y(t))$. Assume y(0) = 0 and x(0) > 0 for simplicity. Let t_1 be the first time when $x(t_1) = 0$. Note $y(t_1) \neq 0$, since $(0,0) \notin \Gamma$, and so we let t_2 be the first time after t_1 when $y(t_2) = 0$. Again $x(t_2) \neq 0$, so let t_3 be the first time after t_2 when $x(t_3) = 0$. Continue this, and assume that after finitely many steps we obtain an even number N so that $t_N = 1$ and $y(t_N) = y(0) = 0$.

(c) Show that
$$\int_{t_i}^{t_{i+1}} \frac{-y(t)x'(t) + x(t)y'(t)}{x(t)^2 + y(t)^2} dt = \pm \frac{\pi}{2}.$$

(d) If *i* is even, show
$$\int_{t_i}^{t_{i+2}} \frac{-y(t)x'(t) + x(t)y'(t)}{x(t)^2 + y(t)^2} dt = \pm \pi \text{ if } \operatorname{sign}(x_i x_{i+2}) < 0$$
and 0 otherwise.

- (e) Show $W(\Gamma) \in \mathbb{Z}$.
- 4. (Optional Challenge that is worth a reward) If $\Gamma \subseteq \mathbb{R}^3$ is a C^2 closed curve, show that $\oint_{\Gamma} \kappa |d\ell|$ is multiple of 2π . Here κ is the curvature defined in your previous homework. [This is called the *turning number* of the curve Γ .]

Assignment 13 (assigned 2020-04-15, due 2020-04-22).

- 1. Let Γ be the curve parametrized by $\gamma(t) = (a\cos^3 t, a\sin^3 t)$, for $t \in [0, 2\pi]$. 1. (a) Let $U \subseteq \mathbb{R}^2$ be a domain, and $f: U \to \mathbb{R}$ be C^1 , and Σ be the graph of fCompute the area of the region enclosed by Γ .
- 2. Let $P \subseteq \mathbb{R}^2$ be a (not necessarily convex) polygon whose vertices, ordered counter clockwise, are $(x_1, y_1), \ldots, (x_N, y_N)$. Show that

area(P) =
$$\frac{(x_1y_2 - x_2y_1) + (x_2y_3 - x_3y_2) + \dots + (x_Ny_1 - x_1y_N)}{2}$$

[This is called the *surveyors formula*. While the statement involves only elementary coordinate geometry, it isn't as easy to prove directly this way. Hint: Use Greens theorem.]

3. Let $U, V \subset \mathbb{R}^2$ be domains, $\varphi: U \to V$ be C^2 , and $F: V \to \mathbb{R}^2$ be C^1 . Define $G: U \to \mathbb{R}^2$ by $G(x) = (D\varphi_x)^T (F \circ \varphi(x))$. Show that

$$\partial_1 G_2 - \partial_2 G_1 = [(\partial_1 F_2 - \partial_2 F_1) \circ \varphi] \det(D\varphi)$$

[This was a detail in class used in the proof of Greens theorem.]

- 4. Let $U \subseteq \mathbb{R}^2$ be a bounded domain such that ∂U is the finite union of closed, piecewise C^1 curve.
 - (a) Given $x \in \partial U$, let $\hat{n}(x)$ be a unit vector that is normal to ∂U and pointing outwards to the domain U. If $F: \overline{U} \to \mathbb{R}^2$ is C^1 , show that $\oint_{\partial U} F \cdot \hat{n} |d\ell| =$ $\int_{U} \nabla \cdot F \, dA. \ (\text{Here } \nabla \cdot F = \partial_1 F_1 + \partial_2 F_2.)$
 - (b) Suppose $0 \notin \partial U$, and U has no holes (i.e. ∂U is connected). What is the winding number of ∂U oriented counter clockwise? Prove it. [Hint 1: Let $V = \ln|x|, F = \nabla V$, and express the winding number in terms of $\oint_{\partial U} \nabla F \cdot \hat{n} |d\ell|$. Hint 2: Divide into two cases: $0 \in U$, and $0 \notin U$. Hint 3: The case $0 \notin U$ is easier.]
- 5. Let a > b > 0, and Σ be the torus obtained by rotating a circle with center (a, 0, 0)and radius b about the z axis. Parametrize the surface, and evaluate the surface integral that computes area(Σ).

Optional problems, and details from class I left for you to check.

* In each of the following cases, show that

$$\int_{U} (\partial_x Q - \partial_y P) \, dA \neq \oint_{\partial U} P \, dx + Q \, dy.$$

Also explain why this does not contradict Greens theorem.

(a) $U = \{(x, y) | y > 0\}, P = \frac{1}{1 + x^2}, \text{ and } Q = 0.$ (b) $U = B(0,1) \subseteq \mathbb{R}^2$, $P = \frac{-y}{x^2 + y^2}$, and $Q = \frac{x}{x^2 + y^2}$. Assignment 14 (assigned 2020-04-22, due 2020-04-29).

(i.e. $\Sigma = \{(x, y, z) \mid (x, y) \in U, \& z = f(x, y)\}$). Show that

area
$$(\Sigma) = \int_U \sqrt{1 + (\partial_x f)^2 + (\partial_y f)^2} \, dA.$$

- (b) Use the previous part to compute the surface area of a sphere of radius r. [The next problem provides an easier way to do this as well.]
- 2. (a) Let $f:[a,b] \to (0,\infty)$ be C^1 , and $\Sigma \subset \mathbb{R}^3$ be the surface formed by rotating the graph of f about the x-axis. Explicitly,

$$\Sigma = \{ (x, y, z) \mid x \in [a, b] \text{ and } y^2 + z^2 = f(x)^2 \}.$$

Show that
$$\operatorname{area}(\Sigma) = 2\pi \int_{a}^{b} f(x) \sqrt{1 + f'(x)^2} \, dx.$$

- (b) Find the surface area of a cylinder with base radius r and height h.
- (c) Find the surface area of a cone with base radius r and height h.
- 3. Let $\Sigma = \{(x, y, z) \mid y^2 + z^2 = 1, -1 < x < 1 \& z > 0\}$, and $F = e_3$. Compute $\int_{\Sigma} F \cdot \hat{n} \, dS$, where at any point on Σ , \hat{n} is the upward pointing unit normal.
- 4. (a) Let A be a 3×3 matrix, and $u, v \in \mathbb{R}^3$. Show $Au \times Av = \operatorname{adj}(A)^T (u \times v)$. where $\operatorname{adj}(A)$ is the adjugate of the matrix A.
 - (b) Let $U \subseteq \mathbb{R}^3$ be a domain, (Σ, \hat{n}) be an oriented surface. Let $\psi : \mathbb{R}^3 \to \mathbb{R}^3$ be an injective C^1 function such that $\det(D\psi) > 0$ on all of U, and $F: \psi(\Sigma) \to \mathbb{R}^3$ be a vector field. Show that

$$\int_{\psi(\Sigma)} F \cdot \hat{n} \, dS = \int_{\Sigma} (\operatorname{adj}(D\psi)(F \circ \psi)) \cdot \hat{n} \, dS$$

5. Let $N(x) = -1/(4\pi |x|)$, and Σ be a closed surface that encloses the origin. Compute $\int_{\Sigma} \nabla N \cdot \hat{n} \, dS$.

Optional problems, and details from class I left for you to check.

* (Greens identity) Let $U \subseteq \mathbb{R}^3$ be a bounded domain whose boundary is a C^1 surface. If $f, g: U \to \mathbb{R}$ be C^2 , show

$$\int_{U} (f\Delta g) \, dV = \int_{\partial U} f\nabla g \cdot \hat{n} \, dS - \int_{U} (\nabla f \cdot \nabla g) \, dV.$$

Recall $\Delta g = \sum_i \partial_i^2 g = \nabla \cdot (\nabla g)$, and by convention \hat{n} is the *outward pointing* unit normal on ∂U .

* (Challenge) Suppose $u: \mathbb{R}^3 \to \mathbb{R}$ is a C^2 function such that $\Delta u = 0$. For any $a \in \mathbb{R}^3$, r > 0 show that $u(a) = \int_{\partial B(a,r)} u \, dS$. [HINT: Use Greens identity and the function N. Google "Mean value property" if you get stuck.]

Assignment 15 (assigned 2020-04-26, due Never).

- 1. Let $u, v : \mathbb{R}^3 \to \mathbb{R}^3$ be C^2 vector fields, $f, g : \mathbb{R}^3 \to \mathbb{R}$ be C^2 scalar functions. Prove the following identities.
 - (a) $\nabla(fg) = f\nabla g + g\nabla f$. (b) $\nabla \cdot (fu) = (\nabla f) \cdot u + f\nabla \cdot u$. (c) $\nabla \times (fu) = f\nabla \times u + (\nabla f) \times u$ (d) $\nabla \times (u \times v) = u(\nabla \cdot v) - v(\nabla \cdot u) + (v \cdot \nabla)u - (u \cdot \nabla)v$ (e) $\nabla \cdot (u \times v) = (\nabla \times u) \cdot v - u \cdot (\nabla \times v)$ (f) $\nabla \times (\nabla u) = 0$ (g) $\nabla \cdot (\nabla \times u) = 0$ (h) $\nabla \times (\nabla \times u) = -\Delta u + \nabla(\nabla \cdot u)$
- 2. Let $F = (2x, y^2, z^2)$.
 - (a) Compute $\int_{\Sigma} F \cdot \hat{n} \, dS$, where $\Sigma \subseteq \mathbb{R}^3$ is the sphere of radius 1.
 - (b) Compute $\oint_{\Gamma} F \cdot d\ell$, where Γ is the intersection of Σ above and the plane x + 2y + 3z = 0.
- 3. In each of the following cases show that

$$\int_{U} (\nabla \cdot v) \, dV \neq \int_{\partial U} v \cdot \hat{n} \, dS$$

Also explain why this does not contradict the divergence theorem.

(a) $U = B(0, 1) \subseteq \mathbb{R}^3$, and $v = x/|x|^4$.

(b)
$$U = \{x \in \mathbb{R}^3 \mid z > x^2 + y^2\}$$
, and $v(x, y, z) = (0, 0, e^{-z})$.

4. (a) Suppose $U \subseteq \mathbb{R}^3$ is a bounded domain with piecewise C^1 boundary. If $u: \bar{U} \to \mathbb{R}^3$ and $f: \bar{U} \to \mathbb{R}$ are C^1 functions, then show that

$$\int_{U} \nabla f \cdot u \, dV = \int_{\partial U} f u \cdot \hat{n} \, dS - \int_{U} f \nabla \cdot u \, dV$$

(b) If $f: \mathbb{R}^3 \to \mathbb{R}$ and $u: \mathbb{R}^3 \to \mathbb{R}^3$ are two C^1 functions such that $|f(x)u(x)| \leq C/|x|^{2+\varepsilon}$ for some constants $C, \varepsilon > 0$ and all $x \notin B(0,1)$, then show that

$$\int_{\mathbb{R}^3} \nabla f \cdot u \, dV = - \int_{\mathbb{R}^3} f \, \nabla \cdot u \, dV$$

5. Suppose $U, V \subseteq \mathbb{R}^3$ are domains, $\varphi : U \to V$ is C^2 and $v : V \to \mathbb{R}^3$ is C^1 . Define $w = \operatorname{adj}(D\varphi)(v \circ \varphi) : U \to \mathbb{R}^3$. Show that

$$\nabla \cdot w = \det(D\varphi)(\nabla \cdot v) \circ \varphi.$$

[This was a detail used in the proof of the divergence theorem.]

6. (Optional challenge) Suppose $u : \mathbb{R}^3 \to \mathbb{R}^3$ is C^1 . Show that $\nabla \cdot u = 0$ if and only if there exists a C^1 vector field $v : \mathbb{R}^3 \to \mathbb{R}^3$ such that $u = \nabla \times v$. [HINT: If you get stuck this is done at the very end of the lecture notes.]