

Putnam $\Sigma.13$

Po-Shen Loh

17 November 2012

1 Problems

Putnam 1991/A4. Does there exist an infinite sequence of closed discs D_1, D_2, D_3, \dots in the plane, with centers c_1, c_2, c_3, \dots , respectively, such that

1. the c_i have no limit point in the finite plane,
2. the sum of the areas of the D_i is finite, and
3. every line in the plane intersects at least one of the D_i ?

Putnam 1991/A5. Find the maximum value of

$$\int_0^y \sqrt{x^4 + (y - y^2)^2} dx$$

for $0 \leq y \leq 1$.

Putnam 1991/A6. Let $A(n)$ denote the number of sums of positive integers

$$a_1 + a_2 + \dots + a_r$$

which add up to n with

$$\begin{aligned} a_1 &> a_2 + a_3, a_2 > a_3 + a_4, \dots, \\ a_{r-2} &> a_{r-1} + a_r, a_{r-1} > a_r. \end{aligned}$$

Let $B(n)$ denote the number of $b_1 + b_2 + \dots + b_s$ which add up to n , with

1. $b_1 \geq b_2 \geq \dots \geq b_s$,
2. each b_i is in the sequence $1, 2, 4, \dots, g_j, \dots$ defined by $g_1 = 1$, $g_2 = 2$, and $g_j = g_{j-1} + g_{j-2} + 1$, and
3. if $b_1 = g_k$ then every element in $\{1, 2, 4, \dots, g_k\}$ appears at least once as a b_i .

Prove that $A(n) = B(n)$ for each $n \geq 1$.

(For example, $A(7) = 5$ because the relevant sums are $7, 6+1, 5+2, 4+3, 4+2+1$, and $B(7) = 5$ because the relevant sums are $4+2+1, 2+2+2+1, 2+2+1+1+1, 2+1+1+1+1+1, 1+1+1+1+1+1+1$.)