

## 2. Polynomials

Po-Shen Loh

CMU Putnam Seminar, Fall 2019

### 1 Classical results

**Algebra.** If  $r$  is a root of the polynomial  $P(x)$ , then  $P$  factors as  $(x - r)Q(x)$  for some polynomial  $Q$ .

**Algebra.** Every polynomial of degree  $n$  has at most  $n$  distinct roots.

**Lagrange Interpolation.** Show that there is a degree-4 polynomial which takes values  $P(0) = 0$ ,  $P(1) = 0$ ,  $P(2) = 0$ ,  $P(3) = 1$ , and  $P(4) = 1$ .

**Reed-Solomon codes.** Automatic spell checkers know to correct “teh” to “the”. More abstractly, an *error-correcting code* with minimum distance  $d$  is a collection of strings of length  $n$  from an alphabet  $A$ , with the property that any two strings differ by at least  $d$  pointwise edits. It turns out that there are nice error-correcting codes with minimum distance  $d$  over alphabets of size  $q$ , for prime powers  $q$ , and these are based on polynomials!

**Multiple roots.** If  $r$  is a real root of the polynomial  $P(x)$ , and  $r$  has multiplicity greater than 1, then both  $P(r) = 0$  and  $P'(r) = 0$ .

**Gauss-Lucas.** The zeros of the derivative  $P'(z)$  of any polynomial lie in the convex hull of the zeros of the polynomial  $P(z)$ .

### 2 Problems

1. Find a polynomial with integer coefficients that has the zero  $\sqrt{2} + \sqrt[3]{3}$ .
2. There is no polynomial which has the property that  $P(k) = 2^k$  for all positive integers  $k$ .
3. Let  $a_1, \dots, a_n$  be positive real numbers. Prove that the polynomial  $P(x) = x^n - a_1x^{n-1} - a_2x^{n-2} - \dots - a_n$  has a unique positive zero.
4. Solve the system

$$\begin{aligned}x + y + z &= 1 \\xyz &= 1\end{aligned}$$

knowing that  $x, y, z$  are complex numbers of absolute value equal to 1.

5. Let  $P(z)$  and  $Q(z)$  be polynomials with complex coefficients of degree greater than or equal to 1 with the property that  $P(z) = 0$  if and only if  $Q(z) = 0$  and  $P(z) = 1$  if and only if  $Q(z) = 1$ . Prove that the polynomials are equal.

6. Let  $P(x)$  and  $Q(x)$  be arbitrary polynomials with real coefficients, and let  $d$  be the degree of  $P(x)$ . Assume that  $P(x)$  is not the zero polynomial. Prove that there exist polynomials  $A(x)$  and  $B(x)$  with real coefficients, such that:
- (i) both  $A$  and  $B$  have degree at most  $d/2$ , and
  - (ii) at most one of  $A$  and  $B$  is the zero polynomial, and
  - (iii)  $\frac{A(x)+Q(x)B(x)}{P(x)}$  is a polynomial with real coefficients. That is, there is some polynomial  $C(x)$  with real coefficients such that  $A(x) + Q(x)B(x) = P(x)C(x)$ .

### 3 Homework

Please write up solutions to two of the problems, to turn in at next week's meeting. One of them may be a problem that we discussed in class. You are encouraged to collaborate with each other. Even if you do not solve a problem, please spend two hours thinking, and submit a list of your ideas.