Putnam $\Sigma.13$

Po-Shen Loh

17 November 2019

1 Problems

- **Putnam 1998/A4.** Let $A_1 = 0$ and $A_2 = 1$. For n > 2, the number A_n is defined by concatenating the decimal expansions of A_{n-1} and A_{n-2} from left to right. For example $A_3 = A_2A_1 = 10$, $A_4 = A_3A_2 = 101$, $A_5 = A_4A_3 = 10110$, and so forth. Determine all n such that 11 divides A_n .
- **Putnam 1998/A5.** Let \mathcal{F} be a finite collection of open discs in \mathbb{R}^2 whose union contains a set $E \subseteq \mathbb{R}^2$. Show that there is a pairwise disjoint subcollection D_1, \ldots, D_n in \mathcal{F} such that

 $E \subseteq \bigcup_{j=1}^{n} 3D_j.$

Here, if D is the disc of radius r and center P, then 3D is the disc of radius 3r and center P.

Putnam 1998/A6. Let A, B, C denote distinct points with integer coordinates in \mathbb{R}^2 . Prove that if

 $(|AB| + |BC|)^2 < 8 \cdot [ABC] + 1$

then A, B, C are three vertices of a square. Here |XY| is the length of segment XY and [ABC] is the area of triangle ABC.