Putnam $\Sigma.8$

Po-Shen Loh

18 October 2020

1 Problems

- **Putnam 2012/A4.** Let q and r be integers with q > 0, and let A and B be intervals on the real line. Let T be the set of all b + mq where b and m are integers with b in B, and let S be the set of all integers a in A such that ra is in T. Show that if the product of the lengths of A and B is less than q, then S is the intersection of A with some arithmetic progression.
- **Putnam 2012/A5.** Let \mathbb{F}_p denote the field of integers modulo a prime p, and let n be a positive integer. Let v be a fixed vector in \mathbb{F}_p^n , let M be an $n \times n$ matrix with entries of \mathbb{F}_p , and define $G : \mathbb{F}_p^n \to \mathbb{F}_p^n$ by G(x) = v + Mx. Let $G^{(k)}$ denote the k-fold composition of G with itself, that is, $G^{(1)}(x) = G(x)$ and $G^{(k+1)}(x) = G(G^{(k)}(x))$. Determine all pairs p, n for which there exist v and M such that the p^n vectors $G^{(k)}(0)$, $k = 1, 2, ..., p^n$ are distinct.
- **Putnam 2012/A6.** Let f(x, y) be a continuous, real-valued function on \mathbb{R}^2 . Suppose that, for every rectangular region R of area 1, the double integral of f(x, y) over R equals 0. Must f(x, y) be identically 0?