## Putnam $\Sigma.13$

Po-Shen Loh

27 November 2022

## 1 Problems

- **Putnam 2012/A4.** Let q and r be integers with q > 0, and let A and B be intervals on the real line. Let T be the set of all b + mq where b and m are integers with b in B, and let S be the set of all integers a in A such that ra is in T. Show that if the product of the lengths of A and B is less than q, then S is the intersection of A with some arithmetic progression.
- **Putnam 2012/A5.** Let  $\mathbb{F}_p$  denote the field of integers modulo a prime p, and let n be a positive integer. Let v be a fixed vector in  $\mathbb{F}_p^n$ , let M be an  $n \times n$  matrix with entries of  $\mathbb{F}_p$ , and define  $G : \mathbb{F}_p^n \to \mathbb{F}_p^n$ by G(x) = v + Mx. Let  $G^{(k)}$  denote the k-fold composition of G with itself, that is,  $G^{(1)}(x) = G(x)$ and  $G^{(k+1)}(x) = G(G^{(k)}(x))$ . Determine all pairs p, n for which there exist v and M such that the  $p^n$ vectors  $G^{(k)}(0)$ ,  $k = 1, 2, ..., p^n$  are distinct.
- **Putnam 2012/A6.** Let f(x, y) be a continuous, real-valued function on  $\mathbb{R}^2$ . Suppose that, for every rectangular region R of area 1, the double integral of f(x, y) over R equals 0. Must f(x, y) be identically 0?