

# Putnam E.3

Po-Shen Loh

14 Sep 2016

## 1 Problems

**Putnam 1987/A1.** Curves  $A$ ,  $B$ ,  $C$ , and  $D$  are defined in the plane as follows:<sup>1</sup>

$$\begin{aligned} A &= \left\{ (x, y) : x^2 - y^2 = \frac{x}{x^2 + y^2} \right\}, \\ B &= \left\{ (x, y) : 2xy + \frac{y}{x^2 + y^2} = 3 \right\}, \\ C &= \left\{ (x, y) : x^3 - 3xy^2 + 3y = 1 \right\}, \\ D &= \left\{ (x, y) : 3x^2y - 3x - y^3 = 0 \right\}. \end{aligned}$$

Prove that  $A \cap B = C \cap D$ .

**Putnam 1987/A2.** The sequence of digits

123456789101112131415161718192021...

is obtained by writing the positive integers in order. If the  $10^n$ -th digit in this sequence occurs in the part of the sequence in which the  $m$ -digit numbers are placed, define  $f(n)$  to be  $m$ . For example,  $f(2) = 2$  because the 100th digit enters the sequence in the placement of the two-digit integer 55. Find, with proof,  $f(1987)$ .

**Putnam 1987/A3.** For all real  $x$ , the real-valued function  $y = f(x)$  satisfies

$$y'' - 2y' + y = 2e^x.$$

- (a) If  $f(x) > 0$  for all real  $x$ , must  $f'(x) > 0$  for all real  $x$ ? Explain.
- (b) If  $f'(x) > 0$  for all real  $x$ , must  $f(x) > 0$  for all real  $x$ ? Explain.

---

<sup>1</sup>The equations defining  $A$  and  $B$  are indeterminate at  $(0, 0)$ . The point  $(0, 0)$  belongs to neither.