

# Putnam $\Sigma.10$

Po-Shen Loh

30 October 2016

## 1 Problems

**Putnam 1994/A4.** Let  $A$  and  $B$  be  $2 \times 2$  matrices with integer entries such that  $A, A+B, A+2B, A+3B,$  and  $A+4B$  are all invertible matrices whose inverses have integer entries. Show that  $A+5B$  is invertible and that its inverse has integer entries.

**Alternative: Putnam 1940/B6 (if you've seen A4 before).** The  $n \times n$  matrix  $(m_{ij})$  is defined as  $m_{ij} = a_i a_j$  for  $i \neq j$ , and  $a_i^2 + k$  for  $i = j$ . Show that  $\det(m_{ij})$  is divisible by  $k^{n-1}$  and find its other factor.

**Putnam 1994/A5.** Let  $(r_n)_{n \geq 0}$  be a sequence of positive real numbers such that  $\lim_{n \rightarrow \infty} r_n = 0$ . Let  $S$  be the set of numbers representable as a sum

$$r_{i_1} + r_{i_2} + \cdots + r_{i_{1994}},$$

with  $i_1 < i_2 < \cdots < i_{1994}$ . Show that every nonempty interval  $(a, b)$  contains a nonempty subinterval  $(c, d)$  that does not intersect  $S$ .

**Putnam 1994/A6.** Let  $f_1, \dots, f_{10}$  be bijections of the set of integers such that for each integer  $n$ , there is some composition  $f_{i_1} \circ f_{i_2} \circ \cdots \circ f_{i_m}$  of these functions (allowing repetitions) which maps 0 to  $n$ . Consider the set of 1024 functions

$$\mathcal{F} = \{f_1^{e_1} \circ f_2^{e_2} \circ \cdots \circ f_{10}^{e_{10}}\},$$

$e_i = 0$  or  $1$  for  $1 \leq i \leq 10$ . ( $f_i^0$  is the identity function and  $f_i^1 = f_i$ .) Show that if  $A$  is any nonempty finite set of integers, then at most 512 of the functions in  $\mathcal{F}$  map  $A$  to itself.