

# Putnam $\Sigma.6$

Po-Shen Loh

1 October 2017

## 1 Problems

**Putnam 1992/B4.** Let  $p(x)$  be a nonzero polynomial of degree less than 1992 having no nonconstant factor in common with  $x^3 - x$ . Let

$$\frac{d^{1992}}{dx^{1992}} \left( \frac{p(x)}{x^3 - x} \right) = \frac{f(x)}{g(x)}$$

for polynomials  $f(x)$  and  $g(x)$ . Find the smallest possible degree of  $f(x)$ .

**Putnam 1992/B5.** Let  $D_n$  denote the value of the  $(n-1) \times (n-1)$  determinant

$$\begin{bmatrix} 3 & 1 & 1 & 1 & \cdots & 1 \\ 1 & 4 & 1 & 1 & \cdots & 1 \\ 1 & 1 & 5 & 1 & \cdots & 1 \\ 1 & 1 & 1 & 6 & \cdots & 1 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 1 & 1 & \cdots & n+1 \end{bmatrix}.$$

Is the set  $\left\{ \frac{D_n}{n!} \right\}_{n \geq 2}$  bounded?

**Putnam 1992/B6.** Let  $\mathcal{M}$  be a set of real  $n \times n$  matrices such that

- (i)  $I \in \mathcal{M}$ , where  $I$  is the  $n \times n$  identity matrix;
- (ii) if  $A \in \mathcal{M}$  and  $B \in \mathcal{M}$ , then either  $AB \in \mathcal{M}$  or  $-AB \in \mathcal{M}$ , but not both;
- (iii) if  $A \in \mathcal{M}$  and  $B \in \mathcal{M}$ , then either  $AB = BA$  or  $AB = -BA$ ;
- (iv) if  $A \in \mathcal{M}$  and  $A \neq I$ , there is at least one  $B \in \mathcal{M}$  such that  $AB = -BA$ .

Prove that  $\mathcal{M}$  contains at most  $n^2$  matrices.