

Putnam $\Sigma.12$

Po-Shen Loh

15 November 2020

1 Problems

Putnam 2013/A4. A finite collection of digits 0 and 1 is written around a circle. An *arc* of length $L \geq 0$ consists of L consecutive digits around the circle. For each arc w , let $Z(w)$ and $N(w)$ denote the number of 0's in w and the number of 1's in w , respectively. Assume that $|Z(w) - Z(w')| \leq 1$ for any two arcs w, w' of the same length. Suppose that some arcs w_1, \dots, w_k have the property that

$$Z = \frac{1}{k} \sum_{j=1}^k Z(w_j) \text{ and } N = \frac{1}{k} \sum_{j=1}^k N(w_j)$$

are both integers. Prove that there exists an arc w with $Z(w) = Z$ and $N(w) = N$.

Putnam 2013/A5. For $m \geq 3$, a list of $\binom{m}{3}$ real numbers a_{ijk} ($1 \leq i < j < k \leq m$) is said to be *area definite* for \mathbb{R}^n if the inequality

$$\sum_{1 \leq i < j < k \leq m} a_{ijk} \cdot \text{Area}(\Delta A_i A_j A_k) \geq 0$$

holds for every choice of m points A_1, \dots, A_m in \mathbb{R}^n . For example, the list of four numbers $a_{123} = a_{124} = a_{134} = 1, a_{234} = -1$ is area definite for \mathbb{R}^2 . Prove that if a list of $\binom{m}{3}$ numbers is area definite for \mathbb{R}^2 , then it is area definite for \mathbb{R}^3 .

Putnam 2013/A6. Define a function $w : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$ as follows. For $|a|, |b| \leq 2$, let $w(a, b)$ be as in the table shown; otherwise, let $w(a, b) = 0$.

$w(a, b)$	b					
	-2	-1	0	1	2	
a	-2	-1	-2	2	-2	-1
	-1	-2	4	-4	4	-2
	0	2	-4	12	-4	2
	1	-2	4	-4	4	-2
	2	-1	-2	2	-2	-1

For every finite subset S of $\mathbb{Z} \times \mathbb{Z}$, define

$$A(S) = \sum_{(\mathbf{s}, \mathbf{s}') \in S \times S} w(\mathbf{s} - \mathbf{s}').$$

Prove that if S is any finite nonempty subset of $\mathbb{Z} \times \mathbb{Z}$, then $A(S) > 0$. (For example, if $S = \{(0, 1), (0, 2), (2, 0), (3, 1)\}$, then the terms in $A(S)$ are 12, 12, 12, 12, 4, 4, 0, 0, 0, 0, -1, -1, -2, -2, -4, -4.)