

Putnam $\Sigma.3$

Po-Shen Loh

6 September 2026

1 Problems

Putnam 2006/A4. Let $S = \{1, 2, \dots, n\}$ for some integer $n > 1$. Say a permutation π of S has a local maximum at $k \in S$ if

- (i) $\pi(k) > \pi(k+1)$ for $k = 1$;
- (ii) $\pi(k-1) < \pi(k)$ and $\pi(k) > \pi(k+1)$ for $1 < k < n$;
- (iii) $\pi(k-1) < \pi(k)$ for $k = n$.

(For example, if $n = 5$ and π takes values at $1, 2, 3, 4, 5$ of $2, 1, 4, 5, 3$, then π has a local maximum of 2 at $k = 1$, and a local maximum of 5 at $k = 4$.) What is the average number of local maxima of a permutation of S , averaging over all permutations of S ?

Putnam 2006/A5. Let n be a positive odd integer and let θ be a real number such that θ/π is irrational. Set $a_k = \tan(\theta + k\pi/n)$, $k = 1, 2, \dots, n$. Prove that

$$\frac{a_1 + a_2 + \dots + a_n}{a_1 a_2 \dots a_n}$$

is an integer, and determine its value.

Putnam 2006/A6. Four points are chosen uniformly and independently at random in the interior of a given circle. Find the probability that they are the vertices of a convex quadrilateral.