

HANF NUMBER OF THE FIRST STABILITY CARDINAL IN AECs

SAMSON LEUNG

ABSTRACT

We show that $\beth_{(2^{\text{LS}(\mathbf{K}))_+}$ is the lower bound to the Hanf numbers for the length of the order property and for stability in stable abstract elementary classes (AECs). Our examples satisfy the joint embedding property, no maximal model, $(< \aleph_0)$ -tameness but not necessarily the amalgamation property. We also define variations on the order and syntactic order properties by allowing the index set to be linearly ordered rather than well-ordered. Combining with Shelah's stability theorem, we deduce that our examples can have the order property up to any $\mu < \beth_{(2^{\text{LS}(\mathbf{K}))_+}$. Boney conjectured that the need for joint embedding property for two type-counting lemmas is necessary. We solved the conjecture by showing it is independent of ZFC. Using Galois Morleyization, we give syntactic proofs to known stability results assuming a monster model.

TABLE OF CONTENTS

1	Introduction	2
2	Preliminaries	3
3	Stability and no order property	8
4	Lower bound for stability and no order property*	11
5	Galois Morleyization and syntactic order property	16
6	Shelah's stability theorem	20
7	Syntactic splitting	24

Date: October 7, 2021

AMS 2020 Subject Classification: Primary 03C48. Secondary: 03C45, 03C55. Key words and phrases. Abstract Elementary Classes; Stability; First stability cardinal; Tameness; Hanf number.

1 INTRODUCTION

Semantic order properties (Definition 3.2) in abstract elementary classes (AECs) are defined in terms of (semantic) Galois types instead of formulas. They are analogs to syntactic order properties in first-order and infinitary logics. In [She72], Shelah showed that in $L_{\lambda^+, \omega}$ the (syntactic) order property of length $\beth_{(2^\lambda)^+}$ implies the order property of arbitrary length. In [GS86], Grossberg and Shelah introduced the Hanf number of the order property of $L_{\lambda^+, \omega}$ and later [GS98, Theorem 2.8] gave a lower bound as \beth_{λ^+} . These bound the Hanf number of order property between \beth_{λ^+} and $\beth_{(2^\lambda)^+}$. However, the example for the lower bound does not readily generalize to (semantic) order properties of AECs. Shelah [She99, Claim 4.6] hinted that the upper bound of the order property in AECs is $\beth_{(2^{\text{LS}(\mathbf{K}))}^+}$ but it was not known whether it is tight. We present examples (Corollary 6.8) that $\beth_{(2^{\text{LS}(\mathbf{K}))}^+}$ is exact. Our examples satisfy the joint embedding property, no maximal model and ($< \aleph_0$)-tameness but the amalgamation property fails. It is open whether the bound can be lowered when one assumes the amalgamation property.

Vasey [Vas16b] extended Shelah [She71], Grossberg and Lessmann's [GL02] results to AECs and showed that assuming the amalgamation property and tameness, the first stability cardinal is bounded above by $\beth_{(2^{\text{LS}(\mathbf{K}))}^+}$. It is open whether this bound can be lowered under the amalgamation property. Our examples, which do not satisfy the amalgamation property, show that the lower bound in general is at least $\beth_{(2^{\text{LS}(\mathbf{K}))}^+}$. From instability, we can apply Vasey's techniques (which are based on [She09b, V.A.]) to derive the order property. This provides an alternative way other than finding the witnesses directly. It is open whether the amalgamation property can lower the bound for the first stability cardinal.

Vasey's result above relies on one direction of [Bon17, Theorem 3.1], which does not use the joint embedding property. The other direction involves lemmas that assume the joint embedding property, which Boney suspected to be necessary. As a side product of our construction, we show in Corollary 4.4 that the need for the joint embedding property is independent of ZFC; and we find an example and a counterexample under different set theoretic assumptions.

In Section 2, we state our notations and definitions. We also give a shorter proof to

Boney’s result to motivate Corollary 4.4. In Section 3, we review results concerning stability and the order property. We give more details for the proof of [She99, Claim 4.6]. In Section 4, we construct our main examples in Proposition 4.1 which set a lower bound to stability and a variation of the order property for stable AECs. The variation of the order property is slightly more general by allowing the index set to be linear ordered rather than well-ordered. We will also show Corollary 4.4 as a side product of our construction. In Section 5, we apply the same variation to the syntactic order property which can be combined with Galois Morleyization. We give analogs to Vasey’s results with our variation on the order property. In Section 6, we write down the details of Vasey’s observation [Vas16b, Fact 4.10] that Shelah’s results in [She09b, V.A.] can be applied to AECs under Galois Morleyization. It allows us to deduce (Corollary 6.8) the order property up to $\beth_{(2^{\text{LS}(\mathbf{K})})^+}$ in our examples in Proposition 4.1, without finding explicit witnesses. We also apply such technique to bound the first stability cardinal under extra hypotheses. In Section 7, we use Galois Morleyization to recover common stability results where types can be over sets under $\text{LS}(\mathbf{K})$. Vasey in [Vas16b, Section 5] has done similarly for coheir while we will work on splitting instead. In particular we prove Theorem 2.10 syntactically which is needed for Vasey’s upper bound to the first stability cardinal, under the amalgamation property and tameness.

This paper was written while the author was working on a Ph.D. under the direction of Rami Grossberg at Carnegie Mellon University and we would like to thank Prof. Grossberg for his guidance and assistance in my research in general and in this work in particular. We also thank John Baldwin, Hanif Cheung, Marcos Mazari-Armida and Wentao Yang for useful comments.

2 PRELIMINARIES

We assume some familiarity with AECs, for example [Bal09, Chapter 4]. We will use $\kappa, \lambda, \mu, \chi$ to denote cardinals, α, β, γ to denote ordinals, n for natural numbers. We define κ^- to be the predecessor cardinal (if it exists) or κ itself. When we write $\alpha - n$, we assume $\alpha = \beta + n$ for some ordinal β .

Let $\mathbf{K} = \langle K, \leq_{\mathbf{K}} \rangle$ be an AEC. If the context is clear, we write \leq in place of $\leq_{\mathbf{K}}$. We

abbreviate *AP* the *amalgamation property*, *JEP* the *joint embedding property* and *NMM* *no maximal model*. For $M \in K$, write $|M|$ the universe of M and $\|M\|$ the cardinality of M . For Galois types (orbital types) of length $(< \alpha)$, we denote them as $\text{gS}^{<\alpha}(\cdot)$ where the argument can be a set A in some model $M \in K$. In general $\text{gS}^{<\alpha}(A) := \bigcup \{ \text{gS}^{<\alpha}(A; M) : M \in K, |M| \supseteq A \}$ (under *AP*, the choice of M does not matter). \mathbf{K} is $(< \alpha)$ -*stable in* λ if for any set A in some model $M \in K$, $|A| \leq \lambda$, then $|\text{gS}^{<\alpha}(A; M)| \leq \lambda$. We omit “ $(< \alpha)$ ” if $\alpha = 2$, while we omit “in λ ” if there exists such a $\lambda \geq \text{LS}(\mathbf{K})$. Similarly \mathbf{K} is α -*stable in* λ if for any set A in some model $M \in K$, $|\text{gS}^\alpha(A)| \leq \lambda$. Tameness will be defined in Definition 5.1. We allow stability and tameness under $\text{LS}(\mathbf{K})$, especially in Section 7.

Given a $\forall\exists$ theory T and a set of $L(T)$ -types Γ , $EC(T, \Gamma)$ is the class of models of T such that Γ is not realized by any elements. If we order $EC(T, \Gamma)$ by L -substructures, it forms an AEC with $\text{LS}(\mathbf{K}) = |L(T)|$. $\delta(\lambda, \kappa)$ is the least ordinal δ such that: for any T, Γ with $|L(T)| \leq \lambda$, $|\Gamma| \leq \kappa$, $\{P, <\} \subseteq L(T)$ where P is a unary predicate, $<$ is a linear order on P , if there is a model $M \in EC(T, \Gamma)$ whose $(P^M, <^M)$ has order type $\geq \delta$, then there is a model $N \in EC(T, \Gamma)$ whose $(P, <)$ is not well-ordered.

Recall the classical theorem: notice in the proof that X can be a linear order while Y' can be its suborder.

Theorem 2.1 (Erdős-Rado Theorem). *Let λ be an infinite cardinal. For $n < \omega$,*

$$\beth_n(\lambda)^+ \rightarrow (\lambda^+)_\lambda^{n+1}$$

In other words, for any $|X| \geq \beth_n(\lambda)^+$, any $f : [X]^{n+1} \rightarrow \lambda$, there is $X' \subseteq X$ such that $|X'| \geq \lambda^+$ and $f \upharpoonright [X']^{n+1}$ is constant.

Proof. We adapt the proof in [Mar02, Theorem 5.1.4] because it does not require the set X to be well-ordered. We prove by induction: When $n = 0$, the statement is $\lambda^+ \rightarrow (\lambda^+)_\lambda^1$. Let X be of size $\geq \lambda^+$. We need to color its elements with λ -many colors. By pigeonhole principle, it is possible to find $X' \subseteq X$ of size $\geq \lambda^+$ such that $f \upharpoonright X'$ is constant.

Assume the statement is true for $n - 1$. We need to show $\beth_n(\lambda)^+ \rightarrow (\lambda^+)_\lambda^{n+1}$. Let X be of size $\beth_n(\lambda)^+$, $f : [X]^{n+1} \rightarrow \lambda$. For $x \in X$, define $f_x : [X - \{x\}]^n \rightarrow \lambda$ by $f_x(Y) := f(Y \cup \{x\})$. We build $\langle X_\alpha : \alpha < \beth_{n-1}(\lambda)^+ \rangle$ increasing and continuous subsets of

X such that for $\alpha < \beth_n(\lambda)^+$, $|X_\alpha| = \beth_n(\lambda)$. For the base step, take any $X_0 \subseteq X$ of size $\beth_n(\lambda)$. Suppose X_α is constructed, we build $X_{\alpha+1}$ satisfying:

1. $X_\alpha \subseteq X_{\alpha+1} \subseteq X$
2. $|X_{\alpha+1}| = \beth_n(\lambda)$
3. For any $Y \subseteq X_\alpha$ of size $\beth_{n-1}(\lambda)$, any $x \in X - Y$, there is $x' \in X_{\alpha+1} - Y$ such that $f_x \upharpoonright [Y]^n = f_{x'} \upharpoonright [Y]^n$.

The above is possible by a counting argument: the number of possible Y is

$$|X_\alpha|^{\beth_{n-1}(\lambda)} = \beth_n(\lambda)^{\beth_{n-1}(\lambda)} = (2^{\beth_{n-1}(\lambda)})^{\beth_{n-1}(\lambda)} = 2^{\beth_{n-1}(\lambda)} = \beth_n(\lambda).$$

Given Y , the number of possible $h : [Y]^n \rightarrow \lambda$ is bounded by

$$\lambda^{\beth_{n-1}(\lambda)} = 2^{\beth_{n-1}(\lambda)} = \beth_n(\lambda)$$

Therefore, it suffices to add $\beth_n(\lambda) \cdot \beth_n(\lambda) = \beth_n(\lambda)$ -many witnesses from X to X_α . Define $X' = \bigcup \{X_\alpha : \alpha < \beth_{n-1}(\lambda)^+\}$. Notice that $|X'| = \beth_n(\lambda) < |X|$. For any $Y \subseteq X'$ of size $\beth_{n-1}(\lambda)$, by a cofinality argument $Y \subseteq X_\alpha$ for some $\alpha < \beth_{n-1}(\lambda)^+$. So for any $x \in X - Y$, there is $x' \in X_{\alpha+1} - Y \subseteq X' - Y$ such that $f_x \upharpoonright [Y]^n = f_{x'} \upharpoonright [Y]^n$.

Pick any $x \in X - X'$ and build $Y = \{y_\alpha : \alpha < \beth_{n-1}(\lambda)^+\} \subseteq X'$ such that $f_{y_\alpha} \upharpoonright [\{y_\beta : \beta < \alpha\}]^n = f_x \upharpoonright [\{y_\beta : \beta < \alpha\}]^n$ ($y_0 \in X'$ can be any element). By inductive hypothesis on Y and f_x , we can find $Y' \subseteq Y$ of size $\geq \lambda^+$ such that $f_x \upharpoonright [Y']^n$ is constant. We check that Y' is as desired: let $A \in [Y']^{n+1}$ and write $A = \{y_{\alpha_1}, \dots, y_{\alpha_{n+1}}\}$ where $\alpha_1 < \dots < \alpha_{n+1} < \beth_{n-1}(\lambda)^+$.

$$f(A) = f_{y_{\alpha_{n+1}}}(A - \{y_{\alpha_{n+1}}\}) = f_x(A - \{y_{\alpha_{n+1}}\})$$

which is constant because f_x is constant on $[Y']^n \ni A - \{y_{\alpha_{n+1}}\}$. □

The following Theorem 2.2 and Theorem 2.10 are only used in the proof of Corollary 6.6(1). We will streamline Boney's proof of Lemma 2.9 by omitting the ambient models (otherwise it would involve a lot of bookkeeping and direct limits). We will clarify the

relationship between Lemma 2.9 and Theorem 2.2 and show that *JEP* in Theorem 2.2 is not needed. If we work in a monster model \mathfrak{C} , we can also allow stability over sets (of size $< \text{LS}(\mathbf{K})$), but we keep the original formulation to state Remark 2.6 more clearly.

Theorem 2.2. *Let \mathbf{K} be an AEC and $\lambda \geq \text{LS}(\mathbf{K})$. Suppose \mathbf{K} has λ -AP and is stable in λ . For any ordinal $\alpha \geq 1$ with $\lambda^{|\alpha|} = \lambda$, \mathbf{K} is α -stable in λ .*

The requirement $\lambda^{|\alpha|} = \lambda$ cannot be improved: let $\lambda^{|\alpha|} > \lambda$, take \mathbf{K} be the well-orderings of type at most λ and $\leq_{\mathbf{K}}$ by initial segments. Then it is stable in λ because there are only λ -many elements in the unique maximal model (which witnesses AP). It is not α -stable because each element has different types, so the number of α -types is exactly $\lambda^{|\alpha|} > \lambda$.

We will prove the theorem through a series of lemmas. We may assume α to be a cardinal $\kappa = |\alpha|$. Denote $\text{gS}_\lambda^1 := \sup\{|\text{gS}^1(M)| : M \in K, \|M\| = \lambda\}$ and similarly $\text{gS}_\lambda^\kappa := \sup\{|\text{gS}^\kappa(M)| : M \in K, \|M\| = \lambda\}$.

Lemma 2.3. *Suppose $\kappa \geq \lambda$, then $(\text{gS}_\lambda^1)^\kappa = \text{gS}_\lambda^\kappa = 2^\kappa$.*

Proof. $2^\kappa \leq (\text{gS}_\lambda^1)^\kappa \leq (2^\lambda)^\kappa = 2^\kappa$. Pick any $\|M\| = \lambda$ and two distinct elements $\{a, b\}$ from $|M|$. Form binary sequences from $\{a, b\}$ of length κ , which shows $2^\kappa \leq \text{gS}_\lambda^\kappa \leq 2^{\lambda+\kappa} = 2^\kappa$. \square

Lemma 2.4. [Bon17, Proposition 2.7] *Suppose $\kappa \leq \lambda$. If in addition \mathbf{K} has λ -JEP and $\text{cf}(\text{gS}_\lambda^\kappa) \leq \lambda$, then there is $M \in K$, $\|M\| = \lambda$ such that $|\text{gS}^\kappa(M)| = \text{gS}_\lambda^\kappa$.*

Proof. Pick $\langle M_i : i < \mu \rangle$ (not necessarily increasing) witnessing $\mu := \text{cf}(\text{gS}_\lambda^\kappa) \leq \lambda$. By λ -AP and λ -JEP, obtain M of size λ such that $M \geq M_i$ for all M_i . $|\text{gS}^\kappa(M)| \geq \sup_{i < \mu} |\text{gS}^\kappa(M_i)| = \text{gS}_\lambda^\kappa$. \square

Lemma 2.5. [Bon17, Theorem 3.2] *Suppose $\kappa \leq \lambda$. If in addition \mathbf{K} has λ -JEP, then $(\text{gS}_\lambda^1)^\kappa \leq \text{gS}_\lambda^\kappa$.*

Proof. Given $M \in K$ of size λ . We show that $|\text{gS}^\kappa(M)| \geq |\text{gS}^1(M)|^\kappa$, which does not use λ -JEP. By λ -AP, pick $N \geq M$ (perhaps of size greater than λ) such that N realizes $\text{gS}(M)$, say by $\langle a_i : i < |\text{gS}(M)| \rangle$. Form sequences of length κ from the a_i , there are $|\text{gS}(M)|^\kappa$ -many sequences. They realize distinct types in $\text{gS}^\kappa(M)$ by checking each coordinate.

Suppose $\text{cf}(\text{gS}_\lambda^1) \leq \kappa$, then $\text{cf}(\text{gS}_\lambda^1) \leq \lambda$. Substitute $\kappa = 1$ in Lemma 2.4 (which uses λ -JEP), there is $M^* \in K_\lambda$ such that $|\text{gS}^1(M^*)| = \text{gS}_\lambda^1$. Thus $(\text{gS}_\lambda^1)^\kappa = |\text{gS}^1(M^*)|^\kappa \leq |\text{gS}^\kappa(M^*)| \leq \text{gS}_\lambda^\kappa$.

Now suppose $\text{cf}(\text{gS}_\lambda^1) > \kappa$, then a cofinality argument shows the second equality below:

$$\begin{aligned} (\text{gS}_\lambda^1)^\kappa &:= (\sup\{|\text{gS}^1(M)| : M \in K, \|M\| = \lambda\})^\kappa \\ &= \sup\{|\text{gS}^1(M)|^\kappa : M \in K, \|M\| = \lambda\} \\ &\leq \sup\{|\text{gS}^\kappa(M)| : M \in K, \|M\| = \lambda\} =: \text{gS}_\lambda^\kappa \end{aligned}$$

□

Remark 2.6. After [Bon17, Proposition 2.7], Boney suggested that λ -JEP might be necessary. We will show in Corollary 4.4 that the need of λ -JEP is independent of ZFC for the above two lemmas.

Question 2.7. In [Bon17, Proposition 2.7], there is an alternative hypothesis to Lemma 2.4 where $\text{cf}(\text{gS}_\lambda^\kappa) \leq \lambda$ is replaced by a stronger assumption $I(K, \lambda) \leq \lambda$. Would λ -JEP be necessary in this case or is it again independent of ZFC? An answer would shed light on the relationship between stability and the number of nonisomorphic models.

Lemma 2.8. [Bon17, Theorem 3.5] *Suppose $\kappa \leq \lambda$, then $(\text{gS}_\lambda^1)^\kappa \geq \text{gS}_\lambda^\kappa$.*

Proof. First we describe the proof strategy: for a fixed model M , we show that $\text{gS}^\kappa(M)$ is bounded above by $(\text{gS}_\lambda^1)^\kappa$. To do so, we build a gS_λ^1 -branching tree of models of height κ and list the possible 1-types of each model. For each κ -type in $\text{gS}^\kappa(M)$, we map it injectively to a branch of the tree (which is a sequence in $(\text{gS}_\lambda^1)^\kappa$), according to the 1-types of the elements from that sequence.

Let $\mu := \text{gS}_\lambda^1$. Fix an arbitrary $M \in K$ with $\|M\| = \lambda$. Write $\text{gS}^\kappa(M) = \langle p_k : k < \chi \rangle$ where p_k are distinct. Fix $\bar{a}_k := \{a_k^\alpha : \alpha < \kappa\} \models p_k$. Construct a tree of models $\langle M_\nu \in K_\lambda : \nu \in \mu^{<\kappa} \rangle$ as follows: $M_\emptyset := M$, take union at limit stages. Suppose M_ν is built for some $\nu \in \mu^{<\kappa}$. Enumerate without repetition $\text{gS}^1(M_\nu) = \langle q_i^\nu : i < \chi_\nu \rangle$ for some $\chi_\nu \leq \mu$. For $i < \chi_\nu$, define $M_{\nu \smallfrown i} \in K_\lambda$ with $M_{\nu \smallfrown i} \geq M_\nu$ and containing some $c_i^\nu \models q_i^\nu$. For $\chi_\nu \leq i < \mu$ (if there is any), give a default value to $M_{\nu \smallfrown i} := M_\nu$. Now we map each $p_k \in \text{gS}^\kappa(M)$ to

$\eta_k \in \mu^\kappa$ as follows: suppose $\nu := \eta_k \upharpoonright \alpha$ has been defined for some $\alpha < \kappa$, we set $\eta_k[\alpha]$ to be the minimum $i < \chi_\nu$ (which is the same as requiring $i < \mu$) such that a_k^α realizes q_i^ν . In other words, we decide the α -th element of the branch based on the type of the α -th element of \bar{a}_k over the current node (model).

It remains to check that the map is injective. Let $k < \chi$, we build $\langle f_\alpha : \alpha \leq \kappa \rangle$ increasing and continuous such that f_α maps a_k^β to $c_{\eta_k[\beta]}^{\eta_k \upharpoonright \beta}$ for all $\beta \leq \alpha$ while fixing $|M|$. Take $f_{-1} := \text{id}_M$ and we handle the successor case: suppose f_α has been constructed. There is $g : a_k^\alpha \mapsto c_{\eta_k[\alpha]}^{\eta_k \upharpoonright \alpha}$ fixing $|M_{\eta_k \upharpoonright \alpha}| \supseteq \{c_{\eta_k[\beta]}^{\eta_k \upharpoonright \beta} : \beta < \alpha\}$ by type equality. Let $f_{\alpha+1} := g \circ f_\alpha$. Now f_κ witnesses that \bar{a}_k and $\langle c_{\eta_k[\alpha]}^{\eta_k \upharpoonright \alpha} : \alpha < \kappa \rangle$ realize the same type over M . Since the latter sequence only depends on the coordinates of η_k , our map $p_k \mapsto \eta_k$ is injective. Therefore, $|\text{gS}^\kappa(M)| \leq \mu^\kappa$. Since M is arbitrary, $\text{gS}_\lambda^\kappa \leq \mu^\kappa = (\text{gS}_\lambda^1)^\kappa$. \square

Lemma 2.9. [Bon17, Theorem 3.1] *Let \mathbf{K} be an AEC with λ -AP and λ -JEP. Let $\kappa \geq 1$ be a cardinal. Then $(\text{gS}_\lambda^1)^\kappa = \text{gS}_\lambda^\kappa$ where λ -JEP is used in the “ \leq ” direction.*

Proof. Combine Lemma 2.3, Lemma 2.5 and Lemma 2.8. \square

Proof of Theorem 2.2. Let $\kappa = |\alpha|$. By Lemma 2.9, $\lambda = \lambda^\kappa = (\text{gS}_\lambda^1)^\kappa \geq \text{gS}_\lambda^\kappa$ which shows that \mathbf{K} is κ -stable in λ . By reordering the index κ , \mathbf{K} is α -stable in λ . \square

Theorem 2.10. [GV06, Corollary 6,4] *Let μ be an infinite cardinal and \mathbf{K} be an AEC with AP. If \mathbf{K} is μ -tame and stable in μ , then \mathbf{K} is stable in all $\lambda = \lambda^\mu$.*

The original proof proceeds semantically and we will give a syntactic proof in Section 7, allowing stability over sets which can be of size $< \text{LS}(\mathbf{K})$.

3 STABILITY AND NO ORDER PROPERTY

To find the upper bound of the first stability cardinal in stable complete first-order theories, one possible way is to establish:

Fact 3.1 (Shelah). Let T be a complete first-order theory. The following are equivalent:

1. T is stable.
2. For all $\lambda = \lambda^{|T|}$, T is stable in λ .

3. T has no (syntactic) order property of length ω .

$2^{|T|}$ is an upper bound for the first stability cardinal. Notice that in showing (3), compactness is used to stretch the order property to arbitrary length. In AECs, we can use the Hanf number to bound the (Galois) order property length. The following definition is based on [She99, Definition 4.3] and [Vas16b, Definition 4.3]:

Definition 3.2. Let μ be an infinite cardinal, $\alpha \geq 2$ and $\beta \geq 1$ be ordinals.

1. K has the β -order property of length μ if there exists some $\langle a_i : i < \mu \rangle \subseteq M \in K$ such that $l(a_i) = \beta$, and for $i_0 < i_1 < \mu$, $j_0 < j_1 < \mu$, $\text{gtp}(a_{i_0}a_{i_1}/\emptyset, M) \neq \text{gtp}(a_{j_0}a_{j_1}/\emptyset, M)$.
2. K has the $(< \alpha)$ -order property of length μ if there is a $\beta < \alpha$ witnessing (1).
3. K has the $(< \alpha)$ -order property if for all μ , K has the $(< \alpha)$ -order property of length μ . In other words, if we fix μ , we can find a suitable β_μ witnessing (1).
4. K has the *no* $(< \alpha)$ -order property if (3) fails. In other words, for each $\beta < \alpha$, there is an upper bound to the length of the β -order property. We omit $(< \alpha)$ if $\alpha = \omega$.

The above definition works fine if one wants an abstract generalization of the order property from the first-order version, in which case the length can be fixed at ω . However, in AECs, it is hard to construct long well-ordered sets without breaking stability or raising $\text{LS}(\mathbf{K})$. We propose the following definition instead:

Definition 3.3. In Definition 3.2, we replace all occurrences of “order property” by “order property*” if we also allow sequences indexed by linear orders instead of well-orderings. For example in (1), we say K has the β -order property* of length μ if there exist some linear order I , some $\langle a_i : i \in I \rangle \subseteq M \in K$ such that $|I| = \mu$, $l(a_i) = \beta$, and for $i_0 < i_1$ in I , $j_0 < j_1$ in I , $\text{gtp}(a_{i_0}a_{i_1}/\emptyset, M) \neq \text{gtp}(a_{j_0}a_{j_1}/\emptyset, M)$. When μ is omitted, we mean for all μ , there is a linear order I of cardinality μ witnessing the β -order property* of length μ .

In the following proposition, item (1) applies Morley’s method [Mor65] (see also [Bal09, Theorem A.3(2)]). The statement we use is from [She99, Claim 4.6] which only hinted at the proof of the Hanf number for arbitrarily large models [She90, VII.5]. We add more

details and explain how to adapt that proof. The proof of item (3) adapts the proof from [BGKV16, Fact 5.13].

Proposition 3.4. *Let \mathbf{K} be an AEC, $\beta \geq 2$ be an ordinal.*

1. *If for all $\mu < \beth_{(2^{<(\text{LS}(\mathbf{K})+|\beta|)})_+}$, \mathbf{K} has the $(< \beta)$ -order property of length μ , then \mathbf{K} has the $(< \beta)$ -order property (and the $(< \beta)$ -order property*).*
2. *If for all $\mu < \beth_{(2^{<(\text{LS}(\mathbf{K})+|\beta|)})_+}$, \mathbf{K} , \mathbf{K} has the $(< \beta)$ -order property* of length μ , then \mathbf{K} has the $(< \beta)$ -order property* (and $(< \beta)$ -order property).*
3. *If \mathbf{K} is $(< \beta)$ -stable (in some $\lambda \geq \text{LS}(\mathbf{K}) + |\beta|$), then there is $\mu < \beth_{(2^{<(\text{LS}(\mathbf{K})+|\beta|)})_+}$ such that \mathbf{K} has no $(< \beta)$ -order property* (and thus no $(< \beta)$ -order property) of length μ .*

Proof sketch. 1. We adapt the usual Hanf number argument. Suppose \mathbf{K} has the $(< \beta)$ -order property and we fix $\gamma < \beta$ such that \mathbf{K} has the γ -order property. By Shelah's Presentation Theorem, we may write $K = PC(T, \Gamma_1, L(\mathbf{K}))$ for some first-order theory T in $L \supseteq L(\mathbf{K})$ and some sets of L -types Γ_1 . Now we refer to the construction of [She90, VII Theorem 5.3] or [Gro21, Chapter 2 Theorem 6.35]. For each $\alpha < (2^{\text{LS}(\mathbf{K})+|\gamma|})_+$, instead of defining $F^{\text{ob}}(\alpha)$ to be some $M \in K$ of size \beth_α , we demand it to be the witness of the γ -order property of length \beth_α (we can also add another function $F_1^{\text{ob}}(\alpha, \cdot)$ to enumerate the elements of $F^{\text{ob}}(\alpha)$). At the end of the construction (which uses Erdős-Rado Theorem), we obtain an L -indiscernible sequence (of γ -tuples) $\langle a_i : i < \omega \rangle$ such that for $n < \omega$, $i_1 < i_2 < \dots < i_n < \omega$, the first-order type of $a_{i_1} \dots a_{i_n}$ is realized by some $d_1 \dots d_n$ that witness the order property. This induces an L -isomorphism between $EM(a_{i_1} \dots a_{i_n}) \cong EM(d_1 \dots d_n)$. Its reduct to $L(\mathbf{K})$ is also an isomorphism between $EM(a_{i_1} \dots a_{i_n}) \upharpoonright L(\mathbf{K}) \cong EM(d_1 \dots d_n) \upharpoonright L(\mathbf{K})$. Since the right-hand-side witnesses the order property in K , so is the left-hand-side. The same argument applies when the indiscernible sequence is stretched to arbitrary length (or any linear order).

2. The same proof of (1) goes through because Erdős-Rado Theorem applies to linear orders (actually any sets) besides well-orderings.

3. Otherwise by (1)(2), \mathbf{K} has the $(< \beta)$ -order property*. For any infinite cardinal $\lambda \geq \text{LS}(\mathbf{K}) + |\beta|$, let I_λ be a linear order of size $> \lambda$ such that it has a dense suborder J_λ of size λ . We stretch the indiscernible $\langle a_i : i < \omega \rangle$ in (2) to be indexed by I_λ . Pick $i < i'$ in I_λ , we can find k in J_λ such that $i < j < i'$. Then $\text{gtp}(a_i a_j / \emptyset) \neq \text{gtp}(a_{i'} a_j / \emptyset)$ are distinct by the order property, which means $\text{gtp}(a_i / a_j) \neq \text{gtp}(a_{i'} / a_j)$. This shows that \mathbf{K} is $(< \beta)$ -unstable in λ . As λ is arbitrary, \mathbf{K} is $(< \beta)$ -unstable above $\text{LS}(\mathbf{K}) + |\beta|$. \square

Remark 3.5. • If we have a specific K in mind, we may replace $(2^{\text{LS}(\mathbf{K})})^+$ by $\delta(\text{LS}(\mathbf{K}), \kappa)$ where $\kappa := |\Gamma_1| \leq 2^{\text{LS}(\mathbf{K})}$ in Shelah's Presentation Theorem.

- We used Galois types over the empty set in proving (2). The same proof goes through if we require the domain of the types to be some fixed (nonempty) model.

4 LOWER BOUND FOR STABILITY AND NO ORDER PROPERTY*

From Proposition 3.4, we saw that a stable AEC cannot have the order property* of length μ for some $\mu < \beth_{(2^{\text{LS}(\mathbf{K})})^+}$. Our goal is to show that this is also a lower bound for no order property* as well as stability.

Proposition 4.1. *Let λ be an infinite cardinal and α be an ordinal with $\lambda \leq \alpha < (2^\lambda)^+$. Then there is a stable AEC \mathbf{K} such that $\text{LS}(\mathbf{K}) = \lambda$, \mathbf{K} has the order property* of length up to $\beth_\alpha(\lambda)$ and is unstable anywhere below $\beth_\alpha(\lambda)$. Moreover, \mathbf{K} has JEP, NMM and $(< \aleph_0)$ -tameness but not AP.*

The proof of Proposition 4.1 will come after Corollary 4.4 and use Lemma 4.2 below.

Lemma 4.2. [She90, VII Theorem 5.5(6)] *Let λ be an infinite cardinal and α be an ordinal with $\lambda \leq \alpha < (2^\lambda)^+$. There is a stable AEC \mathbf{K}_1 such that $\text{LS}(\mathbf{K}_1) = \lambda$ and for all $M \in K_1$, M is well-ordered of order type at most α . Moreover, \mathbf{K}_1 has AP, JEP and is $(< \aleph_0)$ -tame.*

Proof. Let $L_1 := \{<, P_i : i < \lambda\}$ where P_i are unary predicates. T_1 requires $<$ to be a linear order. For $\beta < \alpha$, pick distinct subsets $S_\beta \subseteq \lambda$. Given an element x , we define its *type* by the set of indices i such that $P_i(x)$. We require that each element is characterized by its

type and the only possible types are among $\{S_\beta : \beta < \alpha\}$. If $\beta < \alpha$ and x, y have types S_β and S_α respectively, then we stipulate that $x < y$. These will be coded in the collection of types Γ_1 .

More precisely, for $\beta < \gamma < \alpha$, $S \subseteq \lambda$,

$$p_{\beta,\gamma}(x, y) := \{\neg(x < y)\} \cup \{P_i(x) : i \in S_\beta\} \cup \{\neg P_i(x) : i \notin S_\beta\}$$

$$\cup \{P_i(y) : i \in S_\gamma\} \cup \{\neg P_i(y) : i \notin S_\gamma\}$$

$$p_S(x) := \{P_i(x) : i \in S\} \cup \{\neg P_i(x) : i \notin S\}$$

$$\Gamma_1 := \{p_{\beta,\gamma}(x, y) : \beta < \gamma < \alpha\} \cup \{p_{S'} : S' \subseteq \lambda \text{ such that for } \beta < \alpha, S' \neq S_\beta\}$$

$$\cup \{x \neq y \wedge P_i(x) \leftrightarrow P_i(y) : i < \lambda\}$$

Let $\mathbf{K}_1 := EC(T_1, \Gamma_1)$ ordered by substructures. Notice that $|L(T_1)| = \lambda$ and $\Gamma_1 = 2^\lambda$. By replacing \mathbf{K}_1 by $(\mathbf{K}_1)_{\geq \lambda}$, we may assume $LS(\mathbf{K}_1) = \lambda$. Then $M_1 = \alpha$ is the maximal model (every model can be extended to an isomorphic copy of it) where for $\beta \in M_1$, $i < \lambda$, $P_i^{M_1}(\beta)$ iff $i \in S_\beta$. Hence \mathbf{K}_1 satisfies *AP*, *JEP* (but not *NMM*). As \mathbf{K}_1 has the maximal model of size $|\alpha| \leq 2^\lambda$, \mathbf{K}_1 is trivially stable in $\geq (2^\lambda)^+$. It is $(< \aleph_0)$ -tame because types are decided by the P_i 's they belong to. □

Remark 4.3. By [She90, VII Theorem 5.5(2)], for any λ and $\kappa \leq 2^\lambda$, $\delta(\lambda, \kappa) \leq (2^\lambda)^+$, so the threshold $(2^\lambda)^+$ cannot be improved. If we restrict $1 \leq \kappa < 2^\lambda$, then for $\alpha < \lambda^+$, we can still define \mathbf{K}_1 to be well-orderings of type at most α , where models are ordered by initial segments. Then we get a lower bound $\delta(\lambda, \kappa) \geq \lambda^+$. But the above proof does not go through because it requires $|\Gamma| = 2^\lambda$.

Using Lemma 4.2, we are able to answer Boney's conjecture in Remark 2.6. We will complete the proof of Proposition 4.1 after the proof of Corollary 4.4.

Corollary 4.4. *Under GCH, the λ -JEP assumption in Lemma 2.4 and Lemma 2.5 is not necessary. If $2^{\aleph_0} = \aleph_{\omega_1}$ and $\lambda = \aleph_0$, then λ -JEP is necessary in Lemma 2.4. If $2^{\aleph_0} = \aleph_1 =: \lambda$ and $2^{\aleph_1} = \aleph_{\omega_2}$, then λ -JEP is necessary in Lemma 2.5.*

Proof. By Lemma 2.3, we may assume $\kappa < \lambda$. Suppose \mathbf{K} is κ -stable in λ , then Lemma 2.4 and Lemma 2.5 are always true: for all $\|M\| = \lambda$, $\text{gS}_\lambda^\kappa = \lambda = |\text{gS}^\kappa(M)|$. Also, by taking sequences of length κ from $|M|$ (which give the algebraic types), we have $\text{gS}_\lambda^\kappa \geq |\text{gS}_\lambda^\kappa(M)| \geq \lambda^\kappa \geq (\text{gS}_\lambda^1)^\kappa$ where the last inequality is by stability.

Therefore, we may further assume κ -instability in λ , witnessed by M . Suppose GCH holds, then $|\text{gS}^\kappa(M)| \geq \lambda^+ = 2^\lambda = 2^{\lambda+\kappa} \geq \text{gS}_\lambda^\kappa \geq |\text{gS}^\kappa(M)|$. Also, $\text{gS}_\lambda^\kappa \geq |\text{gS}^\kappa(M)| \geq \lambda^+ = 2^\lambda = (2^\lambda)^\kappa \geq (\text{gS}_\lambda^1)^\kappa$. Hence it is consistent that the lemmas are always true (regardless of λ -JEP or $\text{cf}(\text{gS}_\lambda^\kappa)$).

We show that Lemma 2.4 can be false: let $\lambda = \aleph_0$, $\kappa = 1$ and suppose $2^{\aleph_0} = \aleph_{\omega_1}$ (which is consistent by Easton Theorem, see [Jec03, Theorem 15.18]). Define L_1, T_1 as in Lemma 4.2, pick \aleph_ω many distinct subsets of \aleph_0 , say $\{S_\beta : \beta < \aleph_\omega\}$. We allow $\beta < \gamma < \aleph_\omega$ when constructing $p_{\beta,\gamma}$. Now for $n < \omega$, define Γ_1^n to be the same as Γ_1 except that α is replaced by \aleph_n . Define $\mathbf{K}^n := EC(T_1, \Gamma_1^n)$ and \mathbf{K} be the disjoint union of all \mathbf{K}^n (adding \aleph_0 -many predicates and stipulate that no two elements belong to different predicates — this destroys \aleph_0 -JEP). Notice that \mathbf{K} is still a *EC* class where the language has size \aleph_0 and whose models omit 2^{\aleph_0} -many types. For any $M \in K$, $M \in K^n$ for some $n < \omega$. Since the unique maximal model in \mathbf{K}^n has size \aleph_n , $|\text{gS}(M)| \leq \aleph_n < \aleph_\omega = \sup_{n < \omega} \aleph_n = \text{gS}_{\aleph_0}^1$ where the last equality is due to the fact that distinct two elements in a model satisfy distinct subsets of $\{P_i : i < \aleph_0\}$. Hence $\text{gS}_{\aleph_0}^1$ (and similarly all $\text{gS}_{\aleph_n}^1$, $n < \omega$) is not attained by any model. $\text{cf}(\text{gS}_{\aleph_0}^1) = \aleph_0$ which satisfies the hypothesis.

We show that Lemma 2.5 can also be false: let $\lambda = \aleph_1$, $\kappa = \aleph_0$ and suppose $2^{\aleph_0} = \aleph_1$ and $2^{\aleph_1} = \aleph_{\omega_2}$ (which is consistent by Easton Theorem). This time we pick \aleph_ω many distinct subsets from \aleph_1 rather than from \aleph_0 . Form \mathbf{K} as above which is an *EC* class whose language has size \aleph_1 and whose models omit 2^{\aleph_1} -many types. Now $(\text{gS}_\lambda^1)^\kappa = (\text{gS}_{\aleph_1}^1)^{\aleph_0} = \aleph_\omega^{\aleph_0} > \aleph_\omega$. On the other hand, $\text{gS}_\lambda^\kappa = \text{gS}_{\aleph_1}^{\aleph_0} \leq \sup_{n < \omega} \sup\{|\text{gS}^{\aleph_0}(M)| : M \in (K^n)_{\aleph_1}\} \leq \sup_{n < \omega} \aleph_n^{\aleph_0} = \sup_{n < \omega} (2^{\aleph_0} \cdot \aleph_n) = \sup_{n < \omega} \aleph_1 \cdot \aleph_n = \aleph_\omega < (\text{gS}_\lambda^1)^\kappa$ where the second equality is a special case of the Hausdorff formula [Jec03, Equation 5.22]. \square

Proof of Proposition 4.1. Fix α as in Lemma 4.2, we use \mathbf{K}_1 to build \mathbf{K} as follows: let $L := L_1 \cup \{E, Q, R, f, g, c\}$ where E, Q are binary predicates, R is a ternary predicate, f

is a unary function, g is a symmetric binary function and c is a constant. Thus we have $|L| = \lambda$. A model in $\mathbf{K} = (\mathbf{K}_0, \mathbf{K}_1, \mathbf{K}_2)$ has three sorts (M_0, M_1, M_2) . M_1 is in K_1 , M_2 will take care of NMM while M_0 is the iterated power sets of M_1 . In details, we require:

1. $M_1 \in EC(T_1, \Gamma_1)$ as in Lemma 4.2. We identify it as an ordinal $\leq \alpha$.
2. M_2 is an infinite model of the theory of pure equality.
3. If xEy , then x is in $M_0 \cup M_1$ while y is in M_0 . E also satisfies the extensionality axiom.
4. The first argument of Q is in M_1 . We write $Q_i(\cdot) := Q(i, \cdot)$ and abbreviate Q_i the elements x in M_0 with $Q(i, x)$. For limit ordinal σ in M_1 , we require $Q_\sigma = \bigcup_{i < \sigma} Q_i$.
5. f is the rank function from M_0 to M_1 such that each x is sent to the smallest i with $Q_i(x)$. If $x \in y$, then $f(x) < f(y)$.
6. R , g and c code the total order of M_0 : we define $R(\beta, x, y)$ and $g(x, y)$ as follows:
 - (a) β in M_1 , x, y in M_0
 - (b) If $f(x) \neq f(y)$, then we say x is less than y when x has a smaller rank than y .
 - (c) If $f(x) = f(y) = 0$, then $g(x, y)$ is the $<$ -least element in the symmetric difference of x, y . $R(0, x, y)$ if $g(x, y)Ey$, in which case we say x is less than y .
 - (d) If $f(x) = f(y) = \beta > 0$, then $g(x, y)$ is the least element in the symmetric difference of x, y . $R(\beta, x, y)$ if $g(x, y)Ey$, in which case we say x is less than y .
 - (e) c is the default value for $g(x, y)$ when x, y are in M_1 or M_2 , or when x, y in M_0 are equal or have different ranks.

(If we think of subsets as sequences, we are ordering M_0 by rank, and then by lexicographical order of each rank.)

We order $(M_0, M_1, M_2) \leq_{\mathbf{K}} (N_0, N_1, N_2)$ iff for $i \leq 2$, $M_i, N_i \in K_i$ and $M_i \subseteq N_i$. Notice that we can describe \mathbf{K} as some $EC(T, \Gamma)$ where T is a $\forall\exists$ theory, $|L(T)| = \lambda$ and $|\Gamma| = 2^\lambda$. Also, $LS(\mathbf{K}) = \lambda$ because $LS(\mathbf{K}_1) = \lambda$ and we can close any set in M_0 to a model by adding witnesses for f, g in (5),(6) ω -many times. Therefore \mathbf{K} is an AEC.

The maximal model in $(\mathbf{K}_0, \mathbf{K}_1)$ is $M^* := (V_\alpha(\alpha), \alpha)$, where for $\beta < \alpha$, $Q_\beta^{M^*} := V_{1+\beta}(\alpha)$. M^* witnesses that \mathbf{K} have *JEP*. \mathbf{K} is $(< \aleph_0)$ -tame because elements are determined either by the predicates P_i ; or their ranks and their own elements. With \mathbf{K}_2 , we know that \mathbf{K} has *NMM*. Since M^* has size $\beth_\alpha(\lambda)$, $(\mathbf{K}_0, \mathbf{K}_1)$ is trivially stable in $\geq \beth_\alpha(\lambda)$. As \mathbf{K}_2 is trivially stable everywhere, \mathbf{K} is stable in $\geq \beth_\alpha(\lambda)$. We now show instability in $< \beth_\alpha(\lambda)$ and the order property of lengths up to $\beth_\alpha(\lambda)$.

For instability, $(\mathcal{P}(\lambda), \lambda, \omega)$ witnesses that \mathbf{K} is unstable between $[\lambda, 2^\lambda)$. Consider the maximal model M^* above, for each $\beta < \alpha$, $|Q_\beta^{M^*}| = \beth_{1+\beta}(\lambda)$. By item (6) above, $Q_\beta^{M^*}$ can distinguish all elements in $Q_{\beta+1}^{M^*}$, so $(\mathbf{K}_0, \mathbf{K}_1)$, and hence \mathbf{K} is unstable in $[\beth_{1+\beta}(\lambda), \beth_{1+\beta+1}(\lambda))$. Therefore, we have instability in $[\lambda, \beth_\alpha(\lambda))$.

For the order property*, we apply the order in (6) to $V_\alpha(\alpha)$ of the maximal model. In details: let a less than b while c less than d . If a, b is mapped to d, c respectively, then $f(d) = f(a) \leq f(b) = f(c) \leq f(d)$. It cannot happen to rank 0 because their elements are well-ordered in M_1 . Since a is less than b , $g(a, b) \in b$. Then by mapping $g(d, c) (= g(c, d)) \in c$ which shows that d is less than c , contradiction. As $|V_\alpha(\alpha)| = \beth_\alpha(\lambda)$, we have the order property* of length $\beth_\alpha(\lambda)$.

$(\mathbf{K}_0, \mathbf{K}_1)$ does not have *AP*: Pick an element x from $(V_\alpha(\alpha), \alpha)$ which contains $\geq \lambda^+$ elements. Close x to a substructure N of size λ , then there is yEx in $(V_\alpha(\alpha), \alpha)$ but y is not in N . N can be included in $(V_\alpha(\alpha), \alpha)$ such that x is mapped to $(x - \{y\})$. Suppose the following amalgam exists:

$$\begin{array}{ccc}
 x \in (V_\alpha(\alpha), \alpha) & \cdots \cdots \cdots \rightarrow & x \in W \\
 \uparrow & & \uparrow t \\
 x \in N & \longrightarrow & x - \{y\} \in (V_\alpha(\alpha), \alpha)
 \end{array}$$

Without loss of generality, we may assume the top dotted arrow is identity (hence we can write the image of x to be x itself). Since $(V_\alpha(\alpha), \alpha)$ is maximal, $W = (V_\alpha(\alpha), \alpha)$. Therefore, $t \in \text{Aut}((V_\alpha(\alpha), \alpha))$. By an induction argument, t must be the identity (which boils down to the fact that α , the maximal model in \mathbf{K}_1 , is rigid). From the right dotted arrow, $x - \{y\}$ would be mapped to x , which is a contradiction. \square

Remark 4.5. 1. One way to save *AP* is to redefine $\leq_{\mathbf{K}}$ by the *E*-transitive closure, but

it raises $\text{LS}(\mathbf{K})$ to $V_{\alpha-1}(\lambda)$. In this case, instability and the order property* length are up to $2^{\text{LS}(\mathbf{K})}$.

2. Our total order is ill-founded: $\alpha, \alpha - \{0\}, \alpha - \{0, 1\}, \dots$ form an infinite descending sequence in $Q_0^{M^*}$. It is not clear how to extract a witness to the order property of length $> 2^\lambda$. We will see in Corollary 6.8 that we can refine our example to have the order property at least up to $\beth_{\alpha-3}(\lambda)$.

Corollary 4.6. *1. For stable AECs, the Hanf number for the order property* length is exactly $\beth_{(2^{\text{LS}(\mathbf{K})})^+}$.*

2. *The Hanf number for stability is at least $\beth_{(2^{\text{LS}(\mathbf{K})})^+}$. In other words, let $\lambda \geq \aleph_0$ and $\mu < \beth_{(2^\lambda)^+}$, there is a stable AEC \mathbf{K} such that $\text{LS}(\mathbf{K}) = \lambda$ and the first stability cardinal is greater than μ .*

Proof. Combine Proposition 4.1 and Proposition 3.4 and range α in $[\lambda, (2^\lambda)^+]$. □

In the next two sections, we develop the machinery to show that the lower bound in (2) is tight, based on the arguments in [Vas16b]. Then we conclude: our example witnesses that the bound for the order property is also tight.

5 GALOIS MORLEYIZATION AND SYNTACTIC ORDER PROPERTY

Galois Morleyization is a way to capture tameness syntactically by adding infinitary predicates. First recall the definition of tameness:

Definition 5.1. Let κ be an infinite cardinal.

1. Let $p = \text{gtp}(a/A, N)$ where $a = \langle a_i : i < \alpha \rangle$ may be infinite, $I \subseteq \alpha$, $A_0 \subseteq A$. We write $l(p) := l(a)$, $p \upharpoonright A_0 := \text{gtp}(a/A_0, N)$, $a^I = \langle a_i : i \in I \rangle$ and $p^I := \text{gtp}(a^I/A, N)$.
2. K is $(< \kappa)$ -tame for $(< \alpha)$ -types if for any subset A in some model of K , any $p \neq q \in \text{gS}^{< \alpha}(A)$, there is $A_0 \subseteq A$, $|A_0| < \kappa$ with $p \upharpoonright A_0 \neq q \upharpoonright A_0$. We omit $(< \alpha)$ if $\alpha = 2$.
3. K is $(< \kappa)$ -short if for any $\alpha \geq 2$, any subset A in some model of K , $p \neq q \in \text{gS}^{< \alpha}(A)$, there is $I \subseteq \alpha$, $|I| < \kappa$ with $p^I \neq q^I$.

4. κ -tame means $(< \kappa^+)$ -tame. Similarly for shortness.

Now we construct Galois Morleyization:

Definition 5.2. [Vas16b, Definitions 3.3, 3.13] Let κ be an infinite cardinal and K be an AEC in a (finitary) language L . The $(< \kappa)$ -Galois Morleyization of K is a class \hat{K} of structures in a language \hat{L} such that:

1. \hat{L} is a $(< \kappa)$ -ary language. For convenience we may require $L \subseteq \hat{L}$.
2. For each $p \in \text{gS}^{<\kappa}(\emptyset)$, we add a predicate R_p of length $l(p)$ to \hat{L} .
3. For each $M \in K$, we define $\hat{M} \in \hat{K}$ with $|M| = |\hat{M}|$. For $p \in \text{gS}^{<\kappa}(M)$, $a \in |\hat{M}|^{l(p)}$, $\hat{M} \models R_p[a]$ iff $a \models p$ in K . Extend the definition to quantifier-free formulas of $\hat{L}_{\kappa, \kappa}$.
4. The $(< \kappa)$ -syntactic type of $a \in |\hat{M}|^{<\kappa}$ over $A \subseteq |\hat{M}|$ is defined by $\text{tp}_{\text{qf-}\hat{L}_{\kappa, \kappa}}(a/A; \hat{M})$, namely the quantifier-free formulas of $\hat{L}_{\kappa, \kappa}$ over A that a satisfies. We will abbreviate it as $\text{tp}_{\kappa}(a/A; \hat{M})$.
5. For $\hat{M}, \hat{N} \in \hat{K}$, we order $\hat{M} \leq_{\hat{\mathbf{K}}} \hat{N}$ iff $M \leq_{\mathbf{K}} N$. We will omit the subscripts.

Remark 5.3. 1. If we allow AECs to have infinitary languages, we can view \hat{K} as an AEC.

2. The above is well-defined even for AECs without AP , but readers can assume the existence of a monster model \mathfrak{C} for convenience.

3. $|\hat{L}| = |L| + |\text{gS}^{<\kappa}(\emptyset)| \leq 2^{<(\kappa + \text{LS}(\mathbf{K})^+)}$.

The following justifies the definition of Galois Morleyization in tame AECs:

Proposition 5.4. [Vas16b, Corollary 3.18(2)] *Let K be an AEC and \hat{K} be its $(< \kappa)$ -Galois Morleyization. For each $p = \text{gtp}(b/A; M) \in \text{gS}(A)$, define $p_{\kappa} := \text{tp}_{\kappa}(b/A; \hat{M})$ to be its $(< \kappa)$ -syntactic version. Then K is $(< \kappa)$ -tame iff $p \mapsto p_{\kappa}$ is a 1-1 correspondence.*

Proof. \Rightarrow : The map is well-defined because Galois types are finer than syntactic types. It is a surjection by construction. Suppose $p \neq q \in \text{gS}(A)$, by $(< \kappa)$ -tameness we may assume

the domain A has size $< \kappa$. Let $b \models p$ and $b' \models q$. Then bA and $b'A$ satisfy different types in $\text{gS}^{<\kappa}(\emptyset)$, say r and s respectively. Thus $bA \models R_r \wedge \neg R_s$ while $b'A \models R_s \wedge \neg R_r$.

\Leftarrow : Suppose $p = \text{gtp}(b/A; M) \neq q = \text{gtp}(b'/A; M')$. Then $p_\kappa \neq q_\kappa$ and we can find $r \in \text{gS}^{<\kappa}(\emptyset)$ and (a suitable enumeration of) $A_0 \subseteq A$ such that $M \models R_r[b; A_0]$ but $M' \models \neg R_r[b'; A_0]$. This means $bA_0 \models r$ while $b'A_0 \not\models r$. Hence $\text{gtp}(b/A_0; M) \neq \text{gtp}(b'/A_0; M')$ witnessing $(< \kappa)$ -tameness. \square

Remark 5.5. There is a stronger version assuming $(< \kappa)$ -shortness in [Vas16b, Corollary 3.18(1)] but we have no use of it here.

We now define an infinitary version of the syntactic order property:

Definition 5.6. Let $2 \leq \alpha \leq \kappa$ and $1 \leq \beta < \kappa$ be ordinals.

1. [Vas16b, Definition 4.2] In Definition 3.2, we replace all occurrences of “order property” by “syntactic order property” while requiring the condition in (1) there be: there exist some $\langle a_i : i < \mu \rangle \subseteq \hat{M} \in \hat{K}$ and some quantifier-free $\hat{L}_{\kappa, \kappa}$ formula $\phi(x, y)$ such that $l(a_i) = \beta$, and for $i, j < \mu$, $i < j$ iff $\hat{M} \models \phi[a_i, a_j]$.
2. As in Definition 3.3, we define syntactic order property* if in (1) we allow the index set to be a linear order I with $|I| = \mu$, instead of being a well-ordering μ .

The following links the (Galois) order property in K with the syntactic order property in \hat{K} .

Proposition 5.7. *Let κ be an infinite cardinal and \hat{K} be $(< \kappa)$ -Galois Morleyization of K . Let $1 \leq \beta < \kappa$ be an ordinal and $M \in K$. Let λ, μ be infinite cardinals and $\chi := |\text{gS}^{\beta+\beta}(\emptyset)|$.*

1. [Vas16b, Proposition 4.4]
 - (a) *If \hat{M} has the syntactic β -order property of length μ , then M has the β -order property of length μ .*
 - (b) *If M has the β -order property of length μ for some $\mu \geq (2^{\lambda+\chi})^+$, then \hat{M} has the syntactic β -order property of length λ^+ .*

2. (a) If \hat{M} has the syntactic β -order property* of length μ , then M has the β -order property* of length μ .
- (b) If M has the β -order property* of length μ for some $\mu \geq (2^{\lambda+x})^+$, then \hat{M} has the syntactic β -order property* of length λ^+ .
3. The following are equivalent:
 - (a) K has the β -order property.
 - (b) K has the β -order property*.
 - (c) \hat{K} has the syntactic β -order property.
 - (d) \hat{K} has the syntactic β -order property*.

Proof. 1. (a) is true because Galois types are finer than syntactic types. For (b): suppose $\langle a_i : i < \mu \rangle$ witnesses the β -order property. By Erdős-Rado Theorem, we have $\mu \rightarrow (\lambda^+)_{2^x}^2$. Apply this to $(i < j) \mapsto \text{gtp}(a_i a_j / \emptyset; M)$ and then on $(j < i) \mapsto \text{gtp}(a_j a_i / \emptyset; M)$. We can find $\langle b_i : i < \lambda^+ \rangle$ subsequence of $\langle a_i : i < \mu \rangle$, $p \neq q \in \text{gS}^{\beta+\beta}(\emptyset)$ such that for $i < j < \lambda$, $\text{gtp}(b_i b_j / \emptyset; M) = p$ and $\text{gtp}(b_j b_i / \emptyset; M) = q$. We may choose R_p (or R_q) to witness the β -syntactic order property.

2. The same proof goes through, because Erdős-Rado Theorem applies to linear orders too.

3. (1) gives (a) \Leftrightarrow (c) while (2) gives (b) \Leftrightarrow (d). (a) \Leftrightarrow (b) is by Proposition 3.4(1),(2). □

Definition 5.8. Let κ be an infinite cardinal, $2 \leq \alpha \leq \kappa$ and $1 \leq \beta < \kappa$ be ordinals, \hat{K} be the $(< \kappa)$ -Galois Morleyization of K . Then

1. For $\mu \geq \text{LS}(\mathbf{K}) + |\beta|$, \hat{K} is β -syntactically stable in μ if

$$|\{p_\kappa : p \in \text{gS}^\beta(A; M), A \subseteq |M|, |A| \leq \mu, M \in K\}| \leq \mu.$$

2. For $\mu \geq \text{LS}(\mathbf{K}) + |\alpha|$, \hat{K} is $(< \alpha)$ -syntactically stable in μ if

$$|\{p_\kappa : p \in \text{gS}^{<\alpha}(A; M), A \subseteq |M|, |A| \leq \mu, M \in K\}| \leq \mu.$$

Corollary 5.9. *Let $\beta \geq 1$ be an ordinal and $\mu \geq \text{LS}(\mathbf{K}) + |\beta|$ be a cardinal.*

1. [Vas16b, Fact 4.9] *If K has the β -order property, then K is not β -stable in μ . If also $\beta < \kappa$, then \hat{K} is not β -(syntactically) stable in μ .*
2. *The same conclusion holds when K has the β -order property*.*

Proof. By Proposition 5.7, either assumption gives the syntactic β -order property. This implies β -syntactic instability in μ , using the proof of Proposition 3.4 (in particular replace $F^{\mathfrak{B}}(\alpha)$ by a witness of the syntactic β -order property of length \beth_α). \square

6 SHELAH'S STABILITY THEOREM

We will connect syntactic stability with no syntactic order property. The original result due to Shelah was in a more general context but only proof sketches were given. Vasey [Vas16b, Fact 4.10] applied it to AECs without a complete proof so we write out all the details. We will also remove the requirement that the order property length be a successor (which was hinted in [She09b, Exercise 1.22]).

Theorem 6.1. [She09b, V.A. Theorem 1.19] *Let $\chi \geq 2^{<(\kappa + \text{LS}(\mathbf{K})^+)}$, μ be an infinite cardinal such that $\mu = \mu^\chi + 2^{2^{<\chi}}$. Suppose \hat{K} does not have the $(< \kappa)$ -syntactic order property of length χ , then \hat{K} is $(< \kappa)$ -syntactically stable in μ .*

The proof will be given after Lemma 6.5. Before that we state some relevant definitions and lemmas.

Definition 6.2. Let κ be an infinite cardinal, Π be a set of quantifier-free formulas of $\hat{L}_{\kappa, \kappa}$ over A , and p_κ be a $(< \kappa)$ -syntactic type over A . We say p_κ *splits over* Π if there are $\phi(x; b), \neg\phi(x; c) \in p_\kappa$ such that for any \hat{M} containing b, c and the parameters from Π , any $\psi(y; d) \in \Pi$ with $l(y) = l(b) = l(c)$, we have $\hat{M} \models \psi[b; d] \Leftrightarrow \hat{M} \models \psi[c; d]$ (the choice of \hat{M} does not matter because its interpretation of R_p is external).

If we require the witnesses $\phi(x; b), \neg\phi(x; c)$ to be from a fixed formula $\phi(x; y)$, then we say p_κ ϕ -*splits over* Π .

Lemma 6.3. [She09b, V.A. Fact 1.10(4)] *Using the above notation,*

$$|\{p_\kappa \upharpoonright \phi : p_\kappa \text{ does not } \phi\text{-split over } \Pi\}| \leq 2^{2^{|\Pi|}}$$

$$|\{p_\kappa : p_\kappa \text{ does not split over } \Pi\}| \leq 2^{2^{|\Pi|} \cdot \chi}$$

where $\chi := |\hat{L}|^{<\kappa} = (|\text{gS}^{<\kappa}(\emptyset)|)^{<\kappa} \leq 2^{<(\kappa + \text{LS}(\mathbf{K})^+)}$ is the size of the set of quantifier-free formulas of $\hat{L}_{\kappa, \kappa}$.

Proof. We count the number of combinations to build a $p_\kappa \upharpoonright \phi$ that does not ϕ -split over Π . Partition the parameters of ϕ by their Π -type. Namely, b, c are equivalent iff for any \hat{M} containing b, c and the parameters from Π , any $\psi(y; d) \in \Pi$ with $l(y) = l(b) = l(c)$, we have $\hat{M} \models \psi[b; d] \Leftrightarrow \hat{M} \models \psi[c; d]$. Then there are $2^{|\Pi|}$ -many classes. Within each class, say containing b , it remains to choose whether $\phi(x; b)$ or $\neg\phi(x; b)$ is in $p_\kappa \upharpoonright \phi$. Hence we have $2^{2^{|\Pi|}}$ -many choices.

The second part follows from the observation that a $(< \kappa)$ -syntactic type p_κ is determined by its restrictions $p_\kappa \upharpoonright \phi$ where ϕ is a quantifier-free formula of $\hat{L}_{\kappa, \kappa}$. \square

Definition 6.4. A set A is $(< \chi)$ -compact if for any \hat{M} containing A , any cardinal $\lambda < \chi$, any quantifier-free formulas $\{\phi_i(x) : i < \lambda\}$ in $\hat{L}_{\kappa, \kappa}$ with parameters from A , if $\hat{M} \models \bigwedge_{i < \lambda} \phi_i[b]$ for some $b \in \hat{M}$, then $\hat{M} \models \bigwedge_{i < \lambda} \phi_i[a]$ for some $a \in A$.

Lemma 6.5. [She09b, V.A. Theorem 1.12] *Let χ be an infinite cardinal, A be $(< \chi)$ -compact with $A \subseteq |\hat{M}|$, $\phi(x; y)$ be a quantifier-free formula in $\hat{L}_{\kappa, \kappa}$. Either*

1. *For any $m \in |\hat{M}|$, there is a set $\Pi \subseteq \{\phi(x; a) : a \in [A]^{<\kappa}\}$ such that $|\Pi| < \chi$ and $\text{tp}_\kappa(m/A; \hat{M}) \upharpoonright \phi$ does not ϕ -split over Π ; or*
2. *$A \subseteq |\hat{M}|$ witnesses the $(< \kappa)$ -syntactic order property of length χ .*

Proof. Suppose (1) does not hold, then we can pick $m \in |\hat{M}|$ such that $\text{tp}_\kappa(m/A; \hat{M}) \upharpoonright \phi$ splits over any Π with $|\Pi| < \chi$. Thus we can recursively build

1. $\langle m_i, b_i, c_i : i < \chi \rangle$ inside A .
2. For $j < \chi$, $\hat{M} \models \phi[m; b_j] \Leftrightarrow \neg\phi[m; c_j]$

3. For $i < j < \chi$, $\hat{M} \models \phi[m_i; b_j] \leftrightarrow \phi[m_i; c_j]$.
4. For $j < \chi$, $m_j \models \bigwedge_{i \leq j} (\phi(x; b_i) \leftrightarrow \neg \phi(x; c_i))$.

The construction is possible by the definition of ϕ -splitting and by $(< \chi)$ -compactness of A . The sequence $\langle m_i b_i c_i : i < \chi \rangle$ witnesses the $(< \kappa)$ -syntactic order property of length χ via the formula $\phi(x; y) \leftrightarrow \phi(x; z)$. \square

Proof of Theorem 6.1. Let A be a set of size μ . As in Lemma 6.3, χ bounds the number of quantifier-free formulas of $\hat{L}_{\kappa, \kappa}$. Also $\mu = \mu^{< \chi}$, so we may assume A is $(< \chi)$ -compact (see Definition 6.4). Since Lemma 6.5(2) fails, (1) must hold for each quantifier-free formula ϕ of $\hat{L}_{\kappa, \kappa}$.

Now we count the number of $(< \kappa)$ -syntactic types. Each type p_κ is determined by its restrictions $\{p_\kappa \upharpoonright \phi : \phi \text{ is a quantifier-free formula in } \hat{L}_{\kappa, \kappa}\}$. Since $\mu = \mu^\chi$, we may assume $p_\kappa = p_\kappa \upharpoonright \phi$ for a fixed ϕ (this is where we need $\mu = \mu^\chi$ instead of $\mu = \mu^{< \chi}$). By Lemma 6.5(1), we can find some Π_{p_κ} of size $< \chi$ such that p_κ does not ϕ -split over Π_{p_κ} . There are $|A|^{< \chi} = \mu^{< \chi}$ -many ways to choose Π_{p_κ} . For each fixed $\Pi = \Pi_{p_\kappa}$, Lemma 6.3 gives at most $2^{2^{|\Pi|}} = 2^{2^{< \chi}}$ -many choices for p_κ . So in total there are $\mu^{< \chi} + 2^{2^{< \chi}} = \mu$ -many choices. \square

Corollary 6.6. *Let K be a stable AEC.*

1. [Vas16b, Theorem 4.13] *If K is $(< \kappa)$ -tame, has AP and is stable in some cardinal $\geq \kappa^-$, then the first stability cardinal is bounded above by $\beth_{(2^{< (\kappa + \text{LS}(\mathbf{K}^+))})^+}$.*
2. *If K is $(< \kappa)$ -tame and does not have $(< \kappa)$ -order property of length $\chi := 2^{< (\kappa + \text{LS}(\mathbf{K}^+))}$, then the first stability cardinal is bounded above by $2^{2^{< \chi}}$.*
3. *If K is $\text{LS}(\mathbf{K})$ -tame and does not have $\text{LS}(\mathbf{K})$ -order property of length $2^{\text{LS}(\mathbf{K})}$, then the first stability cardinal is bounded above by $\beth_3(\text{LS}(\mathbf{K}))$.*
4. *Let $|D(T)| := |\text{gS}^{< \omega}(\emptyset)|$. If K is $(< \aleph_0)$ -tame and does not have $(< \omega)$ -order property of length $|D(T)|$, then the first stability cardinal is bounded above by $\beth_2(|D(T)|)$.*

Proof. We prove (1): Since \mathbf{K} is $(< \kappa)$ -tame, by Proposition 5.4 $(< \kappa)$ -syntactic stability in $\hat{\mathbf{K}}$ is equivalent to $(< \kappa)$ -stability in \mathbf{K} . Also, by the contrapositive of Proposition 5.7(1)(a),

no $(< \kappa)$ -order property of length χ in \mathbf{K} implies no $(< \kappa)$ -syntactic order property in $\hat{\mathbf{K}}$ of length χ .

Let \mathbf{K} be stable in some $\lambda \geq \kappa^-$. Since it is $(< \kappa)$ -tame, Theorem 2.10 gives stability in $\beth_{\lambda^+}(\lambda)$ and Theorem 2.2 gives $(< \kappa)$ -stability in $\beth_{\lambda^+}(\lambda)$. By Proposition 3.4(3), there is $\chi < \beth_{(2^{<(\kappa+\text{LS}(\mathbf{K})^+)})^+}$ such that \mathbf{K} does not have $(< \kappa)$ -order property of length χ . We may assume $\chi \geq 2^{<(\kappa+\text{LS}(\mathbf{K})^+)}$. By the first paragraph and Theorem 6.1, \mathbf{K} is $(< \kappa)$ -stable in $2^{2^{<\chi}} < \beth_{(2^{<(\kappa+\text{LS}(\mathbf{K})^+)})^+}$. \square

Remark 6.7. In particular (4) misses the actual lower bound by \beth_2 [She90, III Theorem 5.15].

We now show the promised result in Remark 4.5:

Corollary 6.8. *Let λ be an infinite cardinal and γ be an ordinal with $\lambda \leq \gamma < (2^\lambda)^+$. Then there is a stable AEC \mathbf{K} such that $\text{LS}(\mathbf{K}) = \lambda$, \mathbf{K} has the order property of length up to $\beth_\gamma(\lambda)$. Moreover, \mathbf{K} has JEP, NMM, $(< \aleph_0)$ -tameness but not AP.*

Proof. Let $\chi := \beth_\gamma(\lambda)$. We use the example in Proposition 4.1 with $\alpha = \gamma + 3$. Suppose \mathbf{K} has no $(< \omega)$ -order property of length χ . Since \mathbf{K} is $(< \aleph_0)$ -tame, by Proposition 5.4 $(< \omega)$ -syntactic stability in $\hat{\mathbf{K}}$ is equivalent to $(< \omega)$ -stability in \mathbf{K} . Also, by the contrapositive of Proposition 5.7(1)(a), no $(< \omega)$ -order property of length χ in \mathbf{K} implies no $(< \omega)$ -syntactic order property in $\hat{\mathbf{K}}$ of length χ . Since $\chi \geq 2^{<\aleph_0}$, by Theorem 6.1, \mathbf{K} is $(< \omega)$ -stable in all $\mu = \mu^\chi + 2^{2^{<\chi}}$. In particular, it is $(< \omega)$ -stable in $\beth_2(\chi) = \beth_{\alpha-1}(\lambda) = \beth_{\alpha-1}(\lambda)$, contradicting the fact that $M^* \in \mathbf{K}$ is unstable in any cardinal $< \beth_\alpha(\lambda)$. \square

Remark 6.9. In our example, where exactly is the witness to the $(< \omega)$ -order property of $\beth_\gamma(\lambda)$? Tracing the proofs, the key is the recursive construction in Lemma 6.5, where a long splitting chain is utilized. Fix a cardinal $\chi \leq \beth_{\alpha-1}(\lambda)$. For our \mathbf{K} , we do not even need the b_i and can simply set $\langle c_i, m_i : i < \chi \rangle$ such that all elements are distinct and m_i contains exactly $\{c_j : j \leq i\}$. Then $\langle c_i, m_i : i < \chi \rangle$ witnesses the 2-order property of length χ , via the formula $\phi(x_1 y_1; x_2 y_2) := x_1 E y_2$. Therefore, we have an explicit example of the order property up to length $\beth_{\alpha-1}(\lambda)$ (the subscript cannot go further because most elements on the top rank do not belong to any other elements).

We can conclude:

- Corollary 6.10.** 1. *For stable AECs, the Hanf number for the order property length is exactly $\beth_{(2^{\text{LS}(\mathbf{K})})^+}$.*
2. *For stable AECs, the Hanf number for the order property* length is exactly $\beth_{(2^{\text{LS}(\mathbf{K})})^+}$.*
3. *The Hanf number for stability is at least $\beth_{(2^{\text{LS}(\mathbf{K})})^+}$. In other words, let $\lambda \geq \aleph_0$ and $\mu < \beth_{(2^\lambda)^+}$, there is a stable AEC \mathbf{K} such that $\text{LS}(\mathbf{K}) = \lambda$ and the first stability cardinal is greater than μ .*
4. *With $\text{LS}(\mathbf{K})$ -tameness and AP, the Hanf number for stability is at most $\beth_{(2^{\text{LS}(\mathbf{K})})^+}$. In other words, if \mathbf{K} is a stable AEC with AP and $\text{LS}(\mathbf{K})$ -tameness, then the first stability cardinal is at most $\beth_{(2^{\text{LS}(\mathbf{K})})^+}$.*

- Proof.* 1. Lower bound is by Corollary 6.8 and upper bound is by Proposition 3.4(3).
2. Lower bound is by Proposition 4.1 and upper bound is by Proposition 3.4(3).
3. By Proposition 4.1.
4. By Corollary 6.6(1).

□

We finish this section with the following question: are the bounds in (3) and (4) optimal?

7 SYNTACTIC SPLITTING

We will give a syntactic proof to Theorem 2.10 using Galois Morleyization. The advantage is that types are syntactic and can be over sets of size less than $\text{LS}(\mathbf{K})$; the disadvantage is that we have extra assumptions.

Assumption 7.1. We assume the existence of a monster model \mathfrak{C} ($AP + JEP + NMM$), where each set is inside some model in K and each (set) embedding/isomorphism is extended by a \mathbf{K} -embedding/isomorphism. We also assume AP over set bases: if

$A \subseteq |M_1| \cap |M_2|$ and M_1, M_2 interpret A in the same way, then there are $M_3 \geq M_2$ and $f : M_1 \xrightarrow[A]{} M_3$.


Definition 7.2. Let μ be an infinite cardinal, A, B be sets in some models of K .

1. B is *universal* over A if $A \subset B$ and for any $|B'| \leq |B|$, $B' \supseteq A$, there is $f : B' \xrightarrow[A]{} B$. We write $A \subset_u B$.
2. B is μ -*universal* over A if B' in (1) must have size $\leq \mu$.
3. B is μ -*homogeneous* if it is $< \mu$ -universal over any $C \subset B$ of size $< \mu$.
4. Let $A \subseteq B \in K$, $p \in \text{gS}(B)$. We say p μ -*splits over* A if there exists $A \subseteq B_1, B_2 \subseteq B$, $\|B_1\| = \mu$, $f : B_1 \cong_A B_2$ with $f(p) \upharpoonright B_2 \neq p \upharpoonright B_2$.

[She09a, II 1.16(1)(a)] shows that universal *models* of size μ exist if \mathbf{K} is stable in $\mu \geq \text{LS}(\mathbf{K})$ (and has μ -AP, μ -NMM). We prove universal sets exist:

- Proposition 7.3.**
1. For any A , if \mathbf{K} is stable in $|A|$, there is $|B| = |A|$, $B \supset_u A$.
 2. For any infinite λ, μ and any $|A| \leq \lambda$. If $\lambda^{<\mu} = \lambda$, then there is a μ -homogeneous $B \supset A$ of size λ .

Proof. 1. Let $\mu := |A|$. Build $\langle B_i : i < \mu \rangle$ increasing and continuous such that $B_0 := A$, $B_{i+1} \models \text{gS}(B_i)$. For any $A' \supseteq A$, $|A'| \leq \mu$. We may assume $|A' - A| = \mu$ and write $A' = A \cup \{a_i : i < \mu\}$. Define $A_i := A \cup \{a_j : j \leq i\}$ and $\langle f_i : A_i \xrightarrow[A]{} B_i : i \leq \mu \rangle$ increasing and continuous partial embeddings such that $f_i(a_i) \in B_i$. Set $f_{-1} := \text{id}_A$ and suppose f_i has been constructed, obtain \tilde{A}' , an isomorphic copy of A' over $\text{ran}(f_i)$ and denote \tilde{a}_{i+1} the copy of a_{i+1} . Now $B_{i+1} \models \text{gS}(B_i) \supseteq \text{gS}(\text{ran}(f_i))$ so it realizes the type of \tilde{a}_{i+1} over $\text{ran}(f_i)$, say by b_{i+1} . By AP there is $g : \tilde{a}_{i+1} \mapsto b_{i+1}$ fixing $\text{ran}(f_i)$. Define $f_{i+1}(a_{i+1}) := b_{i+1}$.



2. By an exhaustive argument, we can build a (set) saturated $B \supset A$. We check that it is μ -homogeneous. Let $C \subset B$, $C' \supseteq C$ both of size $< \mu$, the argument from the previous item applies because B is saturated and $|C'| < \mu$.

□

We notice a correspondence between μ -Galois splitting and $(< \mu^+)$ -syntactic splitting (see Definition 6.2); a similar treatment for coheir has already been done in [Vas16b, Section 5]). We write q μ -syn-splits over A to mean q $(< \mu^+)$ -syntactically splits over the quantifier-free formulas of \hat{L}_{μ^+, μ^+} over A .

Proposition 7.4. *Let μ be an infinite cardinal, $\hat{\mathbf{K}}$ be the $(< \mu^+)$ -Galois Morleyization of \mathbf{K} . For any $A \subseteq B \in K$, $p \in \text{gS}(B)$, p μ -splits over A iff p_{μ^+} μ -syn-splits over A .*

Proof. Suppose $d \models p$ μ -splits over A , obtain witness $f : B_1 \cong_A B_2$ as above. Enumerate $A = a$ and B_i as b_i . Since $f(p) \upharpoonright B_2 \neq p \upharpoonright B_2$, $r_1 := \text{gtp}(f(d)b_2/\emptyset) \neq \text{gtp}(db_2/\emptyset)$, so $\hat{\mathcal{C}} \models R_{r_1}[db_1] \wedge R_{r_1}[f(d)b_2] \wedge \neg R_{r_1}[db_2]$. b_1 and b_2 have the same syntactic type over a because of f . Therefore, $p_{\mu^+} = \text{tp}_{\mu^+}(d/B) \supseteq \text{tp}_{\mu^+}(d/B_1 \cup B_2)$ μ -syn-splits over A .

Conversely, suppose p_{μ^+} μ -syn-splits over A . There are $\phi(x; b_1), \neg\phi(x; b_2)$ in p_{μ^+} such that b_1, b_2 have the same syntactic type over a . Pick $d \models p_{\mu^+}$, then $\hat{\mathcal{C}} \models \phi[d; b_1] \wedge \neg\phi[d; b_2]$, $\text{gtp}(db_1/\emptyset) \neq \text{gtp}(db_2/\emptyset)$ (actually ϕ might tell the exact Galois type of db_1). On the other hand, let $r = \text{gtp}(b_1 a/\emptyset)$ and consider $R_r(x; a)$. As b_1, b_2 have the same syntactic type over A , $\hat{\mathcal{C}} \models R_r[b_1; a] \wedge R_r[b_2; a]$, which means $\text{gtp}(b_2 a/\emptyset) = r$. Thus there is $f : b_1 \cong_a b_2$. □

In the above proof, we did not use tameness simply because $(< \mu^+)$ -Galois Morleyization is already large enough to code all types over domains of size μ .

Corollary 7.5. *Let μ be an infinite cardinal and \mathbf{K} be μ -tame. Let $A \subseteq B$ with $|A| \leq \mu$. Then any $p \in S(B)$ ($\geq \mu$)-splits over A iff it μ -splits over A .*

Proof. We adopt the previous proof: suppose $d \models p$ ($\geq \mu$)-splits over A , obtain witness $f : B_1 \cong_A B_2$ as above. Enumerate $A = a$ and B_i as b_i . Since $f(p) \upharpoonright B_2 \neq p \upharpoonright B_2$, $\text{gtp}(f(d)b_2/\emptyset) \neq \text{gtp}(db_2/\emptyset)$. By Proposition 5.4, there is a quantifier-free formula ϕ in \hat{L}_{μ^+, μ^+} so that $\hat{\mathcal{C}} \models \phi[db_1] \wedge \phi[f(d)b_2] \wedge \neg\phi[db_2]$. b_1 and b_2 have the same syntactic type over a because of f . Therefore, $p_{\mu^+} = \text{tp}_{\mu^+}(d/B) \supseteq \text{tp}_{\mu^+}(d/B_1 \cup B_2)$ μ -syn-splits over A . This implies p μ -splits over A by the second paragraph of the previous proof. \square

We now prove a series of results syntactically. The original proofs in [Van06, Theorems I 4.10, 4.12], [GV06, Section 6] are semantic.

Lemma 7.6. *Let μ be an infinite cardinal and \mathbf{K} be μ -tame. For any $B \subseteq C$ both of size $\geq \mu$ and $p \in \text{gS}(C)$, we have $p_{\mu^+} \upharpoonright B = (p \upharpoonright B)_{\mu^+}$.*

Proof. Let $d \models p$,

$$\begin{aligned} p_{\mu^+} \upharpoonright B &= \text{tp}_{\mu^+}(d/C) \upharpoonright B \\ &= \text{tp}_{\mu^+}(d/B) \\ &= (p \upharpoonright B)_{\mu^+} \text{ because } d \models p \upharpoonright B \end{aligned}$$

\square

Lemma 7.7. *Let μ be an infinite cardinal, $A \subset_u B \subseteq C$ all of size μ and $p, q \in \text{gS}(C)$. Suppose p, q do not μ -split over A and $p \upharpoonright B = q \upharpoonright B$. Then $p = q$. We can allow $|B|, |C| \geq \mu$ if we assume μ -tameness.*


Proof. We comment the case $|C| > \mu$ in square brackets. Let $\hat{\mathbf{K}}$ be the ($< \mu^+$)-Galois Morleyization of \mathbf{K} . Since p, q do not μ -split over A , Proposition 7.4 shows that p_{μ^+}, q_{μ^+} do not μ -syn-split over A [use μ -tameness and Corollary 7.5]. Suppose $p_{\mu^+} \neq q_{\mu^+}$, [by μ -tameness] there is $d \subseteq C$ of size μ , $\phi(x; y)$ such that $\phi(x; d) \in p_{\mu^+} - q_{\mu^+}$. As $B \supset_u A$, we may pick $b \models \text{tp}_{\mu^+}(d/A)$. By non-syn-splitting, $\phi(x; b) \in p_{\mu^+} - q_{\mu^+}$ contradicting $p_{\mu^+} \upharpoonright B = q_{\mu^+} \upharpoonright B$ [use Proposition 5.4 and Lemma 7.6]. \square

Extension also holds but it is applicable to μ -sized models.

Lemma 7.8. *Let μ be an infinite cardinal, $A \subset_u B \subseteq C$ all of size μ . Let $p \in \text{gS}(B)$ do not μ -split over A . Then there is $q \in \text{gS}(C)$ extending p and does not μ -split over A . Also, if p is non-algebraic, we can have q non-algebraic.*

Proof. Let $\hat{\mathbf{K}}$ be the $(< \mu^+)$ -Galois Morleyization of \mathbf{K} . First we decide whether $\pm\phi(x; c) \in q_{\mu^+}$ for each quantifier-free formula in \hat{L}_{μ^+, μ^+} over C . Since $A \subset_u B$, there is a copy $C' \subset B$ of C . We want q does not μ -split over A , by Proposition 7.4 we must set $\phi(x; c) \in q_{\mu^+}$ iff $\phi(x; b_c) \in p_{\mu^+}$ where $b_c \in C'$ and $b_c \models \text{tp}_{\mu^+}(c/A)$. Such q_{μ^+} is realized because p_{μ^+} , and thus $p_{\mu^+} \upharpoonright C' \cong_A q_{\mu^+}$ is realized where $C' \subset B$ is the copy of C .

If p is non-algebraic and we can modify the argument above by extending C to a copy of B . Then p_{μ^+} is realized/algebraic iff q_{μ^+} is.



□

Lemma 7.9. *Let μ be an infinite cardinal, \mathbf{K} be μ -tame and stable in μ . For any $|A| \leq \mu$, $A \subset_u C$,*

$$\chi := |\{p \in \text{gS}(C) : p \text{ does not } \geq \mu\text{-split over } A\}| \leq \mu$$

Proof. Pick B of size μ with $A \subset_u B \subseteq C$. By Corollary 7.5, Lemma 7.7 and Lemma 7.8, $\chi = |\{p \in \text{gS}(B) : p \text{ does not } \mu\text{-split over } A\}| \leq |\text{gS}(B)| \leq \mu$. □

The following originates from [She99, Claim 3.3] and is extended to longer types in [GV06, Fact 4.6].

Lemma 7.10. *Let μ be an infinite cardinal and \mathbf{K} be stable in μ . For any $p \in \text{gS}(B)$, there is $A \subseteq B$ of size μ such that p does not μ -splits over A .*

Proof. Suppose the lemma is false, we can find $d \models p \in \text{gS}(B)$, B of size $> \mu$ such that p μ -splits over all A of size μ . Let $\hat{\mathbf{K}}$ be the $(< \mu^+)$ -Galois Morleyization of \mathbf{K} . By Proposition 7.4, p_{μ^+} μ -syn-splits over all A of size μ . Pick any $A_0 \subset B$ of size μ and choose minimum $\kappa \leq \mu$ with $2^\kappa > \mu$. By assumption we can build $\langle a_{\eta \frown 0}, a_{\eta \frown 1}, \phi_\eta, f_\eta : \eta \in 2^{< \kappa} - \{\langle \rangle\} \rangle$ and $\langle A_\alpha : \alpha < \kappa \rangle$ such that

1. $\langle A_\alpha : \alpha < \kappa \rangle$ is increasing and continuous. For $\alpha < \kappa$, $A_\alpha \subset B$ has size μ .
2. For $\eta \in 2^{< \kappa}$, $a_\eta \in A_{l(\eta)}$, $f_\eta : a_{\eta \frown 0} \mapsto a_{\eta \frown 1}$ and f_η fixes $A_{l(\eta)}$.
3. For $\eta \in 2^{< \kappa}$, $\phi_\eta(x; a_{\eta \frown 0}), \neg \phi_\eta(x; a_{\eta \frown 1}) \in p_{\mu^+}$

$$\begin{array}{ccc}
 A_2 & \phi_0(x; a_{00}) \xrightarrow[A_1]{f_0} \phi_0(x; a_{01}) & \phi_1(x; a_{10}) \xrightarrow[A_1]{f_1} \phi_1(x; a_{11}) \\
 A_1 & a_0 \xrightarrow[A_0]{f_\emptyset} a_1 &
 \end{array}$$

For $\nu \in 2^\kappa$, define $q_\nu := \{\phi_\eta(x; a_{\eta \frown i}) : \eta \in 2^{< \kappa}, \eta \frown i \sqsubseteq \nu; i = 1, 2\} \cup \{\neg \phi_\eta(x; a_{\eta \frown 1}) : \eta \in 2^{< \kappa}, \eta \frown 0 \sqsubseteq \nu\}$. $\langle q_\nu : \nu \in 2^\kappa \rangle$ are obviously pairwise contradictory. It remains to show each of them is realized. For any $\nu \in 2^\kappa$, define g_ν to be the composition of $\langle f_{\nu \upharpoonright \alpha}^{\nu \upharpoonright \alpha} : \alpha < \kappa \rangle$ where each $f_\eta^0 := \text{id}$ and $f_\eta^1 := f_\eta$. g_ν is well-defined because for $\eta \in 2^{< \kappa}$, f_η fixes $A_{l(\eta)}$. Extend g_ν to an isomorphism containing d . Inductively we can show $g_\nu(d) \models q_\nu$: let $\alpha < \kappa$. If $\nu \upharpoonright \alpha = 0$, then item (3) in the construction guarantees $g(p_\mu^+) \supseteq g_\nu(\{\phi_{\nu \upharpoonright \alpha}(x; a_{\nu \upharpoonright \alpha \frown 0}), \neg \phi_{\nu \upharpoonright \alpha}(x; a_{\nu \upharpoonright \alpha \frown 1})\}) = \{\phi_{\nu \upharpoonright \alpha}(x; a_{\nu \upharpoonright \alpha \frown 0}), \neg \phi_{\nu \upharpoonright \alpha}(x; a_{\nu \upharpoonright \alpha \frown 1})\}$. If $\nu \upharpoonright \alpha = 1$, then $g(p_\mu^+) \supseteq g_\nu(\{\phi_{\nu \upharpoonright \alpha}(x; a_{\nu \upharpoonright \alpha \frown 0})\}) = \{\phi_{\nu \upharpoonright \alpha}(x; a_{\nu \upharpoonright \alpha \frown 1})\}$. \square

Proof of Theorem 2.10. Let C be of size λ . By an exhaustive argument, we may extend C to be μ^+ -saturated. Let $A \subset C$ of size μ . By Proposition 7.3(1) and (2), we can build B of size μ such that $A \subset_u B \subset C$. Thus Lemma 7.9 applies. Also by Lemma 7.10, each $p \in \text{gS}(C)$ does not μ -split over some $A_p \subset C$ of size μ . There are at most λ^μ of such A_p . Thus

$$|\text{gS}(C)| = \left| \bigcup \{p \in \text{gS}(C) : p \text{ does not } \mu\text{-split over } A_p\} \right| \leq \lambda^\mu \cdot \mu = \lambda$$

\square

Corollary 7.11. *If \mathbf{K} is stable in some $\lambda < \text{LS}(\mathbf{K})$. Then the first stability cardinal $\geq \text{LS}(\mathbf{K})$ is bounded above by $2^{\text{LS}(\mathbf{K})}$.*

Proof. Apply Theorem 2.10 to $(2^{\text{LS}(\mathbf{K})})^\lambda = 2^{\text{LS}(\mathbf{K})}$. □

Our final application is the upward transfer of stability. The original proof of (1) below uses weak tameness (tameness over saturated models). [Vas16a, Lemma 5.5] proves a stronger version of (2) with chain local character instead of set local character, but we do not assume the former here.

Proposition 7.12. *Let $\mu < \lambda$ be infinite cardinals. Assume \mathbf{K} is μ -tame and stable in μ .*

1. [BKV06, Theorem 4.5] \mathbf{K} is also stable in μ^+ .
2. *If in addition $\text{cf}(\lambda) > \mu$ and \mathbf{K} is stable in unbounded many cardinals below λ , then it is stable in λ .*

Proof. 1. Suppose $|A| = \mu^+$ has μ^{++} many types $\langle p_\alpha : \alpha < \mu^{++} \rangle$. Write $A = \bigcup_{i < \mu^+} A_i$ increasing and continuous, with $|A_i| = \mu$. We may assume $A_{i+1} \models \text{gS}(A_i)$ by the following: define another chain $\langle A'_i : i \leq \mu^+ \rangle$ increasing and continuous such that $|A'_{i+1}| = |A_{i+1}| = \mu$ and $A'_{i+1} \supset_u A_i \cup A'_i$ (using Proposition 7.3(1)). Replace A by A'_{μ^+} .

By Lemma 7.10 and $\text{cf} \mu^{++} > \mu^+$, we may assume all p_α does not μ -split over A_0 . By stability in μ and pigeonhole principle, we may assume all p_α has the same type over A_1 . Together with Lemma 7.7, all p_α are equal, contradiction.

2. We consider limit cardinal λ . Pick a cofinal sequence $\langle \lambda_i : i < \text{cf}(\lambda) \rangle$ to λ such that \mathbf{K} is stable in all λ_i . Repeat the same argument as (1) but with $|A| = \lambda_i$ and $A = \bigcup_{i < \text{cf}(\lambda)} A_i$.

□

REFERENCES

- [Bal09] John T. Baldwin. *Categoricity*. University lecture series; v. 50. American Mathematical Society, Providence, R.I, 2009.

- [BGKV16] Will Boney, Rami Grossberg, Alexei Kolesnikov, and Sebastien Vasey. Canonical forking in AECs. *Annals of pure and applied logic*, 167(7):590–613, 2016.
- [BKV06] John Baldwin, David Kueker, and Monica VanDieren. Upward stability transfer for tame abstract elementary classes. *Notre Dame journal of formal logic*, 47(2):291–298, 2006.
- [Bon17] Will Boney. Computing the number of types of infinite length. *Notre Dame journal of formal logic*, 58(1), 2017.
- [GL02] Rami Grossberg and Olivier Lessmann. Shelah’s stability spectrum and homogeneity spectrum in finite diagrams. *Archive for mathematical logic*, 41(1):1–31, 2002.
- [Gro21] Rami Grossberg. *A Course in Model Theory I: Introduction*. 2021. A book in preparation.
- [GS86] Rami Grossberg and Saharon Shelah. On the number of nonisomorphic models of an infinitary theory which has the infinitary order property. I. *J. Symbolic Logic*, 51(2):302–322, 1986.
- [GS98] Rami Grossberg and Saharon Shelah. On Hanf numbers of the infinitary order property. 1998. Preprint.
- [GV06] Rami Grossberg and Monica VanDieren. Galois-stability for tame abstract elementary classes. *Journal of mathematical logic*, 6(1):25–48, 2006.
- [Jec03] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer, Berlin, Germany, 3rd edition, 2003.
- [Mar02] David Marker. *Model Theory : An Introduction*. Graduate Texts in Mathematics, 217. Springer New York, New York, NY, 1st edition, 2002.
- [Mor65] Michael Morley. Omitting classes of elements. In Addison, Henkin, and Tarski, editors, *The Theory of Models*, pages 265–273. North-Holland, Amsterdam, 1965.
- [She71] Saharon Shelah. Finite diagrams stable in power. *Ann. Math. Logic*, 2(1):69–118, 1970/1971.
- [She72] Saharon Shelah. A combinatorial problem; stability and order for models and theories in infinitary languages. *Pacific J. Math.*, 41:247–261, 1972.

- [She90] Saharon Shelah. *Classification theory and the number of non-isomorphic models*. Studies in logic and the foundations of mathematics; v. 92. North-Holland, Amsterdam, rev. edition, 1990.
- [She99] Saharon Shelah. Categoricity for abstract classes with amalgamation. *Ann. Pure Appl. Logic*, 98(1-3):261–294, 1999.
- [She09a] Saharon Shelah. *Classification theory for abstract elementary classes*, volume 18 of *Studies in Logic (London)*. College Publications, London, 2009.
- [She09b] Saharon Shelah. *Classification theory for abstract elementary classes. Vol. 2*, volume 20 of *Studies in Logic (London)*. College Publications, London, 2009.
- [Van06] Monica VanDieren. Categoricity in abstract elementary classes with no maximal models. *Annals of pure and applied logic*, 141(1):108–147, 2006.
- [Vas16a] Sebastien Vasey. Forking and superstability in tame AECs. *The Journal of symbolic logic*, 81(1):357–383, 2016.
- [Vas16b] Sebastien Vasey. Infinitary stability theory. *Archive for mathematical logic*, 55(3):567–592, 2016.

Email: wangchil@andrew.cmu.edu

URL: <http://www.math.cmu.edu/~wangchil/>

Address: Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh PA 15213, USA